

MATHEMATICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer:

1. The absolute difference of the coefficient of x^7 and

x^9 in the expansion of $\left(2x + \frac{1}{2x}\right)^{11}$ is

- (1) 11×2^5 (2) 11×2^7
 (3) 11×2^4 (4) 11×2^3

Answer (2)

Sol. $T_{r+1} = {}^{11}C_r (2x)^{11-r} \left(\frac{1}{2x}\right)^r$

$$= {}^{11}C_r \frac{2^{11-r}}{2^r} x^{11-2r}$$

$$11 - 2r = 7 \text{ and } 11 - 2r = 9$$

$$r = 2 \qquad \qquad \qquad r = 1$$

\therefore Coefficient of x^7 is ${}^{11}C_2 \frac{(2)^9}{2^2} = {}^{11}C_2 (2)^7$

Coefficient of x^9 is ${}^{11}C_1 \frac{(2)^{10}}{2} = {}^{11}C_1 (2)^9$

$${}^{11}C_2 (2)^7 - 11 \times (2)^9$$

$$= 11 \times 2^7$$

2. Let $f(x) = \{1, 2, 3, 4, 5, 6, 7\}$ the relation $R = \{(x, y) \in A \times A, x + y = 7\}$ is

- (1) Symmetric
 (2) Reflexive
 (3) Transitive
 (4) Equivalence

Answer (1)

Sol. $x + y = 7$

$y = 7 - x$

$R = \{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$

$\therefore (a, b) \in R \Rightarrow (b, a) \in R.$

\therefore Relation is symmetric

3. The number of words with or without meaning can be formed from the word MATHEMATICS where C, S not come together is

- (1) $\frac{9}{8} \times 10!$ (2) $\frac{1}{8} \times 10!$
 (3) $\frac{5}{8} \times 10!$ (4) $\frac{1}{2} \times 10!$

Answer (1)

Sol. Total words = $\frac{11!}{2!2!2!}$

When C and S are together = $\frac{10!}{2!2!2!} \times 2!$

$$\therefore \text{ Required number of words} = \frac{11!}{(2!)^3} - \frac{10!}{(2!)^3} \times 2!$$

$$= \frac{10!}{8} [11 - 2]$$

$$= \frac{9}{8} \times 10!$$

4. Let $a_n = 5 + 8 + 14 + 23 + \dots$ upto n terms. If

$S_n = \sum_{k=1}^n a_k$, then $S_{30} - a_{40}$ is equal to

- (1) 78025
 (2) 12800
 (3) 11600
 (4) 12100

Answer (1)

Sol. $a_n = 5 + 8 + 14 + \dots T_n$

$$\frac{a_n}{0} = \frac{5 + 8 + 14 + \dots + T_{n-1} + T_n}{5 + \underbrace{3 + 6 + 9 + \dots}_{(n-1) \text{ terms}} - T_n}$$

$$\Rightarrow T_n = 5 + \left(\frac{n-1}{2}\right)(6 + (n-2)3) = 5 + \frac{3}{2}(n-1)^n$$

$$5 + \frac{3}{2}n^2 - \frac{3}{2}n$$

$$\Rightarrow \frac{1}{2}(10 + 3n^2 - 3n)$$

$$\therefore T_n = \frac{1}{2}(3n^2 - 3n + 10)$$

$$a_n = \sum T_n = \frac{1}{2} \left[\frac{3 \cdot (n)(n+1)(2n+1)}{6} - \frac{3 \cdot (n)(n+1)}{2} + 10n \right]$$

$$= \frac{1}{2} (n) \left(\frac{(n+1)(2n+1)}{2} - \frac{3(n+1)}{2} + 10 \right)$$

$$a_n = \frac{n}{4} (2n^2 + 3n + 1 - 3n - 3 + 20)$$

$$= \frac{n}{4} (2n^2 + 18) = \frac{n}{4} (n^2 + 9)$$

$$a_{40} = \frac{40}{2} (1600 + 9) = 1609 \times 20 = 32180$$

$$S_n = \sum a_n = \frac{1}{2} \left(\left(\frac{(n)(n+1)}{2} \right)^2 + \frac{9 \cdot (n)(n+1)}{2} \right)$$

$$S_{30} = \frac{1}{2} \left(\left(\frac{30 \times 31}{2} \right)^2 + \frac{9}{2} (30)(31) \right)$$

$$= \frac{1}{2} (216225 + 4185)$$

$$= 110205$$

$$S_{30} - a_{40} = 78025$$

5. The equation $ax^2 + bx + c = 0$ has roots α and β .

Then find $\lim_{x \rightarrow \frac{1}{\alpha}} \frac{1 - \cos(cx^2 + bx + a)}{2(1 - \alpha x)^2}$ is

- | | |
|--|--|
| (1) $\frac{c^2(\alpha - \beta)^2}{4\alpha^4\beta^2}$ | (2) $\frac{c^2(\alpha - \beta)^2}{\alpha^4\beta^2}$ |
| (3) $\frac{c^2(\alpha - \beta)^2}{2\alpha^4\beta^2}$ | (4) $\frac{c^2(\alpha - \beta)^2}{4\alpha^2\beta^2}$ |

Answer (1)

Sol. $\lim_{x \rightarrow \frac{1}{\alpha}} \frac{2 \sin^2 \left(\frac{cx^2 + bx + a}{2} \right)}{2\alpha^2 \left(x - \frac{1}{\alpha} \right)^2}$

$$= \frac{c^2(\alpha - \beta)^2}{4\alpha^2\beta^2}$$

6. $\theta \in (0, 2\pi)$ and $\frac{1 + 2i \sin \theta}{1 - i \sin \theta}$ is purely imaginary then

the value of θ is

- | | |
|------------|---------------------|
| (1) π | (2) 0 |
| (3) 2π | (4) $\frac{\pi}{4}$ |

Answer (4)

Sol. Real part has to be zero

$$\Rightarrow \frac{1 - 2 \sin^2 \theta}{1 + \sin^2 \theta} = 0$$

$$\sin^2 \theta = \frac{1}{2}$$

$$\theta = n\pi \pm \frac{\pi}{4}, n \in I$$

$$\theta = \frac{\pi}{4}, \frac{5\pi}{4}, \frac{3\pi}{4}$$

7. The statement $(p \wedge (\sim q)) \vee (\sim p)$ is equivalent to

- | | |
|------------------|----------------------------|
| (1) $p \wedge q$ | (2) $\sim p \vee \sim q$ |
| (3) $p \vee q$ | (4) $\sim p \wedge \sim q$ |

Answer (2)

Sol. $(p \wedge (\sim q)) \vee (\sim p)$

$$= (p \vee \sim p) \wedge (\sim q \vee \sim p)$$


$$= T \wedge (\sim q \vee \sim p)$$

$$= \sim q \vee \sim p$$

8. From $O(0, 0)$, two tangents OA and OB are drawn to a circle $x^2 + y^2 - 6x + 4y + 8 = 0$, then the equation of circumcircle of $\triangle OAB$.

- | | |
|-------------------------------|-------------------------------|
| (1) $x^2 + y^2 - 3x + 2y = 0$ | (2) $x^2 + y^2 + 3x - 2y = 0$ |
| (3) $x^2 + y^2 + 3x + 2y = 0$ | (4) $x^2 + y^2 - 3x - 2y = 0$ |

Answer (1)



Sol. $(0, 0)$

$(0, 0)$ and $(3, -2)$ are diametric end points

$$\therefore (x - 0)(x - 3) + (y - 0)(y + 2) = 0$$

$$\boxed{x^2 + y^2 - 3x + 2y = 0}$$


9. The mid points of side of a triangle are $(0, 1), (1, 0), (1, 1)$, where incentre is D . A parabola $y^2 = 4ax$ passes through D whose focus is $(\alpha + \beta\sqrt{2}, 0)$ then

$$\frac{\beta^2}{\alpha}$$

- | | |
|-------------------|---------|
| (1) $\frac{1}{2}$ | (2) 2 |
| (3) $\frac{1}{8}$ | (4) 4 |

Answer (3)

Sol.



\therefore Mid-point is $(0, 1)$, $(1, 0)$ and $(1, 1)$

$$I = \left(\frac{4}{4+2\sqrt{2}}, \frac{4}{4+2\sqrt{2}} \right)$$

$$y^2 = 4ax$$

$\therefore y^2 = 4ax$ passes through I

$$\left(\frac{4}{4+2\sqrt{2}} \right)^2 = 4a \left(\frac{4}{4+2\sqrt{2}} \right) \Rightarrow a = \frac{1}{4+2\sqrt{2}}$$

Focus = $(a, 0)$

$$= \left(\frac{1}{4+2\sqrt{2}}, 0 \right)$$

$$= \left(\frac{4-2\sqrt{2}}{8}, 0 \right)$$

$$\therefore \alpha = \frac{4}{8} = \frac{1}{2}, \beta = \frac{-2}{8} = \frac{-1}{4}$$

$$\frac{\beta^2}{\alpha} = \frac{1}{8}$$

10. Let $R = \{a, b, c, d, e\}$ and $S = \{1, 2, 3, 4\}$. Then number of onto functions $f(x) : R \rightarrow S$ such that $f(a) \neq 1$ is

- (1) 240 (2) 180
(3) 204 (4) 216

Answer (2)

Sol. Total number of onto functions

$$= \frac{5!}{3!2!} \times 4!$$

Now, when $f(a) = 1$

$$\frac{4!}{2!2!} \times 3! + 4!$$

$$\therefore \text{Required functions} = 240 - 36 - 24 = 180$$

11. A parabola with focus $(3, 0)$ and directrix $x = -3$. Points P and Q lie on the parabola and their ordinates are in the ratio 3 : 1. The point of intersection of tangents drawn at points P and Q lies on the parabola

- (1) $y^2 = 16x$ (2) $y^2 = 4x$
(3) $y^2 = 8x$ (4) $x^2 = 4y$

Answer (1)

Sol. Given parabola $y^2 = 12x$

$$P(3t_1^2, 6t_1), Q(3t_2^2, 6t_2)$$

$$\frac{t_1}{t_2} = 3 \Rightarrow t_1 = 3t_2 \quad \dots(i)$$

Let point of intersection be (h, k)

$$h = 3t_1t_2 \quad \dots(ii)$$

$$\text{and } k = 3(t_1 + t_2) \quad \dots(iii)$$

$$(i) \text{ and } (iii) \Rightarrow t_2 = \frac{k}{12}$$

$$(ii) \Rightarrow h = 9t_2^2 = 9 \times \frac{k^2}{144} \Rightarrow k^2 = 16h$$

$$\Rightarrow y^2 = 16x$$

12. In probability distribution for discrete variable $x = 0, 1, 2 \dots P(x = x) = k(x + 1) \cdot 3^{-x}$. The probability of $P(x \geq 2)$ is equal to

(1) $\frac{5}{18}$ (2) $\frac{10}{18}$

(3) $\frac{20}{27}$ (4) $\frac{7}{27}$

Answer (4)

Sol. $\Sigma P = 1$

$$\Rightarrow k(1 + 2 \cdot 3^{-1} + 3 \cdot 3^{-2} + \dots) = 1$$

$$\text{Let } S = 1 + \frac{2}{3} + \frac{3}{3^2} + \dots$$

$$\frac{S}{3} = \frac{1}{3} + \frac{2}{3^2} + \dots$$

$$\frac{2S}{3} = 1 + \frac{1}{3} + \frac{1}{3^2} + \dots = \frac{1}{1 - \frac{1}{3}} = \frac{3}{2}$$

$$\Rightarrow S = \frac{9}{4}$$

$$\therefore k \cdot \frac{9}{4} = 1 \Rightarrow k = \frac{4}{9}$$

Now $P(x \geq 2) = 1 - P(x = 0, 1)$

$$= 1 - \left(k + k \cdot \frac{2}{3} \right)$$

$$= 1 - \frac{5k}{3}$$

$$= 1 - \frac{5}{3} \cdot \frac{4}{9}$$

$$= \frac{7}{27}$$

13. If $f(x) = \begin{cases} 3x^2 + k\sqrt{x+1} & 0 < x < 1 \\ 3mx^2 + k^2 & x \geq 1 \end{cases}$ is

differentiable at $x > 1$ then $\frac{8f'(8)}{f'\left(\frac{1}{8}\right)}$ is for $k \neq 0$

(1) 309

(2) 311

(3) 306

(4) 305

Answer (1)

Sol. $f(x) = \begin{cases} 3x^2 + k\sqrt{x+1} & 0 < x < 1 \\ 3mx^2 + k^2 & x \geq 1 \end{cases}$

$$3 + k\sqrt{2} = 3m + k^2 \quad \dots(i)$$

$$f'(x) = \begin{cases} 6x + \frac{k}{2\sqrt{x+1}} & 0 < x < 1 \\ 6mx & x \geq 1 \end{cases}$$

$$6 + \frac{k}{2\sqrt{2}} = 6m \quad \dots(ii)$$

$$3 + k\sqrt{2} = 3 + \frac{k}{4\sqrt{2}} + k^2$$

$$k = 0 \text{ or } \frac{7\sqrt{2}}{8}$$

If $k = 0$

If $k = \frac{7\sqrt{2}}{8}$

$m = 1$

$m = \frac{103}{96}$

(Rejected)

Now, $\frac{8f'(8)}{f'\left(\frac{1}{8}\right)} = \frac{48m}{\frac{6}{8} + \frac{k}{2\sqrt{\frac{9}{8}}}} = \frac{48m}{\frac{6}{8} + \frac{\sqrt{2}k}{3}}$

$$\frac{8f'(8)}{f'\left(\frac{1}{8}\right)} = 309$$

14.

15.

16.

17.

18.

19.

20.

SECTION - B

Numerical Value Type Questions: This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g., 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.

21. The area of quadrilateral having vertices as (1, 2), (5, 6), (7, 6), (-1, -6)

Answer (24)

Sol. Area = $\frac{1}{2} \begin{vmatrix} 1 & 2 \\ 5 & 6 \\ 7 & 6 \\ -1 & -6 \\ 1 & 2 \end{vmatrix}$

$$= \frac{1}{2} [6 + 30 - 42 - 2 - 10 - 42 + 6 + 6]$$

$$= \frac{1}{2} [48] = 24$$

22. The value of $\int_0^{2.4} [x^2] dx$ is $\alpha + \beta\sqrt{2} + \gamma\sqrt{3} + \delta\sqrt{5}$

then $(a + b + c + d + e)$ is equal to

Answer (06)

Sol. $\int_0^{2.4} [x^2] dx = \int_0^1 0 dx + \int_1^{\sqrt{2}} 1 dx + \int_{\sqrt{2}}^{\sqrt{3}} 2 dx + \int_{\sqrt{3}}^{\sqrt{4}} 3 dx + \int_{\sqrt{4}}^{\sqrt{5}} 4 dx + \int_{\sqrt{5}}^{2.4} 5 dx$

$$= (\sqrt{2} - 1) + 2(\sqrt{3} - \sqrt{2}) + 3(\sqrt{4} - \sqrt{3}) + 4(\sqrt{5} - \sqrt{4}) + 5(2.4 - \sqrt{5})$$

$$= 9 - \sqrt{2} - \sqrt{3} - \sqrt{5}$$

$\therefore \alpha = 9, \beta = -1, \gamma = -1, \delta = -1$

$\therefore \alpha + \beta + \gamma + \delta = 6$

23. $\frac{dx}{dy} - \frac{3\sin y}{\cos y(\ln \cos y)} x = \frac{\sin y}{(\ln \cos y)^2 \cos y}$ and

$x\left(\frac{\pi}{3}\right) = \frac{1}{2\ln 2}, x\left(\frac{\pi}{6}\right) = \frac{1}{\ln(m) - \ln(n)}$ then the value

of mn is

Answer (12)

Sol. $I = e^{\int \frac{-3\sin y}{\cos y(\ln \cos y)} dy}$

Put $\ln(\cos y) = t$

$\frac{-1}{\cos y} \sin y dy = dt$

$= e^{\int \frac{3}{t} dt}$

$= (\ln \cos y)^3$

$x(\ln \cos y)^3 = \int \frac{\sin y}{\cos y} \ln \cos y dy$

$x(\ln \cos y)^3 = \frac{-(\ln(\cos y))^2}{2} + C$

$x\left(\frac{\pi}{3}\right) = \frac{1}{2\ln 2}$

$\Rightarrow C = 0$

$\therefore x = -\frac{1}{2\ln(\cos y)}$

$x\left(\frac{\pi}{6}\right) = \frac{1}{\ln 4 - \ln 3}$

$m = 4$

$n = 3$

24. If m is the number of solution of $x^2 - 12x + 31 + [x] = 0$ and n be the number of solution of $x^2 - 5|x+2| - 4 = 0$, then the value of $m^2 + mn + n^2$ is


Answer (19)

Sol. $x^2 - 12x + 31 - [x] = 0$

$x^2 - 12x + 31 = [x]$

$(x-6)^2 - 5 = [x]$


So, by graph



\therefore Two points of intersects

$\therefore m = 2$

$x^2 - 5|x-2| - 4 = 0$



$\therefore n = 3$

$m^2 + mn + n^2 = 4 + 6 + 9 = 19$

25.
 26.
 27.
 28.
 29.
 30.

