

08/04/2025

Evening



Aakash

Medical | IIT-JEE | Foundations

Corporate Office : AESL, 3rd Floor, Incuspaze Campus-2, Plot-13, Sector-18, Udyog Vihar,
Gurugram, Haryana-122015

Answers & Solutions

Time : 3 hrs.

for

M.M. : 300

JEE (Main)-2025 (Online) Phase-2

(Mathematics, Physics and Chemistry)

IMPORTANT INSTRUCTIONS:

- (1) The test is of **3 hours** duration.
- (2) This test paper consists of **75** questions. Each subject (MPC) has 25 questions. The maximum marks are 300.
- (3) This question paper contains **Three Parts**. **Part-A** is Mathematics, **Part-B** is Physics and **Part-C** is Chemistry. Each part has only two sections: **Section-A** and **Section-B**.
- (4) **Section - A** : Attempt all questions.
- (5) **Section - B** : Attempt all questions.
- (6) **Section - A (01 - 20)** contains 20 multiple choice questions which have **only one correct answer**. Each question carries **+4 marks** for correct answer and **-1 mark** for wrong answer.
- (7) **Section - B (21 - 25)** contains 5 **Numerical value** based questions (MCQ). The answer to each question should be rounded-off to the **nearest integer**. Each question carries **+4 marks** for correct answer and **-1 mark** for wrong answer.

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
IN PHYSICS
(MPC)

1000+ 99
PERCENTILERS
IN MATHS

4000+ 95
PERCENTILERS
IN CHEMISTRY

100
Percentile
in
Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in
Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in
Physics
&
Chemistry



99.99 **Devya Rustagi**
PSID: 00014768785

99.99
Percentile



99.99 **Amogh Bansal**
PSID: 00014769016

OUR JEE CHAMPIONS



Chirag Falor
4 Year Classroom
1 AIR
JEE (Adv.)
2020



Tanishka Kabra
4 Year Classroom
1 AIR-16 CRL
JEE (Adv.)
2022
ALL
INDIA
FEMALE
TOPPER



Sanvi Jain
4 Year Classroom
1 AIR-34 CRL
JEE (Main)
2024
ALL
INDIA
FEMALE
TOPPER

MATHEMATICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer:

1. Let the ellipse $3x^2 + py^2 = 4$ pass through the centre C of the circle $x^2 + y^2 - 2x - 4y - 11 = 0$ of radius r . Let f_1, f_2 be the focal distances of the point C on the ellipse. Then $6f_1f_2 - r$ is equal to

- (1) 68
- (2) 74
- (3) 70
- (4) 78

Answer (3)

Sol. S: $x^2 + y^2 - 2x - 4y - 11 = 0$

Centre C(1, 2)

radius = $\sqrt{1+4+11}$

= 4

Ellipse $3x^2 + py^2 = 4$ passes through (1, 2)

$3(1) + p(4) = 4$

$4p = 1$

$p = \frac{1}{4}$

E: $3x^2 + \frac{y^2}{4} = 4$

or $\frac{x^2}{4} + \frac{y^2}{16} = 1$

$e = \sqrt{1 - \frac{4}{16}}$

$e = \sqrt{1 - \frac{1}{12}}$

$e = \sqrt{\frac{11}{12}}$

Focus = $(0, \pm 2\sqrt{\frac{11}{3}})$

$f_1 = \sqrt{1 + \left(2 - 2\sqrt{\frac{11}{3}}\right)^2}$ and $f_2 = \sqrt{1 + \left(2 + 2\sqrt{\frac{11}{3}}\right)^2}$

$f_1f_2 = \frac{37}{3}$

$\Rightarrow 6f_1f_2 - r = 74 - 4$

= 70

2. The integral $\int_{-1}^{\frac{3}{2}} \left(\left| \pi^2 x \sin(\pi x) \right| \right) dx$ is equal to:

- (1) $3 + 2\pi$
- (2) $2 + 3\pi$
- (3) $4 + \pi$
- (4) $1 + 3\pi$

Answer (4)

Sol. $I = \int_{-1}^{\frac{3}{2}} \left| \pi^2 x \sin(\pi x) \right| dx$

$= \int_{-1}^1 \left| \pi^2 x \sin(\pi x) \right| dx + \int_1^{\frac{3}{2}} \left| \pi^2 x \sin(\pi x) \right| dx$

$= 2 \int_0^1 \left| \pi^2 x \sin(\pi x) \right| dx - \pi^2 \int_1^{\frac{3}{2}} x \sin(\pi x) dx$

$= 2\pi^2 \int_0^1 x \sin(\pi x) dx - \pi^2 \int_1^{\frac{3}{2}} x \sin(\pi x) dx$

$\therefore \int x \sin(\pi x) dx = x \left(\frac{-\cos \pi x}{\pi} \right) - \int \frac{-\cos \pi x}{\pi} dx$

$= -\frac{x}{\pi} \cos \pi x + \frac{1}{\pi^2} \sin \pi x + C$

$\therefore I = 2\pi^2 \left(\frac{1}{\pi} \right) - \pi^2 \left(-\frac{1}{\pi^2} - \frac{1}{\pi} \right)$

= $2\pi + 1 + \pi$

= $3\pi + 1$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths



Shreyas Lohiya
PSID: 00003389699

100 Percentile in Physics



Harsh Jha
PSID: 00014863322

100 Percentile in Physics & Chemistry



Devya Rustagi
PSID: 00014768785

99.99 Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS

Chirag Falor
4 Year Classroom
1 AIR JEE (Adv.) 2020

Tanishka Kabra
4 Year Classroom
1 AIR-16 CRL JEE (Adv.) 2022

Sanvi Jain
4 Year Classroom
1 AIR-34 CRL JEE (Main) 2024

3. The value of

$$\cot^{-1}\left(\frac{\sqrt{1+\tan^2(2)}-1}{\tan(2)}\right) - \cot^{-1}\left(\frac{\sqrt{1+\tan^2\left(\frac{1}{2}\right)+1}}{\tan\left(\frac{1}{2}\right)}\right)$$

is equal to

- (1) $\pi - \frac{3}{2}$ (2) $\pi - \frac{5}{4}$
 (3) $\pi + \frac{3}{2}$ (4) $\pi + \frac{5}{2}$

Answer (2)

Sol.
$$\cot^{-1}\left(\frac{\sqrt{1+\tan^2 2}-1}{\tan 2}\right) - \cot^{-1}\left(\frac{\sqrt{1+\tan^2\left(\frac{1}{2}\right)+1}}{\tan\frac{1}{2}}\right)$$

$$= \cot^{-1}\left(\frac{|\sec 2|-1}{\tan 2}\right) - \cot^{-1}\left(\frac{\sec\left(\frac{1}{2}\right)+1}{\tan\frac{1}{2}}\right)$$

$$= \cot^{-1}\left(\frac{-\sec 2-1}{\tan 2}\right) - \cot^{-1}\left(\frac{\sec\frac{1}{2}+1}{\tan\frac{1}{2}}\right)$$

$$= \pi - \cot^{-1}\left(\frac{1+\cos 2}{\sin 2}\right) - \cot^{-1}\left(\frac{1+\cos\frac{1}{2}}{\sin\frac{1}{2}}\right)$$

$$= \pi - \cot^{-1}\left(\frac{2\cos^2 1}{2\sin 1 \cdot \cos 1}\right) - \cot^{-1}\left(\frac{2\cos^2\frac{1}{4}}{2\sin\frac{1}{4} \cdot \cos\frac{1}{4}}\right)$$

$$= \pi - \cot^{-1}(\cot 1) - \cot^{-1}\left(\cot\frac{1}{4}\right)$$

$$= \pi - 1 - \frac{1}{4}$$


$$= \pi - \frac{5}{4}$$

4. A line passing through the point $P(a, 0)$ makes an acute angle α with the positive x -axis. Let this line be rotated about the point P through an angle $\frac{\alpha}{2}$ in the clock-wise direction. If in the new position, the slope of the line is $2 - \sqrt{3}$ and its distance from the origin is $\frac{1}{\sqrt{2}}$, then the value of $3a^2 \tan^2 \alpha - 2\sqrt{3}$ is

- (1) 8 (2) 6
 (3) 5 (4) 4

Answer (4)

Sol.



$$\tan\frac{\alpha}{2} = 2 - \sqrt{3}$$

$$\Rightarrow \tan\alpha = \frac{1}{\sqrt{3}}$$

Equation of new line: $(y - 0) = (2 - \sqrt{3})(x - a)$
 $y = (2 - \sqrt{3})x - (2 - \sqrt{3})a$

Distance from origin = $\frac{1}{\sqrt{2}}$

$$\left| \frac{-(2 - \sqrt{3})a}{4 + 3 - 4\sqrt{3} + 1} \right| = \frac{1}{\sqrt{2}}$$

$$|a| = \frac{\sqrt{8 - 4\sqrt{3}}}{\sqrt{2}(2 - \sqrt{3})}$$

$$|a| = \frac{2\sqrt{2 - \sqrt{3}}}{\sqrt{2}(2 - \sqrt{3})}$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS

1000+ 99
PERCENTILERS

4000+ 95
PERCENTILERS

100
Percentile



Shreyas Lohiya
PSID: 00003389699

100
Percentile



Harsh Jha
PSID: 00014863322

100
Percentile



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



Chirag Falor
4 Year Classroom
1 AIR
JEE (Adv.)
2020



Tanishka Kabra
4 Year Classroom
1 AIR-16 CRL
JEE (Adv.)
2022



Sanvi Jain
4 Year Classroom
1 AIR-34 CRL
JEE (Main)
2024

$$|a| = \frac{\sqrt{2}}{\sqrt{2-\sqrt{3}}}$$

$$a^2 = \frac{2}{2-\sqrt{3}} = 2(2+\sqrt{3})$$

$$3a^2 \tan^2 \alpha - 2\sqrt{3} = 3(4+2\sqrt{3}) \times \frac{1}{3} - 2\sqrt{3}$$

$$= 4$$

5. Let $A = \begin{bmatrix} 2 & 2+p & 2+p+q \\ 4 & 6+2p & 8+3p+2q \\ 6 & 12+3p & 20+6p+3q \end{bmatrix}$.

If $\det(\text{adj}(\text{adj}(3A))) = 2^m \cdot 3^n$, $m, n \in \mathbb{N}$, then $m+n$ is equal to

(1) 20 (2) 26

(3) 22 (4) 24

Answer (4)

Sol. $\begin{vmatrix} 2 & 2+p & 2+p+q \\ 4 & 6+2p & 8+3p+2q \\ 6 & 12+3p & 20+6p+3q \end{vmatrix}$

$$= \begin{vmatrix} 2 & 2 & 2+p+q \\ 4 & 6 & 8+3p+2q \\ 6 & 12 & 20+6p+3q \end{vmatrix} + \underbrace{\begin{vmatrix} 2 & p & 2+p+q \\ 4 & 2p & 8+3p+2q \\ 6 & 3p & 20+6p+3q \end{vmatrix}}_{=0}$$

$$= 2 \times 2 \begin{vmatrix} 1 & 1 & 2+p+q \\ 2 & 3 & 8+3p+2q \\ 3 & 6 & 20+6p+3q \end{vmatrix}$$

$$C_3 \rightarrow C_3 \rightarrow pC_2$$

$$= 4 \begin{vmatrix} 1 & 1 & 2+q \\ 2 & 3 & 8+2q \\ 3 & 6 & 20+3q \end{vmatrix} = 4 \begin{vmatrix} 1 & 1 & 2 \\ 2 & 3 & 8 \\ 3 & 6 & 20 \end{vmatrix} + 0$$

$$= 4 \begin{vmatrix} 1 & 1 & 1 \\ 2 & 3 & 4 \\ 3 & 6 & 10 \end{vmatrix}$$

$$= 8(1(6) - 1(8) + 1(3))$$

$$= 8$$

$$|\text{adj}(\text{adj}(3A))| = (|3A|)^{2^2} = |3A|^4$$

$$= (3^3 |A|)^4 = 3^{12} \cdot |A|^4$$

$$= 3^{12} \cdot (2^3)^4$$

$$= 3^{12} \cdot 2^{12}$$

6. Given below are two statements:

Statement I: $\lim_{x \rightarrow 0} \left(\frac{\tan^{-1} x + \log_e \sqrt{\frac{1+x}{1-x}} - 2x}{x^5} \right) = \frac{2}{5}$

Statement II: $\lim_{x \rightarrow 1} \left(\frac{2}{x^{1-x}} \right) = \frac{1}{e^2}$

In the light of the above statements, choose the **correct** answer from the options given below

- (1) Statement I is false but Statement II is true
- (2) Both Statement I and Statement II are false
- (3) Both Statement I and Statement II are true
- (4) Statement I is true but Statement II is false

Answer (3)

Sol. $\lim_{x \rightarrow 0} \left(\frac{\tan^{-1} x + \ln \sqrt{\frac{1+x}{1-x}} - 2x}{x^5} \right)$

$$\left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right) + \frac{1}{2} \left(x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \frac{x^5}{5} + \dots \right) - 2x$$

$$\Rightarrow \lim_{x \rightarrow 0} \frac{-\frac{1}{2} \left(x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \frac{x^5}{5} + \dots \right) - 2x}{x^5}$$

$$\left(x + \frac{1}{2}(x+x) - 2x \right) + x^3 \left(\frac{-1}{3} + \frac{1}{2} \times \frac{1}{3} + \frac{1}{2} \times \frac{1}{3} \right)$$

$$+ x^5 \left(\frac{1}{5} + \frac{1}{2} \times \frac{1}{5} + \frac{1}{2} \times \frac{1}{5} \right) + \dots$$

$$\Rightarrow \lim_{x \rightarrow 0} \frac{\left(\frac{1}{5} + \frac{1}{10} + \frac{1}{10} \right) x^5}{x^5} = \frac{2}{5}$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
in Physics

1000+ 99
PERCENTILERS
in Chemistry

4000+ 95
PERCENTILERS
in Biology

100
Percentile
in Physics
& Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in Physics
& Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



$$\Rightarrow \lim_{x \rightarrow 1} x^{\left(\frac{2}{1-x}\right)} = \lim_{x \rightarrow 1} \left([1 + (x-1)]^{\frac{1}{x-1}} \right)^{(x-1)2}$$

$$= e^{-2} = \frac{1}{e^2}$$

7. Let $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$ and $\vec{b} = 2\hat{i} + \hat{j} - \hat{k}$. Let \hat{c} be a unit vector in the plane of the vector \vec{a} and \vec{b} and be perpendicular to \vec{a} . Then such a vector \hat{c} is:

- (1) $\frac{1}{\sqrt{3}}(\hat{i} - \hat{j} + \hat{k})$
- (2) $\frac{1}{\sqrt{5}}(\hat{j} - 2\hat{k})$
- (3) $\frac{1}{\sqrt{2}}(-\hat{i} + \hat{k})$
- (4) $\frac{1}{\sqrt{3}}(-\hat{i} + \hat{j} - \hat{k})$

Answer (3)

Sol. $\vec{c} = x\vec{a} + y\vec{b}$

$$\vec{c} = x(\hat{i} + 2\hat{j} + \hat{k}) + y(2\hat{i} + \hat{j} - \hat{k})$$

$$\vec{a} \cdot \vec{c} = (\hat{i} + 2\hat{j} + \hat{k}) \cdot (x(\hat{i} + 2\hat{j} + \hat{k}) + y(2\hat{i} + \hat{j} - \hat{k}))$$

$$(\hat{i} + 2\hat{j} + \hat{k}) \cdot (x\hat{i} + 2x\hat{j} + x\hat{k} + 2y\hat{i} + y\hat{j} - y\hat{k}) = 0$$

$$\Rightarrow (x + 2y) + 2(x + 9) + (x - y) = 0$$

$$\Rightarrow y = -2x$$

$$\therefore \vec{c} = x(-3\hat{i} + 3\hat{k})$$

$$|\vec{c}| = |x| \sqrt{9+9} = 3|x| \sqrt{2}$$

$$\therefore |\vec{c}| = 1$$

$$3|x| \sqrt{2} = 1$$

$$|x| = \frac{1}{3\sqrt{2}}$$

$$\text{Let } x = \frac{1}{3\sqrt{2}}$$

$$\vec{c} = \frac{1}{3\sqrt{2}}(-3\hat{i} + 3\hat{k})$$

$$\text{or } \vec{c} = \frac{1}{\sqrt{2}}(-\hat{i} + \hat{k})$$

8. Let the value of λ for which the shortest distance between the lines $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ and

$$\frac{x-\lambda}{3} = \frac{y-4}{4} = \frac{z-5}{5} \text{ is } \frac{1}{\sqrt{6}} \text{ be } \lambda_1 \text{ and } \lambda_2. \text{ Then the}$$

radius of the circle passing through the points $(0, 0)$, (λ_1, λ_2) and (λ_2, λ_1) is

- (1) $\frac{\sqrt{2}}{3}$
- (2) $\frac{5\sqrt{2}}{3}$
- (3) 3
- (4) 4

Answer (2)

Sol. $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4} \dots(1)$

$$\frac{x-\lambda}{3} = \frac{y-4}{4} = \frac{z-5}{5} \dots(2)$$

$$\vec{n}_1 \times \vec{n}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{vmatrix}$$

$$= \hat{i}(15 - 16) - \hat{j}(10 - 12) + \hat{k}(8 - 9)$$

$$= -\hat{i} + 2\hat{j} - \hat{k}$$

L_1 passing through $(1, 2, 3)$ and L_2 through $(\lambda, 4, 5)$

$$d = \frac{1}{\sqrt{6}}$$

$$\Rightarrow \frac{|(\lambda - 1)(-1) - 2(-2) + 2(-1)|}{\sqrt{1^2 + 4^2 + 1}} = \frac{1}{\sqrt{6}}$$

$$|-\lambda + 1 + 4 - 2| = 1$$

$$|-\lambda + 3| = 1$$

$$\lambda - 3 = \pm 1$$

$$\lambda = 4, 2$$

Circle passing through $(0, 0)$, $(1, 4)$, $(4, 1)$

$$\therefore \text{Area} = \frac{1}{2} \begin{vmatrix} 0 & 0 & 1 \\ 1 & 4 & 1 \\ 4 & 1 & 1 \end{vmatrix}$$

$$= \left| \frac{1}{2} (1 - 16) \right| = \frac{15}{2}$$

$$\therefore r = \frac{abc}{4\Delta}$$

$$= \frac{5\sqrt{2}}{3}$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
in Physics

1000+ 99
PERCENTILERS
in Maths

4000+ 95
PERCENTILERS
in Biology

100
Percentile
in Physics
& Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in Physics
& Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



9. Let $A = \{0, 1, 2, 3, 4, 5\}$. Let R be a relation on A defined by $(x, y) \in R$ if and only if $\max\{x, y\} \in \{3, 4\}$. Then among the statements

(S_1): The number of elements in R is 18, and
(S_2): The relation R is symmetric but neither reflexive nor transitive.

- (1) only (S_2) is true (2) both are false
(3) only (S_1) is true (4) both are true

Answer (1)

Sol. Let's write the pairs (x, y) in R

$(0, 3), (1, 3), (2, 3), (3, 3), (4, 3), (5, 3), (3, 0), (3, 1), (3, 2), (3, 4), (3, 5), (0, 4), (1, 4), (2, 4), (4, 4), (5, 4), (4, 0), (4, 1), (4, 2), (4, 5)$

There are total 20 pairs

If $(x, y) \in R$, then $\max(x, y) \in \{3, 4\}$

This means, $\max(y, x) \in \{3, 4\}$. So, $(y, x) \in R$.

Thus R is symmetric

Since $\max(5, 5) = 5 \notin \{3, 4\}$, $(5, 5) \notin R$

Thus R is not reflexive

$(3, 4) \in R$ & $(4, 2) \in R$, but $(3, 2) \notin R$, $2 \notin \{3, 4\}$

Thus, R is not transitive

Therefore, only S_2 is true

10. If $\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \dots = \frac{\pi^4}{90}$,

$$\frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots = \alpha$$

$$\frac{1}{2^4} + \frac{1}{4^4} + \frac{1}{6^4} + \dots = \beta,$$

Then $\frac{\alpha}{\beta}$ is equal to

- (1) 14 (2) 15
(3) 18 (4) 23

Answer (2)

Sol. $\alpha = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

$$\beta = \frac{1}{2^4} + \frac{1}{4^4} + \frac{1}{6^4} + \dots$$

$$= \frac{1}{2^4} \left[\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right]$$

$$\Rightarrow 16\beta = \left[\frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots \right] + \left[\frac{1}{2^4} + \frac{1}{4^4} + \frac{1}{6^4} + \dots \right]$$

$$= \alpha + \beta \Rightarrow 15\beta = \alpha \Rightarrow \frac{\alpha}{\beta} = 15$$


11. Let the function $f(x) = \frac{x}{3} + \frac{3}{x} + 3$, $x \neq 0$ be strictly increasing in $(-\infty, \alpha_1) \cup (\alpha_2, \infty)$ and strictly decreasing in $(\alpha_3, \alpha_4) \cup (\alpha_4, \alpha_5)$. Then $\sum_{i=1}^5 \alpha_i^2$ is

- equal to
(1) 48 (2) 40
(3) 36 (4) 28

Answer (3)

Sol. $f(x) = \frac{x}{3} + \frac{3}{x} + 3, x \neq 0$

$$f'(x) = \frac{1}{3} - \frac{3}{x^2} = \left(\frac{x^2 - 9}{3x^2} \right) = \frac{(x-3)(x+3)}{3x^2}$$



$$\Rightarrow f(x) > 0 \forall x \in (-\infty, -3) \cup (3, \infty)$$

$$f(x) < 0 \forall x \in (-3, 0) \cup (0, 3)$$

$$\Rightarrow \alpha_1 = -3, \alpha_2 = 3, \alpha_3 = -3, \alpha_4 = 0, \alpha_5 = 3$$

$$\Rightarrow \sum_{i=1}^5 \alpha_i^2 = (-3)^2 + (3^2) + 0^2 + (-3)^2 + (3)^2 = 4(9) = 36$$

12. If A and B are two events such that $P(A) = 0.7$, $P(B) = 0.4$ and $P(A \cap \bar{B}) = 0.5$, where \bar{B} denotes the complement of B , then $P(B | (A \cup \bar{B}))$ is equal to

- (1) $\frac{1}{2}$ (2) $\frac{1}{4}$
(3) $\frac{1}{3}$ (4) $\frac{1}{6}$

Answer (2)

Sol. $P(A) = 0.7$
 $P(B) = 0.4$
 $P(A \cap B^c) = 0.5$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths



Shreyas Lohiya
PSID: 00003389699

100 Percentile in Physics



Harsh Jha
PSID: 00014863322

100 Percentile in Physics & Chemistry



Devya Rustagi
PSID: 00014768785

99.99 Percentile



Amogh Bansal
PSID: 00014769016


OUR JEE CHAMPIONS



$$P\left(\frac{B}{A \cup B^C}\right) = \frac{P(B \cap (A \cup B^C))}{P(A \cup B^C)}$$

$$= \frac{P(A \cap B)}{P(A \cup B^C)}$$

Using Venn diagram



$$\Rightarrow \frac{P(A \cap B)}{P(A \cup B^C)} = \frac{0.2}{0.5 + 0.2 + 0.1} = \frac{2}{8} = \frac{1}{4}$$

13. Let $f(x) = x - 1$ and $g(x) = e^x$ for $x \in \mathbb{R}$. If

$$\frac{dy}{dx} = \left(e^{-2\sqrt{x}} g(f(f(x))) - \frac{y}{\sqrt{x}} \right), y(0) = 0, \text{ then } y(1) \text{ is}$$

(1) $\frac{2e-1}{e^3}$ (2) $\frac{1-e^2}{e^4}$

(3) $\frac{1-e^3}{e^4}$ (4) $\frac{e-1}{e^4}$

Answer (4)

Sol. $f(x) = x - 1$

$$f(f(x)) = (x - 1) - 1 = x - 2$$

$$g(f(f(x))) = e^{x-2}$$

$$\frac{dy}{dx} = e^{-2\sqrt{x}}(x-2) - \frac{y}{\sqrt{x}}$$

$$\Rightarrow \frac{dy}{dx} + \frac{y}{\sqrt{x}} = e^{-2\sqrt{x}} e^{x-2}$$

$$\text{IF} = e^{\int x^{-1/2} dx} = e^{2\sqrt{x}}$$

$$y \cdot e^{2\sqrt{x}} = \int e^{-2\sqrt{x}} e^{x-2} e^{2\sqrt{x}} dx$$

$$= y \cdot e^{2\sqrt{x}} = e^{x-2} + c$$

$$y(0) = 0 \Rightarrow c = e^{-2}$$

$$\therefore y \cdot e^{2\sqrt{x}} = e^{x-2} - e^{-2}$$

$$y \cdot e^2 = e^{-1} - e^{-2} = \frac{1}{e} - \frac{1}{e^2} = \frac{e-1}{e^2}$$

$$y = \frac{e-1}{e^4}$$

14. The sum of the squares of the roots of $|x-2|^2 + |x-2| - 2 = 0$ and the squares of the roots of $x^2 - 2|x-3| - 5 = 0$, is

- (1) 24 (2) 30
(3) 36 (4) 26

Answer (3)

Sol. $|x-2|^2 + |x-2| - 2 = 0$

$$\Rightarrow (|x-2| + 2)(|x-2| - 1) = 0$$

$$\Rightarrow |x-2| = 1$$

$$\Rightarrow x = 3, 1$$

$$x^2 - 2|x-3| - 5 = 0$$

$$x \geq 3$$

$$x^2 - 2(x-3) - 5 = 0$$

$$\Rightarrow x^2 - 2x + 1 = 0$$

$$\Rightarrow (x-1)^2 = 0$$

$$\Rightarrow x = 1$$

(rejected)

$$x < 3$$

$$x^2 + 2(x-3) - 5 = 0$$

$$\Rightarrow x^2 + 2x - 11 = 0$$

$$x = -1 \pm 2\sqrt{3}$$

Sum of square of roots

$$= 3^2 + 1^2 + (-1 + 2\sqrt{3})^2 + (-1 - 2\sqrt{3})^2$$

$$= 10 + (1+12)2$$

$$= 36$$

15. Let

$$A = \left\{ \theta \in [0, 2\pi] : 1 + 10 \operatorname{Re} \left(\frac{2 \cos \theta + i \sin \theta}{\cos \theta - 3i \sin \theta} \right) = 0 \right\}.$$

Then $\sum_{\theta \in A} \theta^2$ is equal to

(1) $\frac{27}{4} \pi^2$ (2) $8\pi^2$

(3) $6\pi^2$ (4) $\frac{21}{4} \pi^2$

Answer (4)

THE LEGACY OF SUCCESS CONTINUES

OUR JEE CHAMPIONS

4 STATE TOPPERS

70+ 100 PERCENTILERS
(PHYSICS & MATHS)

1000+ 99 PERCENTILERS
(PHYSICS & CHEMISTRY)

4000+ 95 PERCENTILERS
(9 ABOVE)

100 Percentile
Shreyas Lohiya
PSID: 00003389699

100 Percentile
Harsh Jha
PSID: 00014863322

99.99 Percentile
Devyu Rustagi
PSID: 00014768785

99.99 Percentile
Amogh Bansal
PSID: 00014769016

Chirag Falor
4 Year Classroom
1 AIR
JEE (Adv.)
2020

Tanishka Kabra
4 Year Classroom
1 AIR-16 CRL
JEE (Adv.)
2022

Sanvi Jain
4 Year Classroom
1 AIR-34 CRL
JEE (Main)
2024

Sol. $\operatorname{Re}\left(\frac{2\cos\theta + i\sin\theta}{\cos\theta - 3i\sin\theta}\right)$

$$\operatorname{Re}\left(\frac{(2\cos\theta + i\sin\theta)(\cos\theta + 3i\sin\theta)}{\cos^2\theta + 9\sin^2\theta}\right)$$

$$= \frac{2\cos^2\theta - 3\sin^2\theta}{1 + 8\sin^2\theta}$$

Now, $1 + 10\left(\frac{2\cos^2\theta - 3\sin^2\theta}{1 + 8\sin^2\theta}\right) = 0$

$$= 1 + 8\sin^2\theta + 20\cos^2\theta - 30\sin^2\theta = 0$$

$$= 1 - 22\sin^2\theta + 20\cos^2\theta = 0$$

$$= 1 + 20(\cos 2\theta) - 2\sin^2\theta = 0$$

$$= 20\cos 2\theta + \cos 2\theta = 0$$

$$= 21\cos 2\theta = 0$$

$$2\theta = (2n+1)\frac{\pi}{2}, n \in \mathbb{I}$$

$$2\theta = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \frac{7\pi}{2}$$

$$\Rightarrow \theta = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$$

$$\sum_{\theta \in A} \theta^2 = \frac{\pi^2}{16} + \frac{9\pi^2}{16} + \frac{25\pi^2}{16} + \frac{49\pi^2}{16}$$

$$= \frac{84\pi^2}{16}$$

$$= \frac{21\pi^2}{4}$$

16. Let a be the length of a side of a square $OABC$ with O being the origin. Its side OA makes an acute angle α with the positive x -axis and the equations of its diagonals are $(\sqrt{3}+1)x + (\sqrt{3}-1)y = 0$ and $(\sqrt{3}-1)x - (\sqrt{3}+1)y + 8\sqrt{3} = 0$. Then a^2 is equal to


(1) 24

(2) 32

(3) 16

(4) 48

Answer (4)



Sol.

$$OB: (\sqrt{3}+1)x + (\sqrt{3}-1)y = 0$$

$$AC: (\sqrt{3}-1)x - (\sqrt{3}+1)y + 8\sqrt{3} = 0$$

$$\Rightarrow (x, y) \equiv P(3 - \sqrt{3}, 3 + \sqrt{3})$$

Let $AB = a = OA$

$$\Rightarrow OA^2 + AB^2 = OB^2$$

$$2a^2 = 4\left[(3 - \sqrt{3})^2 + (3 + \sqrt{3})^2\right]$$

$$a^2 = 2 \times 24 = 48$$

17. Let $f(x)$ be a positive function and

$$I_1 = \int_{-\frac{1}{2}}^1 2xf(2x(1-2x))dx \text{ and } I_2 = \int_{-1}^2 f(x(1-x))dx.$$

Then the value of $\frac{I_2}{I_1}$ is equal to _____.

(1) 12

(2) 4

(3) 6

(4) 9

Answer (2)

Sol. $I_1 = \int_{-\frac{1}{2}}^1 2xf(2x(1-2x))dx$

$$I_1 = \int_{-\frac{1}{2}}^1 2\left(\frac{1}{2} - x\right)f\left(2\left(\frac{1}{2} - x\right)\left(1 - 2\left(\frac{1}{2} - x\right)\right)\right)dx$$

$$I_1 = \int_{-\frac{1}{2}}^1 (1-2x)f((1-2x)(2x))dx$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS

1000+ 99
PERCENTILERS

4000+ 95
PERCENTILERS

100
Percentile
in Physics
& Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in Physics
& Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



$$I_1 = \int_{-\frac{1}{2}}^1 f((1-2x)(2x)) dx - \int_{-\frac{1}{2}}^1 2xf((1-2x)(2x)) dx$$

$$2I_1 = \int_{-\frac{1}{2}}^1 f((1-2x)(2x)) dx$$

Put $2x = t$

$$2dx = dt$$

$$dx = \frac{dt}{2}$$

$$2I_1 = \frac{1}{2} \int_{-1}^2 f((1-t)(t)) dt$$

$$I_1 = \frac{1}{4} \int_{-1}^2 f((1-x)(x)) dx$$

$$I_1 = \frac{1}{4} I_2$$

$$4 \Rightarrow \frac{I_2}{I_1}$$

18. The number of integral terms in the expansion of

$$\left(\frac{1}{5^2} + \frac{1}{7^8} \right)^{1016}$$

- (1) 127
- (2) 128
- (3) 129
- (4) 130

Answer (2)

$$\text{Sol. } T_{r+1} = 1016 C_r \cdot \left(\frac{1}{5^2} \right)^r \cdot \left(\frac{1}{7^8} \right)^{1016-r}$$

$$\Rightarrow \frac{r}{8} \text{ and } \frac{1016-r}{2} \in \text{Integer}$$

$$\Rightarrow 8/r \Rightarrow r = 0, 8, 16, \dots, 1016$$

$$\Rightarrow r = 0 \times 8, 1 \times 8, \dots, 127 \times 8$$

$$\Rightarrow \text{Total } 128 \text{ } r \text{ such that } T_{r+1} \text{ is rational}$$

19. There are 12 points in a plane, no three of which are in the same straight line, except 5 points which are collinear. Then the total number of triangles that can be formed with the vertices at any three of these 12 points is

- (1) 230
- (2) 200
- (3) 220
- (4) 210

Answer (4)

Sol.



$$\text{Number of triangle} = {}^5C_1 \times {}^7C_2 + {}^7C_3 + {}^5C_2 + {}^7C_1 = 210$$

20. Let α be a solution of $x^2 + x + 1 = 0$, and for some a

$$\text{and } b \text{ in } \mathbf{R}, \begin{bmatrix} 4 & a & b \\ -1 & -1 & 2 \\ -2 & -14 & -8 \end{bmatrix} = [0 \ 0 \ 0].$$

If $\frac{4}{\alpha^4} + \frac{m}{\alpha^a} + \frac{n}{\alpha^b} = 3$, then $m + n$ is equal to _____.

- (1) 7
- (2) 11
- (3) 3
- (4) 8

Answer (2)

$$\text{Sol. } x^2 + x + 1 = 0$$



$$\begin{bmatrix} 4 & a & b \\ -1 & -1 & 2 \\ -2 & -14 & -8 \end{bmatrix}_{3 \times 3} = [0 \ 0 \ 0]$$

$$\Rightarrow [4 - a - 2b, 64 - a - 14b, 52 + 2a - 8b] = [0 \ 0 \ 0]$$

$$a + 2b = 4 \quad \dots(i)$$

$$a + 14b = 64 \quad \dots(ii)$$

Solving (i) and (ii)

$$\text{We get } a = -6, b = 5$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths



Shreyas Lohiya
PSID: 00003389699

100 Percentile in Physics



Harsh Jha
PSID: 00014863322

100 Percentile in Physics & Chemistry



Devya Rustagi
PSID: 00014768785

99.99 Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



$$\therefore \frac{4}{\alpha^4} + \frac{m}{\alpha^a} + \frac{n}{\alpha^b} = 3$$

$$\Rightarrow \frac{4}{w^4} + \frac{m}{w^{-6}} + \frac{n}{w^5} = 3$$

$$\Rightarrow 4w^2 + m + nw = 3 \quad \dots(i)$$

For $\alpha = w^2$,

$$\frac{4}{w^8} + \frac{m}{w^{-12}} + \frac{n}{w^{10}} = 3$$

$$\Rightarrow \frac{4}{w^2} + m + \frac{n}{w} = 3$$

$$\Rightarrow 4w + m + nw^2 = 3 \quad \dots(ii)$$

Adding (i) & (ii)

$$\Rightarrow 4(w^2 - w) + n(w - w^2) = 0$$

$$\Rightarrow (w^2 - w)(4 - n) = 0$$

$$\Rightarrow n = 4$$

$$\therefore 4w + m + 4w^2 = 3$$

$$\Rightarrow -4 + m = 3$$

$$\Rightarrow m = 7$$

$$\therefore m + n = 11$$

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

21. The product of the last two digits of $(1919)^{1919}$ is _____.

Answer (63)

$$\begin{aligned} \text{Sol. } (1919)^{1919} &= (1920 - 1)^{1919} \\ &= {}^{1919}C_0(1920)^{1919} - {}^{1919}C_1(1920)^{1918} + \dots + \\ &\quad \dots + {}^{1919}C_{1918}(1920) - {}^{1919}C_{1919} \end{aligned}$$

$$\text{For last two digit} \Rightarrow {}^{1919}C_{1919}(1920) - 1$$

$$= 3684479$$

\therefore Product of last two digit = 63

22. Let the domain of the function $f(x) = \cos^{-1}\left(\frac{4x+5}{3x-7}\right)$

be $[\alpha, \beta]$ and the domain of $g(x) = \log_2(2 - 6\log_{27}(2x + 5))$ be (γ, δ) .

Then $|7(\alpha + \beta) + 4(\gamma + \delta)|$ is equal to _____.

Answer (96)

$$\text{Sol. } f(x) = \cos^{-1}\left(\frac{4x+5}{3x-7}\right)$$

$$-1 \leq \frac{4x+5}{3x-7} \leq 1$$

$$\frac{7x-2}{3x-7} \geq 0, \frac{x+12}{3x-7} \leq 0$$

$$x \in \left(-\infty, \frac{2}{7}\right] \cup \left(\frac{7}{3}, \infty\right), x \in \left[-12, \frac{7}{3}\right)$$

$$\Rightarrow x \in \left[-12, \frac{2}{7}\right] \Rightarrow \alpha = -12, \beta = \frac{2}{7}$$

$$g(x) = \log_2(2 - 6\log_{27}(2x+5))$$

$$\Rightarrow 2 - 6\log_{27}(2x+5) > 0, \boxed{2x+5 > 0}$$

$$\log_{27}(2x+5) < \frac{1}{3} \Rightarrow (2x+5)^3 < 27$$

$$\Rightarrow 2x+5 < 3 \Rightarrow \boxed{x < -1}$$

$$\Rightarrow x \in \left(-\frac{5}{2}, -1\right) \Rightarrow \gamma = -\frac{5}{2}, \delta = -1$$

$$\Rightarrow |7(\alpha + \beta) + 4(\gamma + \delta)| = 96$$

23. Let r be the radius of the circle, which touches x -axis at point $(a, 0)$, $a < 0$ and the parabola $y^2 = 9x$ at the point $(4, 6)$. Then r is equal to _____.

Answer (30)

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
IN PHYSICS

1000+ 99
PERCENTILERS
IN CHEMISTRY

4000+ 95
PERCENTILERS
IN MATHS

100
Percentile
in Physics
& Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in Physics
& Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



Chirag Falor
4 Year Classroom
1 AIR
JEE (Adv.)
2020



Tanishka Kabra
4 Year Classroom
1 AIR-16 CRL
JEE (Adv.)
2022



Sanvi Jain
4 Year Classroom
1 AIR-34 CRL
JEE (Main)
2024

Sol. Equation of tangent to $y^2 = gx$ at $(4, 6)$ is $3x - 4y + 12 = 0$

Equation of circle is $(x - 4)^2 + (y - 6)^2 + \lambda(3x - 4y + 12) = 0$

$$\Rightarrow x^2 + y^2 + (3\lambda - 8)x + (-12 - 4\lambda)y + 52 + 12\lambda = 0$$

$$\therefore 2\sqrt{g^2 - c} = 0 \Rightarrow g^2 = c$$

$$\Rightarrow \left(-\frac{3\lambda - 8}{2}\right)^2 = 52 + 12\lambda$$

$$\Rightarrow 9\lambda^2 + 64 - 48\lambda = 208 + 48\lambda \Rightarrow 9\lambda^2 - 96\lambda - 144 = 0$$

$$\Rightarrow \lambda = 12, -\frac{2}{3} \Rightarrow f = -30, -\frac{14}{3}$$

$$\Rightarrow r = \sqrt{g^2 + f^2 - c} = |f| = |-(2\lambda + 6)|$$

\therefore centre lies in 2nd quadrant

$$\Rightarrow 3\lambda - 8 > 0 \Rightarrow \lambda > \frac{8}{3}$$


$$\Rightarrow \lambda = 12, f = -30, r = 30$$

24. Let the area of the bounded region $\{(x, y) : 0 \leq 9x \leq y^2, y \geq 3x - 6\}$ be A. Then 6A is equal to _____.

Answer (Bonus)

Sol. The given problem has area bounded $\rightarrow \infty$

However, if we correct in to $y^2 \leq 4x$ then here is the solution.



Required area

$$= \int_{-3}^3 \left(\frac{y+6}{3} - \frac{y^2}{9} \right) dy$$

$$= \frac{27}{2} \text{ sq. units}$$

25. Let the area of the triangle formed by the lines $x + 2 = y - 1 = z$, $\frac{x-3}{5} = \frac{y}{-1} = \frac{z-1}{1}$ and

$$\frac{x}{-3} = \frac{y-3}{3} = \frac{z-2}{1} \text{ be } A. \text{ Then } A^2 \text{ is equal to } \underline{\hspace{2cm}}.$$

$$\frac{x}{-3} = \frac{y-3}{3} = \frac{z-2}{1} \text{ be } A. \text{ Then } A^2 \text{ is equal to } \underline{\hspace{2cm}}.$$

Answer (56)

Sol. $L_1 = \frac{x+2}{1} = \frac{y-1}{1} = \frac{z}{1} = \lambda$, any point on it $(\lambda - 2, \lambda + 1, \lambda)$

$L_2 = \frac{x-3}{5} = \frac{y}{-1} = \frac{z-1}{1} = \mu$, any point on it $(5\mu + 3, -\mu, \mu + 1)$

$L_3 = \frac{x}{-3} = \frac{y-3}{3} = \frac{z-2}{1} = k$, any point on it $(-3k, 3k + 3, k + 2)$

$P \equiv$ point of intersection of L_1 and $L_2 = (-2, 1, 0)$

$Q \equiv$ point of intersection of L_1 and $L_3 = (0, 3, 2)$

$R \equiv$ point of intersection of L_2 and $L_3 = (3, 0, 1)$

$$\overline{PQ} = 2\hat{i} + 2\hat{j} + 2\hat{k}$$

$$\overline{PR} = 5\hat{i} - \hat{j} + \hat{k}$$

$$A = \frac{1}{2} |\overline{PQ} \times \overline{PR}| = \sqrt{56}$$

$$A^2 = 56$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile



Shreyas Lohiya
PSID: 00003389699

100 Percentile



Harsh Jha
PSID: 00014863322

100 Percentile



Devya Rustagi
PSID: 00014768785

99.99 Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



PHYSICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answers :

26. A rod of linear mass density ' λ ' and length ' L ' is bent to form a ring of radius ' R '. Moment of inertia of ring about any of its diameter is

- (1) $\frac{\lambda L^3}{4\pi^2}$ (2) $\frac{\lambda L^3}{8\pi^2}$
 (3) $\frac{\lambda L^3}{12}$ (4) $\frac{\lambda L^3}{16\pi^2}$

Answer (2)

Sol. $2\pi R = L \Rightarrow M = \lambda L$

$$R = \frac{L}{2\pi}$$

Moment of Inertia about diameter

$$= \frac{MR^2}{2} = \frac{M}{2} \left(\frac{L}{2\pi} \right)^2 = \frac{\lambda L^3}{8\pi^2}$$

27. In a Young's double slit experiment, the source is white light. One of the slits is covered by red filter and another by a green filter. In this case

- (1) There shall be alternate interference fringes of red and green.
 (2) There shall be an interference pattern for red distinct from that for green.
 (3) There shall be no interference fringes.
 (4) There shall be an interference pattern, where each fringe's pattern center is green and outer edges is red.

Answer (3)

Sol. Frequency of Green and frequency of Red will be different. No interference pattern is observed for two lights of different frequencies as phase difference does not remain constant.

28. Two balls with same mass and initial velocity, are projected at different angles in such a way that maximum height reached by first ball is 8 times higher than that of the second ball. T_1 and T_2 are the total flying times of first and second ball, respectively, then the ratio of T_1 and T_2 is

- (1) $2\sqrt{2} : 1$ (2) $\sqrt{2} : 1$
 (3) $4 : 1$ (4) $2 : 1$

Answer (1)

Sol. $H_1 = \frac{u^2 \sin^2 \theta_1}{2g}$ $H_2 = \frac{u^2 \sin^2 \theta_2}{2g}$

$T_1 = \frac{2u \sin \theta_1}{g}$ $T_2 = \frac{2u \sin \theta_2}{g}$

$\frac{H_1}{H_2} = \frac{\sin^2 \theta_1}{\sin^2 \theta_2}$ $\frac{T_1}{T_2} = \frac{\sin \theta_1}{\sin \theta_2}$

$8 = \left(\frac{T_1}{T_2} \right)^2 \Rightarrow \frac{T_1}{T_2} = 2\sqrt{2}$

$T_1 : T_2 = 2\sqrt{2} : 1$

29. For a nucleus of mass number A and radius R , the mass density of nucleus can be represented as

- (1) A^3 (2) $A^{\frac{2}{3}}$
 (3) $A^{\frac{1}{3}}$ (4) Independent of A

Answer (4)

Sol. $R = R_0 A^{1/3}$

Density = $\frac{\text{Mass number}}{\text{Volume}} = \frac{A}{\frac{4}{3}\pi R_0^3 A} = \text{constant}$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
in Physics

1000+ 99
PERCENTILERS
in Physics

4000+ 95
PERCENTILERS
in Physics

100 Percentile in Physics & Maths

Shreyas Lohiya
 PSID: 00003389699

100 Percentile in Physics

Harsh Jha
 PSID: 00014863322

100 Percentile in Physics & Chemistry

Devya Rustagi
 PSID: 00014768785

99.99 Percentile

Amogh Bansal
 PSID: 00014769016


OUR JEE CHAMPIONS

Chirag Falor
 4 Year Classroom
1 AIR JEE (Adv.) 2020

Tanishka Kabra
 4 Year Classroom
1 AIR-16 CRL JEE (Adv.) 2022


Sanvi Jain
 4 Year Classroom
1 AIR-34 CRL JEE (Main) 2024

30. Figure shows a current carrying square loop ABCD of edge length is 'a' lying in a plane. If the resistance of the ABC part is r and that of ADC part is 2r, then the magnitude of the resultant magnetic field at centre of the square loop is



- (1) $\frac{\sqrt{2}\mu_0 I}{3\pi a}$ (2) $\frac{\mu_0 I}{2\pi a}$
 (3) $\frac{2\mu_0 I}{3\pi a}$ (4) $\frac{3\pi\mu_0 I}{\sqrt{2}a}$

Answer (1)



$$R_{ABC} = r \qquad R_{ADC} = 2r$$

$$i_1 = \frac{2I}{3} \qquad i_2 = \frac{I}{3}$$

$$B_{\text{centre}} = \frac{2(\mu_0 \sqrt{2})}{4\pi \left(\frac{a}{2}\right)} \left[\frac{2I}{3} - \frac{I}{3} \right] = \sqrt{2} \frac{\mu_0 I}{3\pi a}$$

31. A convex lens of focal length 30 cm is placed in contact with a concave lens of focal length 20 cm. An object is placed at 20 cm to the left of this lens system. The distance of the image from the lens in cm is _____.

- (1) 30 (2) $\frac{60}{7}$
 (3) 15 (4) 45

Answer (3)

Sol. $\frac{1}{F} = \frac{1}{f_1} + \frac{1}{f_2} = \frac{1}{30} + \frac{1}{-20} = -\frac{1}{60}$

$$v = \frac{Fu}{u+F} = \frac{-60 \times -20}{-20 + -60} = -15 \text{ cm}$$

32. The amplitude and phase of a wave that is formed by the superposition of two harmonic travelling waves, $y_1(x, t) = 4 \sin(kx - \omega t)$ and $y_2(x, t) = 2 \sin\left(kx - \omega t + \frac{2\pi}{3}\right)$, are :

(Take the angular frequency of initial waves same as ω)

- (1) $\left[2\sqrt{3}, \frac{\pi}{6}\right]$ (2) $\left[6, \frac{2\pi}{3}\right]$
 (3) $\left[6, \frac{\pi}{3}\right]$ (4) $\left[\sqrt{3}, \frac{\pi}{6}\right]$

Answer (1)

Sol. $A = \sqrt{A_1^2 + A_2^2 + 2A_1A_2 \cos \phi}$

$$\phi = \frac{2\pi}{3}$$

$$A = \sqrt{4^2 + 2^2 + 2 \times 4 \times 2 \cos \frac{2\pi}{3}} = 2\sqrt{3}$$

$$\tan \alpha = \frac{2 \sin \phi}{4 + 2 \cos \phi} = \frac{1}{\sqrt{3}}$$

$$\alpha = \frac{\pi}{6}$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths

Shreyas Lohiya
 PSID: 00003389699

100 Percentile in Physics

Harsh Jha
 PSID: 00014863322

100 Percentile in Physics & Chemistry

Devya Rustagi
 PSID: 00014768785

99.99 Percentile

Amogh Bansal
 PSID: 00014769016

OUR JEE CHAMPIONS

Chirag Falor
 4 Year Classroom
1 AIR JEE (Adv.) 2020

Tanishka Kabra
 4 Year Classroom
1 AIR-16 CRL JEE (Adv.) 2022

Sanvi Jain
 4 Year Classroom
1 AIR-34 CRL JEE (Main) 2024

33. A body of mass 2 kg moving with velocity of $\vec{v}_{in} = 3\hat{i} + 4\hat{j} \text{ ms}^{-1}$ enters into a constant force field of 6N directed along positive z-axis. If the body remains in the field for a period of $\frac{5}{3}$ seconds, then velocity of the body when it emerges from force field is :

- (1) $3\hat{i} + 4\hat{j} + 5\hat{k}$ (2) $3\hat{i} + 4\hat{j} - 5\hat{k}$
(3) $4\hat{i} + 3\hat{j} + 5\hat{k}$ (4) $3\hat{i} + 4\hat{j} + \sqrt{5}\hat{k}$

Answer (1)

Sol. $\vec{F} \cdot t = m(\vec{v}_2 - \vec{v}_1)$

$$\frac{6}{2} \times \frac{5}{3} \hat{k} = \vec{v}_2 - \vec{v}_1$$

$$\vec{v}_2 = \vec{v}_1 + 5\hat{k} = 3\hat{i} + 4\hat{j} + 5\hat{k}$$

34. A quantity Q is formulated as $X^{-2}Y^{\frac{3}{2}}Z^{\frac{2}{5}}$. X, Y and Z are independent parameters which have fractional errors of 0.1, 0.2 and 0.5, respectively in measurement. The maximum fractional error of Q is

- (1) 0.1 (2) 0.6
(3) 0.7 (4) 0.8

Answer (3)

Sol. $\frac{\Delta Q}{Q} = -2 \frac{(\pm \Delta x)}{x} + \frac{3}{2} \frac{(\pm \Delta y)}{y} + \frac{2}{5} \frac{(\pm \Delta z)}{z}$

For maximum fractional error

$$\frac{\Delta Q}{Q} = \frac{2\Delta x}{x} + \frac{3}{2} \frac{\Delta y}{y} + \frac{2}{5} \frac{\Delta z}{z}$$

$$= 2 \times 0.1 + \frac{3}{2} \times 0.2 + \frac{2}{5} \times 0.5$$

$$= 0.7$$

35. A monoatomic gas having $\gamma = \frac{5}{3}$ is stored in a thermally insulated container and the gas is suddenly compressed to $\left(\frac{1}{8}\right)^{\text{th}}$ of its initial volume.

The ratio of final pressure and initial pressure is (γ is the ratio of specific heats of the gas at constant pressure and at constant volume)


- (1) 40 (2) 28
(3) 32 (4) 16

Answer (3)

Sol. $P_1 V_1^\gamma = P_2 V_2^\gamma$ $v_2 = \frac{v_1}{8}$

$$\frac{P_2}{P_1} = \left(\frac{v_1}{v_2}\right)^\gamma = (8)^{\frac{5}{3}} = 32$$

36. Electric charge is transferred to an irregular metallic disk as shown in figure. If $\sigma_1, \sigma_2, \sigma_3$ and σ_4 are charge densities at given points then, choose the correct answer from the options given below:



- A. $\sigma_1 > \sigma_3; \sigma_2 = \sigma_4$
B. $\sigma_1 > \sigma_2; \sigma_3 > \sigma_4$
C. $\sigma_1 > \sigma_3 > \sigma_2 = \sigma_4$
D. $\sigma_1 < \sigma_3 < \sigma_2 = \sigma_4$
E. $\sigma_1 = \sigma_2 = \sigma_3 = \sigma_4$
(1) A and C Only (2) D and E Only
(3) A, B and C Only (4) B and C Only

Answer (3)

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
max 1 (each 1 below)

1000+ 99
PERCENTILERS
& ABOVE

4000+ 95
PERCENTILERS
& ABOVE

100
Percentile
in
Physics
& Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in
Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in
Physics
& Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



Sol. $\sigma \propto \frac{1}{R}$ $R \rightarrow$ radius of curvature

$R_2 = R_4 > R_3 > R_1$

$\Rightarrow \sigma_1 > \sigma_3 > \sigma_4 = \sigma_2$

37. Two strings with circular cross section and made of same material are stretched to have same amount of tension. A transverse wave is then made to pass through both the strings. The velocity of the wave in the first string having the radius of cross section R is v_1 , and that in the other string having radius of cross

section $R/2$ is v_2 . Then $\frac{v_2}{v_1} =$

- (1) 8
- (2) 2
- (3) $\sqrt{2}$
- (4) 4

Answer (2)

Sol. $v = \sqrt{\frac{T}{\mu}}$ $\mu = \rho\pi R^2$

$\frac{v_1}{v_2} = \sqrt{\frac{R_2^2}{R_1^2}} = \frac{R_2}{R_1} = \frac{1}{2}$

$\frac{v_2}{v_1} = 2$

38. An infinitely long wire has uniform linear charge density $\lambda = 2$ nC/m. The net flux through a Gaussian cube of side length $\sqrt{3}$ cm, if the wire passes through any two corners of the cube, that are maximally displaced from each other, would be x Nm²C⁻¹, where x is

[Neglect any edge effects and use $\frac{1}{4\pi\epsilon_0} = 9 \times 10^9$

SI units]

- (1) 0.72π
- (2) 2.16π
- (3) 1.44π
- (4) 6.48π

Answer (2)

Sol. $\phi = \frac{q_{\text{enclosed}}}{\epsilon_0}$ $l = \sqrt{3}a = 3$

$q = \lambda l$

$\phi = \frac{\lambda l}{\epsilon_0} = 2 \times 10^{-9} \times 3 \times 10^{-2} \times 4\pi \times 9 \times 10^9$

$= 2.16 \pi$

39. A block of mass 2 kg is attached to one end of a massless spring whose other end is fixed at a wall. The spring-mass system moves on a frictionless horizontal table. The spring's natural length is 2 m and spring constant is 200 N/m. The block is pushed such that the length of the spring becomes 1 m and then released. At distance x m ($x < 2$) from the wall, the speed of the block will be


(1) $10[1 - (2 - x)^2]^2$ m/s

(2) $10[1 - (2 - x)^2]^{1/2}$ m/s

(3) $10[1 - (2 - x)]^{3/2}$ m/s

(4) $10[1 - (2 - x)^2]$ m/s

Answer (2)



Sol.

Energy conservation

$\frac{1}{2}k(1)^2 = \frac{1}{2}mv^2 + \frac{1}{2}k(2 - x)^2$

Compression in the spring = $(2 - x)$

$\Rightarrow \frac{1}{2}k(1 - (2 - x)^2) = \frac{1}{2}mv^2$

$\Rightarrow v = [100(1 - (2 - x)^2)]^{1/2}$

$v = 10[1 - (2 - x)]^{1/2}$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths

Shreyas Lohiya
 PSID: 00003389699

100 Percentile in Physics

Harsh Jha
 PSID: 00014863322

100 Percentile in Physics & Chemistry

Devya Rustagi
 PSID: 00014768785

99.99 Percentile

Amogh Bansal
 PSID: 00014769016

OUR JEE CHAMPIONS

Chirag Falor
 4 Year Classroom
1 AIR JEE (Adv.) 2020

Tanishka Kabra
 4 Year Classroom
1 AIR-16 CRL JEE (Adv.) 2022

Sanvi Jain
 4 Year Classroom
1 AIR-34 CRL JEE (Main) 2024

40. Given below are two statements: one is labelled as **Assertion A** and the other is labelled as **Reason R**.

Assertion A : Work done in moving a test charge between two points inside a uniformly charged spherical shell is zero, no matter which path is chosen.

Reason R : Electrostatic potential inside a uniformly charged spherical shell is constant and is same as that on the surface of the shell.

In the light of the above statements, choose the **correct** answer from the options given below

- (1) **A** is false but **R** is true
- (2) Both **A** and **R** are true and **R** is the correct explanation of **A**
- (3) Both **A** and **R** are true but **R** is **NOT** the correct explanation of **A**
- (4) **A** is true but **R** is false

Answer (2)

Sol. Both A & R are correct and R is correct explanation

$$\Delta W = q(\Delta V)$$

$$\Delta V = 0$$

$$\Rightarrow \Delta W = 0$$

41. Two metal spheres of radius R and $3R$ have same surface charge density σ . If they are brought in contact and then separated, the surface charge density on smaller and bigger sphere becomes σ_1 and σ_2 ,

respectively. The ratio $\frac{\sigma_1}{\sigma_2}$ is


- (1) 9
- (2) 3
- (3) $\frac{1}{9}$
- (4) $\frac{1}{3}$

Answer (2)

Sol. $v_1 = v_2 \Rightarrow \frac{Q_1}{R_1} = \frac{Q_2}{R_2} \Rightarrow \frac{\sigma_1 R_1^2}{R_1} = \frac{\sigma_2 R_2^2}{R_2}$

$$\Rightarrow \frac{\sigma_1}{\sigma_2} = \frac{R_2}{R_1} = 3$$


42. The output voltage in the following circuit is (Consider ideal diode case)



- (1) 10 V
- (2) 0 V
- (3) +5 V
- (4) -5 V

Answer (2)

Sol.



$D_1 =$ Reverse biased

$D_2 =$ Forward biased

$$V_{(out)} = 0 \text{ V}$$

43. Water falls from a height of 200 m into a pool. Calculate the rise in temperature of the water assuming no heat dissipation from the water in the pool.

(Take $g = 10 \text{ m/s}^2$, specific heat of water = 4200 J/(kg K))

- (1) 0.36 K
- (2) 0.23 K
- (3) 0.48 K
- (4) 0.14 K

Answer (3)

Sol. $mgh = ms\Delta T$

$$\Delta T = \frac{gh}{s} = \frac{10 \times 200}{4200} = 0.48 \text{ K}$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Math



Shreyas Lohiya
PSID: 00003389699

100 Percentile in Physics



Harsh Jha
PSID: 00014863322

100 Percentile in Physics & Chemistry



Devya Rustagi
PSID: 00014768785

99.99 Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



44. A 3 m long wire of radius 3 mm shows an extension of 0.1 mm when loaded vertically by a mass of 50 kg in an experiment to determine Young's modulus. The value of Young's modulus of the wire as per this experiment is $P \times 10^{11} \text{ Nm}^{-2}$, where the value of P is: (Take $g = 3\pi \text{ m/s}^2$)

- (1) 2.5 (2) 25
(3) 5 (4) 10

Answer (3)

Sol. $Y = \frac{F/A}{\Delta L/L} = \frac{FL}{A\Delta L} = \frac{(500N)(3)}{\pi \times 9 \times 10^{-6} \times 1 \times 10^{-4}}$
 $= 5 \times 10^{11} \text{ N/m}^2$

45. A concave-convex lens of refractive index 1.5 and the radii of curvature of its surfaces are 30 cm and 20 cm, respectively. The concave surface is upwards and is filled with a liquid of refractive index 1.3. The focal length of the liquid-glass combination will be

- (1) $\frac{600}{11}$ cm (2) $\frac{500}{11}$ cm
(3) $\frac{700}{11}$ cm (4) $\frac{800}{11}$ cm

Answer (1)



Sol.

$$\frac{1}{F} = \frac{1}{F_1} + \frac{1}{F_2}$$

$$\frac{1}{F} = \frac{1}{120} + \frac{1}{100} = \frac{10+12}{1200} = \frac{0.3}{30} = \frac{1}{100}$$


$$\frac{1}{F} = \frac{22}{1200}$$

$$\frac{1}{F_2} = (1.5 - 1) \left(\frac{-1}{30} - \frac{1}{-20} \right)$$

$$F = \frac{600}{11} = 0.5 \left(\frac{3}{60} - \frac{2}{60} \right) = \frac{1}{120}$$

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.



46.

Space between the plates of a parallel plate capacitor of plate area 4 cm^2 and separation of (d) 1.77 mm, is filled with uniform dielectric materials with dielectric constants (3 and 5) as shown in figure. Another capacitor of capacitance 7.5 pF is connected in parallel with it. The effective capacitance of this combination is _____ pF.

(Given $\epsilon_0 = 8.85 \times 10^{-12} \text{ F/m}$)

Answer (15)

Sol. $\frac{1}{C} = \frac{1}{C_1} + \frac{1}{C_2} = \frac{d/2}{Ak_1\epsilon_0} + \frac{d/2}{Ak_2\epsilon_0}$
 $= \left(\frac{1}{k_1} + \frac{1}{k_2} \right) \frac{d}{2A\epsilon_0}$
 $= \left(\frac{1}{3} + \frac{1}{5} \right) \frac{d}{2A\epsilon_0}$
 $\frac{1}{C} = \frac{4}{15 A\epsilon_0}$
 $C = \frac{15}{4} \times \frac{A\epsilon_0}{d} = \frac{15}{4} \times \frac{4 \times 10^{-4} \times 8.85 \times 10^{-12}}{1.77 \times 10^{-3}}$
 $= 7.5 \text{ pF}$
 $C + C_3 = (7.5 + 7.5) \text{ pF} = 15 \text{ pF}$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths

Shreyas Lohiya
 PSID: 00003389699

100 Percentile in Physics

Harsh Jha
 PSID: 00014863322

100 Percentile in Physics & Chemistry

Devya Rustagi
 PSID: 00014768785

99.99 Percentile

Amogh Bansal
 PSID: 00014769016

OUR JEE CHAMPIONS

Chirag Falor
 4 Year Classroom
1 AIR-16 CRL JEE (Adv.) 2020

Tanishka Kabra
 4 Year Classroom
1 AIR-16 CRL JEE (Adv.) 2022

Sanvi Jain
 4 Year Classroom
1 AIR-34 CRL JEE (Main) 2024

47. An electron is released from rest near an infinite non-conducting sheet of uniform charge density ' $-\sigma$ '. The rate of change of de-Broglie wave length associated with the electron varies inversely as n^{th} power of time. The numerical value of n is _____.

Answer (2)

Sol. $F = qE = q \left(\frac{\sigma}{2\epsilon_0} \right) = \frac{\Delta p}{\Delta t}$

$$\lambda = \frac{h}{p} \quad \frac{d\lambda}{dt} = \frac{h}{p^2} \times \frac{\Delta p}{\Delta t} = \frac{h}{p^2} \times \frac{q\sigma}{2\epsilon_0}$$

$$\frac{d\lambda}{dt} \propto \frac{1}{p^2} \propto \frac{1}{t^2} \propto \frac{1}{t^n}$$


$$p = \frac{q\sigma}{2\epsilon_0} t$$

$$n = 2$$

48. A cube having a side of 10 cm with unknown mass and 200 gm mass were hung at two ends of a uniform rigid rod of 27 cm long. The rod along with masses was placed on a wedge keeping the distance between wedge point and 200 gm weight as 25 cm. Initially the masses were not at balance. A beaker is placed beneath the unknown mass and water is added slowly to it. At given point the masses were in balance and half volume of the unknown mass was inside the water.

(Take the density of unknown mass is more than that of the water, the mass did not absorb water and water density is 1 gm/cm³.) The unknown mass is _____ kg.

Answer (3)



$$25 \times 0.2 \times g = 2 \times (m - \rho \times v) g$$

$$m - \rho v = 2.5 \text{ kg}$$

$$\rho v = \frac{1 \times 10^{-3} \text{ kg}}{\text{cm}^3} \times \frac{10^3 \text{ cm}^3}{2} = \frac{1}{2} \text{ kg}$$

$$m = 3 \text{ kg}$$

49. A sample of a liquid is kept at 1 atm. It is compressed to 5 atm which leads to change of volume of 0.8 cm³. If the bulk modulus of the liquid is 2 GPa, the initial volume of the liquid was _____ litre.

(Take 1 atm = 10⁵ Pa)

Answer (4)

Sol. $-\frac{\Delta v B}{v} = \Delta P \Rightarrow 4 \times 10^5 \text{ Pa} = \frac{2 \times 10^9 \text{ Pa} \times 10^{-6} \times 0.8}{v}$

$$v = \frac{2 \times 10^9 \times 10^{-6} \times 0.8}{4 \times 10^5} = 4 \times 10^{-3} \text{ m}^3$$

$$\Rightarrow 4 \text{ litre}$$

50. A thin solid disk of 1 kg is rotating along its diameter axis at the speed of 1800 rpm. By applying an external torque of 25 π Nm for 40s, the speed increases to 2100 rpm. The diameter of the disk is _____ m.

Answer (40)

Sol. $\tau dt = I \Delta \omega$

$$\Rightarrow 25\pi \times 40 = I(300) \times \frac{2\pi}{60}$$

$$I = \frac{25 \times 60 \times 40}{300 \times 2} = 100 = \frac{MR^2}{4}$$

$$R^2 = 400$$

$$R = 20 \text{ m}$$

$$D = 40 \text{ m}$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
(max. 1 score, 1 subject)

1000+ 99
PERCENTILERS
& ABOVE

4000+ 95
PERCENTILERS
& ABOVE

100
Percentile
in
Physics
&
Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in
Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in
Physics
&
Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



CHEMISTRY

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer :

51. Correct statements for an element with atomic number 9 are

- A. There can be 5 electrons for which $m_s = +\frac{1}{2}$ and 4 electrons for which $m_s = -\frac{1}{2}$
- B. There is only one electron in p_z orbital
- C. The last electron goes to orbital with $n = 2$ and $l = 1$
- D. The sum of angular nodes of all the atomic orbitals is 1


Choose the correct answer from the options given below :

- (1) A and B only (2) C and D only
- (3) A and C only (4) A, C and D only

Answer (3)

Sol. $Z = 9$

Electronic configuration \Rightarrow



- There can be 5e⁻ for which $m_s = +\frac{1}{2}$ and 4e⁻ for which $m_s = -\frac{1}{2}$

- There can be 2e⁻ in p_z orbital
 - Last e⁻ enters 2p ($n = 2, l = 1$)
 - Angular nodes = l
- \therefore Sum of angular nodes = $0 + 0 + 3 = 3$

52. Match List-I with List-II :

	List-I		List-II
A.	[Ni(CO) ₄]	I.	Tetrahedral, 2.8 BM
B.	[Ni(CN) ₄] ²⁻	II.	Square planar, 0 BM
C.	[NiCl ₄] ²⁻	III.	Tetrahedral, 0 BM
D.	[MnBr ₄] ²⁻	IV.	Tetrahedral, 5.9 BM

Choose the correct answer from the options given below

- (1) A(III), B(II), C(I), D(IV)
- (2) A(I), B(II), C(III), D(IV)
- (3) A(IV), B(I), C(III), D(II)
- (4) A(III), B(IV), C(II), D(I)

Answer (1)

Sol. A) Ni(CO)₄ \rightarrow Ni(0) \Rightarrow sp³, tetrahedral, 0 BM (3d¹⁰)(pairing)

B) [Ni(CN)₄]²⁻ \rightarrow Ni²⁺ \Rightarrow dsp², square planar, 0 BM (3d⁸)(pairing)

C) [NiCl₄]²⁻ \rightarrow Ni²⁺(no pairing) \Rightarrow sp³, tetrahedral, 2.8 BM



D) [MnBr₄]²⁻ \Rightarrow Mn²⁺ \Rightarrow 3d⁵ (no pairing)

THE LEGACY OF SUCCESS CONTINUES

OUR JEE CHAMPIONS

JEE Main (Session-1) 2025

4 STATE TOPPERS



100 Percentile in Physics & Maths
100 Percentile in Chemistry
100 Percentile in Biology & Computer

Shreyas Lohiya
PSID: 00003389699

70+ 100 PERCENTILERS (max. 1 each, 1 total)



100 Percentile in Physics & Chemistry

Harsh Jha
PSID: 00014863322

1000+ 99 PERCENTILERS & ABOVE



99.99 Percentile

Devyu Rustagi
PSID: 00014768785

4000+ 95 PERCENTILERS & ABOVE



99.99 Percentile

Amogh Bansal
PSID: 00014769016




Chirag Falor
4 Year Classroom
1 AIR-36 (Male)
JEE (Main) 2020



Tanishka Kabra
4 Year Classroom
1 AIR-36 CBZ (Female)
JEE (Main) 2022




Sanvi Jain
4 Year Classroom
1 AIR-36 CBZ (Female)
JEE (Main) 2024



$$\mu = 5.9 \text{ BM}$$

(A-III, B-II, C-I, D-IV)



The freezing point depression of a 0.1 m aqueous solution of a monobasic weak acid HA is 0.20°C . The dissociation constant for the acid is

Given: $K_f(\text{H}_2\text{O}) = 1.8 \text{ K kg mol}^{-1}$, molality \equiv molarity

- (1) 1.1×10^{-2} (2) 1.38×10^{-3}
(3) 1.90×10^{-3} (4) 1.89×10^{-1}

Answer (2)

Sol. $\Delta T_f = iK_f m$

$$i = \frac{\Delta T_f}{K_f \cdot m}$$

$$i = \frac{0.20}{1.8 \times 0.1} = 1.11$$

$$i = 1.11$$

$$\alpha = \frac{i-1}{n-1} \text{ (for HA, } n=2\text{)}$$

$$\alpha = \frac{1.11-1}{1} = 0.11$$

$$K_a = \frac{\alpha^2}{1-\alpha} = \frac{0.1 \times (0.11)^2}{1-0.11} = 1.38 \times 10^{-3}$$

54. Given below are two statements :

Statement-I : A homoleptic octahedral complex, formed using monodentate ligands, will not show stereoisomerism

Statement-II : cis and trans platin are heteroleptic complexes of Pd

In the light of the above statements, choose the correct answer from the options given below


- (1) Both Statement I and Statement II are true
(2) Statement I is true but Statement II is false

- (3) Both Statement I and Statement II are false
(4) Statement I is false but Statement II is true

Answer (2)

Sol. $[\text{Ma}_6]$ type complex will not show stereoisomerism, where a is monodentate ligand

Cis and trans-platin are heterolytic complexes of Pt(Platinum). Formula is $[\text{Pt}(\text{NH}_3)_2 \text{Cl}_2]$



55. Which of the following binary mixture does not show the behaviour of minimum boiling azeotropes?

- (1) $\text{C}_6\text{H}_5\text{OH} + \text{C}_6\text{H}_5\text{NH}_2$ (2) $\text{H}_2\text{O} + \text{CH}_3\text{COC}_2\text{H}_5$
(3) $\text{CH}_3\text{OH} + \text{CHCl}_3$ (4) $\text{CS}_2 + \text{CH}_3\text{COCH}_3$

Answer (1)

Sol. The solution showing positive deviation from Raoult's law will form minimum boiling azeotrope. Phenol + Aniline shows negative deviation, so they will not form minimum boiling azeotrope

56. Which one of the following reactions will not lead to the desired ether formation in major proportion ?

(iso-Bu \Rightarrow isobutyl, sec-Bu \Rightarrow sec-butyl, nPr \Rightarrow n-propyl, tBu \Rightarrow tert-butyl, Et \Rightarrow ethyl)

- (1) $\text{iso-Bu-O}^-\text{Na}^+ + \text{sec-BuBr} \rightarrow \text{sec-Bu-O-iso-Bu}$
(2) $\text{tBu-O}^-\text{Na}^+ + \text{EtBr} \rightarrow \text{tBu-O-Et}$
(3) $\text{Cyclohexyl-O}^-\text{Na}^+ + \text{CH}_3\text{Br} \rightarrow \text{Cyclohexyl-O-CH}_3$
(4) $\text{Na}^+ \text{O}^- \text{C}_6\text{H}_5 + \text{n-PrBr} \rightarrow \text{n-Pr-O-C}_6\text{H}_5$

Answer (1)

THE LEGACY OF SUCCESS CONTINUES

OUR JEE CHAMPIONS

JEE Main (Session-1) 2025

4 STATE TOPPERS



100 Percentile in Physics & Maths
100 Percentile

Shreyas Lohiya
PSID: 00003389699

70+ 100 PERCENTILERS (max. 1 each, 1 below)



100 Percentile in Physics & Maths
100 Percentile

Harsh Jha
PSID: 00014863322

1000+ 99 PERCENTILERS & ABOVE



100 Percentile in Physics & Chemistry
99.99 Percentile

Devyas Rustagi
PSID: 00014768785

4000+ 95 PERCENTILERS & ABOVE



100 Percentile
99.99 Percentile

Amogh Bansal
PSID: 00014769016




Chirag Falor
4 Year Classroom
1 AIR-36 (Male)
JEE (Main) 2020



Tanishka Kabra
4 Year Classroom
1 AIR-16 CBZ
JEE (Main) 2022




Sanvi Jain
4 Year Classroom
1 AIR-34 CBZ
JEE (Main) 2024



For 2° RX \Rightarrow elimination would be more favourable than substitution

For Williamson's synthesis of ether





57. Match the List-I with List-II

	List-I		List-II
A.	Carbocation	I.	Species that can supply a pair of electrons
B.	C-Free radical	II.	Species that can receive a pair of electrons
C.	Nucleophile	III.	sp^2 hybridized carbon with empty p-orbital
D.	Electrophile	IV.	sp^2/sp^3 hybridised carbon with one unpaired electron

Choose the correct answer from the options given below:

- (1) A(III), B(IV), C(II), D(I)
- (2) A(II), B(III), C(I), D(IV)
- (3) A(IV), B(II), C(III), D(I)
- (4) A(III), B(IV), C(I), D(II)

Answer (4)



Nucleophile $\Rightarrow e^-$ rich species like anions etc.





\Rightarrow can supply a pair of electrons

Electrophile $\Rightarrow e^-$ deficient species \Rightarrow can receive a pair of electrons.


58. The correct decreasing order of spin only magnetic moment values (BM) of Cu^+ , Cu^{2+} , Cr^{2+} and Cr^{3+} ions is :

- (1) $\text{Cu}^+ > \text{Cu}^{2+} > \text{Cr}^{3+} > \text{Cr}^{2+}$
- (2) $\text{Cu}^{2+} > \text{Cu}^+ > \text{Cr}^{2+} > \text{Cr}^{3+}$
- (3) $\text{Cr}^{3+} > \text{Cr}^{2+} > \text{Cu}^+ > \text{Cu}^{2+}$
- (4) $\text{Cr}^{2+} > \text{Cr}^{3+} > \text{Cu}^{2+} > \text{Cu}^+$

Answer (4)



So order :



THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
IN PHYSICS
(OVER 1 MILLION)

1000+ 99
PERCENTILERS
IN CHEMISTRY

4000+ 95
PERCENTILERS
IN MATHS

100
Percentile
in Physics
& Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in Physics
& Chemistry



Devya Rustagi
PSID: 00014768785


99.99
Percentile




Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS







'P' is



Answer (1)




60. Choose the correct set of reagents for the following conversion.




- (1) $\text{Cl}_2/\text{anhy. AlCl}_3$; Br_2/Fe ; alc. KOH
 (2) Br_2/Fe ; Cl_2, Δ ; alc. KOH
 (3) $\text{Br}_2/\text{anhy. AlCl}_3$; Cl_2, Δ ; aq. KOH
 (4) Cl_2/Fe ; $\text{Br}_2/\text{anhy. AlCl}_3$; aq. KOH

Answer (2)

Sol.



61.



Choose the correct option for structures of A and B, respectively

- (1) $\text{H}_2\text{N}-\text{CH}-\text{COO}^-$ and $\text{H}_3\text{N}^+-\text{CH}-\text{COOH}$
 $\quad \quad \quad |$ $\quad \quad \quad |$
 $\quad \quad \quad \text{CH}(\text{CH}_3)_2$ $\quad \quad \quad \text{CH}(\text{CH}_3)_2$
 (2) $\text{H}_2\text{N}-\text{CH}-\text{COO}^-$ and $\text{H}_3\text{N}^+-\text{CH}-\text{COO}^-$
 $\quad \quad \quad |$ $\quad \quad \quad |$
 $\quad \quad \quad \text{CH}(\text{CH}_3)_2$ $\quad \quad \quad \text{CH}(\text{CH}_3)_2$
 (3) $\text{H}_3\text{N}^+-\text{CH}-\text{COO}^-$ and $\text{H}_3\text{N}^+-\text{CH}-\text{COOH}$
 $\quad \quad \quad |$ $\quad \quad \quad |$
 $\quad \quad \quad \text{CH}(\text{CH}_3)_2$ $\quad \quad \quad \text{CH}(\text{CH}_3)_2$
 (4) $\text{H}_3\text{N}^+-\text{CH}-\text{COOH}$ and $\text{H}_2\text{N}-\text{CH}-\text{COO}^-$
 $\quad \quad \quad |$ $\quad \quad \quad |$
 $\quad \quad \quad \text{CH}(\text{CH}_3)_2$ $\quad \quad \quad \text{CH}(\text{CH}_3)_2$

Answer (4)

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
(max. 1 each, 1 below)

1000+ 99
PERCENTILERS
& ABOVE

4000+ 95
PERCENTILERS
& ABOVE

100
Percentile
in
Physics
&
Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in
Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in
Physics
&
Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile




Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS





Sol. At pH = 2 \Rightarrow Medium is acidic so $-\text{NH}_2$ group will convert to $-\text{NH}_3^+$







\therefore At pH = 10 \Rightarrow medium is basic

So $-\text{COOH}$ group will convert to $-\text{COO}^-$




62. When  undergoes intramolecular aldol condensation, the major product formed is :




Answer (2)

Sol.




63. What is the correct IUPAC name of



- (1) 4-Ethylcyclopent-2-en-1-ol
- (2) 4-Ethyl-1-hydroxycyclopent-2-ene
- (3) 1-Ethyl-3-hydroxycyclopent-2-ene
- (4) 1-Ethylcyclopent-2-en-3-ol

Answer (1)



4-Ethylcyclopent-2-en-1-ol

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
AND ABOVE

1000+ 99
PERCENTILERS
AND ABOVE

4000+ 95
PERCENTILERS
AND ABOVE

100
Percentile
in
Physics
& Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in
Physics
& Chemistry



Harsh Jha
PSID: 00014863322

100
Percentile
in
Physics
& Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile
in
Physics
& Chemistry



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



64. Given below are two statements:

Statement I : H_2Se is more acidic than H_2Te

Statement II : H_2Se has higher bond enthalpy for dissociation than H_2Te

In the light of the above statement, choose the correct answer from the options given

- (1) Both Statement I and Statement II are false
 (2) Both Statement I and Statement II are true
 (3) Statement I is false but Statement II is true
 (4) Statement I is true but Statement II is false

Answer (3)

Sol. $\therefore \text{H}_2\text{Te}$ is more acidic than H_2Se

because of lesser bond dissociation energy of H_2Te

It can release H^+ more easily

$\text{pK}_{a_1} : \text{H}_2\text{Se}(3.89) > \text{H}_2\text{Te}(2.6)$

65. In a first order decomposition reaction, the time taken for the decomposition of reactant to one fourth and one eighth of its initial concentration are t_1 and t_2 (s), respectively. The ratio t_1/t_2 will be:

- (1) $\frac{4}{3}$ (2) $\frac{3}{2}$
 (3) $\frac{3}{4}$ (4) $\frac{2}{3}$

Answer (4)

Sol. $t_1 = t_{\frac{1}{4}} = \frac{1}{k} \ln \frac{A_0}{\frac{A_0}{4}} = \frac{1}{k} \ln 4$

$t_2 = t_{\frac{1}{8}} = \frac{1}{k} \ln \frac{A_0}{\frac{A_0}{8}} = \frac{1}{k} \ln 8$

$\frac{t_1}{t_2} = \frac{\ln 4}{\ln 8} = \frac{2 \ln 2}{3 \ln 2} = \frac{2}{3}$

66. On combustion of 0.210 g of an organic compound containing C, H and O gave 0.127 g H_2O and 0.307 g CO_2 . The percentage of hydrogen and oxygen in the given organic compound respectively are:

- (1) 53.41, 39.6 (2) 6.72, 39.87
 (3) 6.72, 53.41 (4) 7.55, 43.85

Answer (3)

Sol. Mass of organic compound = 0.210 g

Mass of water formed = 0.127 g

Mass of CO_2 formed = 0.307 g

Mass of hydrogen = $\frac{0.127 \times 2}{18} = 0.014$ g

Percentage of hydrogen = $\frac{0.014 \times 100}{0.210} = 6.72\%$

Mass of carbon = $\frac{0.307 \times 12}{44} = 0.084$ g

Percentage of carbon = $\frac{0.084 \times 100}{0.210} = 39.87\%$

Percentage of oxygen = 53.41%


67. The number of species from the following that are involved in sp^3d^2 hybridization is $[\text{Co}(\text{NH}_3)_6]^{3+}$, SF_6 , $[\text{CrF}_6]^{3-}$, $[\text{CoF}_6]^{3-}$, $[\text{Mn}(\text{CN})_6]^{3-}$ and $[\text{MnCl}_6]^{3-}$

- (1) 4 (2) 6
 (3) 5 (4) 3

Answer (4)

Sol. (i) $[\text{Co}(\text{NH}_3)_6]^{3+}$

$\text{Co}^{3+} \Rightarrow 3d^6 ; t_{2g}^6 e_g^0 d^2 sp^3$ hybridisation



sp^3d^2 hybridisation

(iii) $[\text{CrF}_6]^{3-}$

$\text{Cr}^{3+} \Rightarrow 3d^3 ; t_{2g}^3 e_g^0 d^2 sp^3$ hybridisation

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
(max. 1 each, 1 total)

1000+ 99
PERCENTILERS
5 ABOVE

4000+ 95
PERCENTILERS
9 ABOVE

100
Percentile
in
Physics
& Math



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in Physics
& Chemistry



Devya Rustagi
PSID: 00014768785


99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS






'A' shows positive Lassaigne's test for N and its molar mass is 121.

'B' gives effervescence with aq NaHCO₃


'C' gives fruity smell

Identify A, B and C from the following



Answer (4)


Sol. Compound (A) is likely to be amide which gives carboxylic acid (B) after hydrolysis. Compound (B) reacts with alcohol to give ester



SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

71. Consider the following half cell reaction



The reaction was conducted with the ratio of

$$\frac{[\text{Cr}^{3+}]^2}{[\text{Cr}_2\text{O}_7^{2-}]} = 10^{-6}$$

The pH value at which the EMF of


the half cell will become zero is _____. (nearest integer value)

[Given : standard half cell reduction potential

$$E^0_{\text{Cr}_2\text{O}_7^{2-}/\text{H}^+/\text{Cr}^{3+}} = 1.33\text{V}, \frac{2.303RT}{F} = 0.059\text{V}]$$

Answer (10)

Sol.



Using Nernst equation


$$E_{\text{Cr}_2\text{O}_7^{2-}/\text{H}^+/\text{Cr}^{3+}} = E^0_{\text{Cr}_2\text{O}_7^{2-}/\text{H}^+/\text{Cr}^{3+}} - \frac{2.303RT}{6F} \log \frac{[\text{Cr}^{3+}]^2}{[\text{Cr}_2\text{O}_7^{2-}][\text{H}^+]^{14}}$$

$$0 = E^0_{\text{Cr}_2\text{O}_7^{2-}/\text{H}^+/\text{Cr}^{3+}} - \frac{0.059}{6} \log(10^{-16}[\text{H}^+]^{-14})$$

$$0 = 1.33 + 0.059 - \frac{0.059 \times 14}{6} \text{pH}$$

$$\text{pH} = \frac{1.389 \times 6}{0.059 \times 14} = 10.10 \approx 10$$

72. The equilibrium constant for decomposition of H₂O(g)



is 8.0×10^{-3} at 2300 K and total pressure at equilibrium is 1 bar. Under this condition, the degree of dissociation (α) of water is _____ $\times 10^{-2}$ (nearest integer value)

[Assume α is negligible with respect to 1]

Answer (5)

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS
(max. 1 score, 1 bonus)

1000+ 99 PERCENTILERS
5 ABOVE

4000+ 95 PERCENTILERS
5 ABOVE

100
Percentile
in
Physics
& Math



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in
Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in
Physics
& Chemistry



Devya Rustagi
PSID: 00014768785


99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS





Initial mole 1 - -

Moles at equil. $1 - \alpha$ α $\frac{\alpha}{2}$

Equilibrium pressure = 1 bar

$$K_p = \frac{\left(\frac{2\alpha}{2+\alpha}\right)\left(\frac{\alpha}{2+\alpha}\right)^{\frac{1}{2}}}{2\left(\frac{1-\alpha}{2+\alpha}\right)} = 8.0 \times 10^{-3}$$

$$\frac{\alpha^2}{\sqrt{2}} = 8.0 \times 10^{-3}$$

$$\alpha = \left(8\sqrt{2} \times 10^{-3}\right)^{\frac{2}{3}}$$


$$= \left(2^{\frac{7}{2}} \times 10^{-3}\right)^{\frac{2}{3}} = 2^{\frac{7}{3}} \times 10^{-2} \approx 5 \times 10^{-2}$$

73. 20 mL of sodium iodide solution gave 4.74 g silver iodide when treated with excess of silver nitrate solution. The molarity of the sodium iodide solution is _____ M. (Nearest Integer Value)

(Given : Na = 23, I = 127, Ag = 108, N = 14, O = 16 g mol⁻¹)

Answer (1)

Sol. Let molarity of NaI solution be x M



$$\text{Moles of AgI formed} = \frac{4.74}{235} = 0.02$$

$$\text{Moles of NaI} = \frac{20 \times x}{1000} = 0.02x$$

$$0.02x = 0.02$$

$$x = 1$$

∴ Molarity of NaI solution = 1 M

74. The energy of an electron in first Bohr orbit of H-atom is -13.6 eV. The magnitude of energy value of electron in the first excited state of Be³⁺ is _____ eV (nearest integer value)

Answer (54)

Sol. E₁ of H-atom = -13.6 eV

$$\begin{aligned} E_2 \text{ of Be}^{3+} &= \frac{-13.6 \times Z^2}{n^2} \\ &= \frac{-13.6 \times (4)^2}{(2)^2} \\ &= -54.4 \text{ eV} \end{aligned}$$

$$|E_2| \text{ of Be}^{3+} = 54 \text{ eV}$$

75. Resonance in X₂Y can be represented as




The enthalpy of formation of X₂Y (X ≡ X(g) + $\frac{1}{2}$ Y = Y(g) → X₂Y(g)) is 80 kJ mol⁻¹.

The magnitude of resonance energy of X₂Y is _____ kJ mol⁻¹ (nearest integer value)

Given : Bond energies of X ≡ X, X = X, Y = Y and X = Y are 940, 410, 500 and 602 kJ mol⁻¹ respectively.

Valence X : 3, Y : 2

Answer (98)



$$[\Delta H_f(X_2Y)]_{\text{Actual}} = 80 \text{ kJ mol}^{-1}$$

$$\begin{aligned} [\Delta H_f(X_2Y)]_{\text{Theoretical}} &= 940 + \frac{1}{2}(500) - (410 + 602) \\ &= 1190 - 1012 \\ &= 178 \text{ kJ mol}^{-1} \end{aligned}$$

$$\begin{aligned} \text{Resonance energy} &= |(\Delta H_f)_{\text{Actual}} - (\Delta H_f)_{\text{Theoretical}}| \\ &= |80 - 178| = 98 \text{ kJ mol}^{-1} \end{aligned}$$



THE LEGACY OF SUCCESS CONTINUES

OUR JEE CHAMPIONS

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS
(max. 1 each, 1 below)

1000+ 99 PERCENTILERS
5 ABOVE

4000+ 95 PERCENTILERS
5 ABOVE



100 Percentile
Shreyas Lohiya
PSID: 00003389699



100 Percentile
Harsh Jha
PSID: 00014863322



99.99 Percentile
Devya Rustagi
PSID: 00014768785



99.99 Percentile
Amogh Bansal
PSID: 00014769016



Chirag Falor
4 Year Classroom
1 AIR
JEE (Adv.)
2020



Tanishka Kabra
4 Year Classroom
1 AIR-16 CBCL
JEE (Adv.)
2022



Sanvi Jain
4 Year Classroom
1 AIR-34 CBCL
JEE (Adv.)
2024