

## CHEMISTRY

### SECTION - A

**Multiple Choice Questions:** This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

**Choose the correct answer :**

31. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A):**  $\text{Cu}^{2+}$  in water is more stable than  $\text{Cu}^+$ .

**Reason (R):** Enthalpy of hydration for  $\text{Cu}^{2+}$  is much less than that of  $\text{Cu}^+$ .

In the light of the above statements, choose the **correct** answer from the options given below:

- (1) Both **(A)** and **(R)** are correct and **(R)** is the correct explanation of **(A)**
- (2) **(A)** is not correct but **(R)** is correct
- (3) **(A)** is correct but **(R)** is not correct
- (4) Both **(A)** and **(R)** are correct but **(R)** is not the correct explanation of **(A)**

**Answer (3)**

**Sol.**  $\text{Cu}^{2+}$  in water is more stable than  $\text{Cu}^+$  due to much higher hydration enthalpy of  $\text{Cu}^{2+}$  ion. Hence correct answer is option (3)

32. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A):**  $\alpha$ -halocarboxylic acid on reaction with dil  $\text{NH}_3$  gives good yield of  $\alpha$ -amino carboxylic acid whereas the yield of amines is very low when prepared from alkyl halides.


**Reason (R):** Amino acids exist in zwitter ion form in aqueous medium.

In the light of the above statements, choose the **correct** answer from the options given below:

- (1) **(A)** is not correct but **(R)** is correct
- (2) **(A)** is correct but **(R)** is not correct
- (3) Both **(A)** and **(R)** are correct and **(R)** is the correct explanation of **(A)**
- (4) Both **(A)** and **(R)** are correct but **(R)** is **not** the correct explanation of **(A)**

**Answer (3)**

**Sol. Statement-I:**



**Statement-II :** Reason is a correct statement as amino do exist as a zwitter ion. Reason is also a correct explanation.


33. Which element is not present in Nessler's reagent?

- (1) Potassium
- (2) Oxygen
- (3) Mercury
- (4) Iodine

**Answer (2)**


**Sol.** Nessler's reagent is  $\text{K}_2[\text{HgI}_4]$

34. All structures given below are of vitamin C. Most stable of them is:



**Answer (1)**

**Sol.** Most stable structure of vitamin(C) is :



35. Which one of the following sets of ions represents a collection of isoelectronic species?

(Given : Atomic Number : F : 9, Cl : 17, Na = 11, Mg = 12, Al = 13, K = 19, Ca = 20, Sc = 21)

- (1)  $\text{Ba}^{2+}$ ,  $\text{Sr}^{2+}$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$
- (2)  $\text{N}^{3-}$ ,  $\text{O}^{2-}$ ,  $\text{F}^-$ ,  $\text{S}^{2-}$
- (3)  $\text{K}^+$ ,  $\text{Cl}^-$ ,  $\text{Ca}^{2+}$ ,  $\text{Sc}^{3+}$
- (4)  $\text{Li}^+$ ,  $\text{Na}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$

**Answer (3)**

**Sol.** Isoelectronic species have same number of electrons.

$\text{K}^+$ ,  $\text{Cl}^-$ ,  $\text{Ca}^{2+}$  and  $\text{Sc}^{3+}$  all have 18 electrons, hence these are isoelectronic.

36. The correct order of bond enthalpy ( $\text{kJ mol}^{-1}$ ) is

- (1)  $\text{C}-\text{C} > \text{Si}-\text{Si} > \text{Sn}-\text{Sn} > \text{Ge}-\text{Ge}$
- (2)  $\text{Si}-\text{Si} > \text{C}-\text{C} > \text{Sn}-\text{Sn} > \text{Ge}-\text{Ge}$
- (3)  $\text{C}-\text{C} > \text{Si}-\text{Si} > \text{Ge}-\text{Ge} > \text{Sn}-\text{Sn}$
- (4)  $\text{Si}-\text{Si} > \text{C}-\text{C} > \text{Ge}-\text{Ge} > \text{Sn}-\text{Sn}$

**Answer (3)**

Bond	Bond energy ( $\text{kJ mol}^{-1}$ )
C-C	348
Si-Si	297
Ge-Ge	260
Sn-Sn	240

Correct answer will be (3)


37. The industrial activity held least responsible for global warming is

- (1) industrial production of urea
- (2) manufacturing of cement
- (3) steel manufacturing
- (4) Electricity generation in thermal power plants

**Answer (1)**


**Sol.** Industrial production of urea is least responsible for global warming.

38. The structures of major products A, B and C in the following reaction are sequence.



**Answer (1)**

**Sol.**






39. The starting material for convenient preparation of deuterated hydrogen peroxide ( $\text{D}_2\text{O}_2$ ) in laboratory is

- (1) 2-ethylanthraquinol
- (2) BaO
- (3) BaO<sub>2</sub>
- (4)  $\text{K}_2\text{S}_2\text{O}_8$


**Answer (4)**

**Sol.**  $\text{K}_2\text{S}_2\text{O}_8$  is used in the laboratory preparation of  $\text{D}_2\text{O}_2$





Answer (3)



41. Given below are two statements: one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A)** : An aqueous solution of KOH when used for volumetric analysis, its concentration should be checked before the use.


**Reason (R)** : On aging, KOH solution absorbs atmospheric  $\text{CO}_2$ .

In the light of the above statements, choose the **correct** answer from the options given below :

- (1) Both **(A)** and **(R)** are correct but **(R)** is **not** the correct explanation of **(A)**
- (2) Both **(A)** and **(R)** are correct and **(R)** is the correct explanation of **(A)**
- (3) **(A)** is correct but **(R)** is not correct
- (4) **(A)** is not correct but **(R)** is correct

Answer (2)

Sol. KOH absorbs  $\text{CO}_2$  get converted to  $\text{K}_2\text{CO}_3$



42. Given below are two statements :

**Statement I** : Sulphanilic acid gives esterification test for carboxyl group.

**Statement II** : Sulphanilic acid gives red colour in Lassaigne's test for extra element detection.

In the light of the above statements, choose the **most appropriate** answer from the options given below :

- (1) **Statement I** is incorrect but **Statement II** is correct
- (2) Both **Statement I** and **Statement II** are incorrect
- (3) **Statement I** is correct but **Statement II** is incorrect
- (4) Both **Statement I** and **Statement II** are correct

Answer (1)

Sol. Sulphanilic acid is p-amino benzene sulphonc acid



Since it contain both N and S so it give red colour in Lassaigne's test.



43. The complex cation which has two isomers is:

- (1)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]^+$  (2)  $[\text{Co}(\text{H}_2\text{O})_6]^{3+}$
- (3)  $[\text{Co}(\text{NH}_3)_5\text{NO}_2]^{2+}$  (4)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]^{2+}$

Answer (3)

Sol. Complex  $[\text{Co}(\text{NH}_3)_5\text{NO}_2]^{2+}$  will have two isomer one linked through N (Nitro) and one through O (Nitrite).


44. The graph which represents the following reaction is :




**Answer (2)**

**Sol.** Rate =  $K[(C_6H_5)_3C - Cl]$

The correct mechanism is  $S_N1$ .



45. In a reaction,




reagents 'X' and 'Y' respectively are :


- (1)  $(CH_3CO)_2O/H^+$  and  $(CH_3CO)_2O/H^+$
- (2)  $CH_3OH/H^+, \Delta$  and  $CH_3OH/H^+, \Delta$
- (3)  $(CH_3CO)_2O/H^+$  and  $CH_3OH/H^+, \Delta$
- (4)  $CH_3OH/H^+, \Delta$  and  $(CH_3CO)_2O/H^+$

**Answer (3)**

**Sol.**



46. In figure, a straight line is given for Freundlich Adsorption ( $y = 3x + 2.505$ ). The value of  $\frac{1}{n}$  and  $\log K$  are respectively.



- (1) 3 and 2.505
- (2) 0.3 and 0.7033
- (3) 0.3 and  $\log 2.505$
- (4) 3 and 0.7033

**Answer (1)**

**Sol.**  $\log \frac{x}{m} = \log k + \frac{1}{n} \log p$

On comparing, we get

$$\frac{1}{n} = 3 \Rightarrow n = 0.3 \text{ and } \log k = 2.505$$

47. The effect of addition of helium gas to the following reaction in equilibrium state, is



- (1) the equilibrium will go backward due to suppression of dissociation of  $PCl_5$
- (2) addition of helium will not affect the equilibrium
- (3) the equilibrium will shift in the forward direction and more of  $Cl_2$  and  $PCl_3$  gases will be produced
- (4) helium will deactivate  $PCl_5$  and reaction will stop

**Answer (3)**

**Sol.** If we consider addition of He gas at constant pressure, the reaction will shift in forward direction [As rigid container is not given]

48. For electron gain enthalpies of the elements denoted as  $\Delta_{eg}H$ , the incorrect option is

- (1)  $\Delta_{eg}H(Cl) < \Delta_{eg}H(F)$
- (2)  $\Delta_{eg}H(Se) < \Delta_{eg}H(S)$
- (3)  $\Delta_{eg}H(I) < \Delta_{eg}H(At)$
- (4)  $\Delta_{eg}H(Te) < \Delta_{eg}H(Po)$

**Answer (2)**

**Sol.**  $\Delta_{eg}H(Cl) = -349 \text{ kJ/mole}$   $\Delta_{eg}H(F) = -333 \text{ kJ/mole}$

$$\Delta_{eg}H(I) = -296 \text{ kJ/mole}$$

$$\Delta_{eg}H(Se) = -195 \text{ kJ/mole}$$

$$\Delta_{eg}H(S) = -200 \text{ kJ/mole}$$

$$\Delta_{eg}H(Te) = -190 \text{ kJ/mole}$$

$$\Delta_{eg}H(Po) = -174 \text{ kJ/mole}$$

Electron gain enthalpy of Se is less negative than that of sulphur.

49. Given below are two statements: one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A):** Gypsum is used for making fireproof wall boards.

**Reason (R):** Gypsum is unstable at high temperatures.

In the light of the above statements, choose the **correct** answer from the options given below

- (1) Both **(A)** and **(R)** are correct and **(R)** is the correct explanation of **(A)**
- (2) **(A)** is correct but **(R)** is not correct
- (3) Both **(A)** and **(R)** are correct but **(R)** is not the correct explanation of **(A)**
- (4) **(A)** is not correct but **(R)** is correct

**Answer (3)**

**Sol.** Both statements are correct. However, II<sup>nd</sup> statement has no relation with I<sup>st</sup> Statement.

50. O – O bond length in H<sub>2</sub>O<sub>2</sub> is X than the O – O bond length in F<sub>2</sub>O<sub>2</sub>. The O – H bond length in H<sub>2</sub>O<sub>2</sub> is Y than that of the O – F bond in F<sub>2</sub>O<sub>2</sub>.

Choose the correct option for X and Y from those given below

- (1) X – shorter, Y - shorter
- (2) X – shorter, Y - longer
- (3) X – longer, Y - shorter
- (4) X – longer, Y - longer

**Answer (3)**


**Sol.** X – longer [because of more p-character in O – F bond]

Y – shorter [size of H is very small as compared to F]

## SECTION - B

**Numerical Value Type Questions:** This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g. 06.25, 07.00, –00.33, –00.30, 30.27, –27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.


51. Testosterone, which is a steroidal hormone, has the following structure.



The total number of asymmetric carbon atom/s in testosterone is \_\_\_\_\_.

**Answer (6)**

**Sol.**



The total number of asymmetric carbon atoms in testosterone is 6.

52. A metal M crystallizes into two lattices: face centred cubic (fcc) and body centred cubic (bcc) with unit cell edge length of 2.0 and 2.5 Å respectively. The ratio of densities of lattices fcc to bcc for the metal M is \_\_\_\_\_.

(Nearest integer)

**Answer (4)**

**Sol.**  $d_1$ , Density of fcc lattice of metal  $M = \frac{4 \times M}{N_0(a_{\text{fcc}})^3}$

$d_2$ , Density of bcc lattice of metal  $M = \frac{2 \times M}{N_0(a_{\text{bcc}})^3}$

$$\frac{d_1}{d_2} = \frac{4}{2} \left( \frac{a_{\text{bcc}}}{a_{\text{fcc}}} \right)^3 = 2 \left( \frac{2.5}{2} \right)^3 = 3.90 \approx 4$$

53.  $A \rightarrow B$

The above reaction is of zero order. Half life of this reaction is 50 min. The time taken for the concentration of A to reduce to one-fourth of its initial value is \_\_\_\_\_ min.

**Answer (75)**

**Sol.**  $A \xrightarrow{a-x} B \xrightarrow{x}$  (Zero Order reaction)

$$a - x = \frac{a}{4} \Rightarrow x = \frac{3a}{4}$$

$$t_{1/2} = \frac{a}{2K} = 50 \text{ min.} \Rightarrow \frac{a}{K} = 100 \text{ min.}$$

$$t = \frac{x}{K} = \frac{3a}{4K} = 75 \text{ min.}$$

54. 0.3 g of ethane undergoes combustion at  $27^\circ\text{C}$  in a bomb calorimeter. The temperature of calorimeter system (including the water) is found to rise by  $0.5^\circ\text{C}$ . The heat evolved during combustion of ethane at constant pressure is \_\_\_\_\_  $\text{kJ mol}^{-1}$ .


(Nearest integer)

[Given: The heat capacity of the calorimeter system is  $20 \text{ kJ K}^{-1}$ ,  $R = 8.3 \text{ JK}^{-1} \text{ mol}^{-1}$ .

Assume ideal gas behaviour.

Atomic mass of C and H are 12 and 1  $\text{g mol}^{-1}$  respectively]

**Answer (1006)**



No. of moles of ethane =  $\frac{0.3}{30} = 0.01$

Heat evolved in Bomb calorimeter =  $20 \times 0.5$   
= 10 kJ

$$\Delta U = -\frac{10}{0.01} = -1000 \text{ kJ mol}^{-1}$$

$$\Delta H = \Delta U + \Delta n_g RT$$

$$= -1000 + (-2.5) \times \frac{8.3 \times 300}{1000}$$

$$= -1000 - 6.225$$


$$= -1006.225$$

$$|\Delta H| \approx 1006 \text{ kJ mol}^{-1}$$

55. The spin only magnetic moment of  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$  complexes is \_\_\_\_\_ B.M. (Nearest integer)

(Given: Atomic no. of Mn is 25)

**Answer (6)**



No. of unpaired electrons = 5


$$\mu = \sqrt{35} \text{ BM} = 6 \text{ BM}$$

56. 20% of acetic acid is dissociated when its 5 g is added to 500 mL of water. The depression in freezing point of such water is \_\_\_\_\_  $\times 10^{-3}^\circ\text{C}$ .

Atomic mass of C, H and O are 12, 1 and 16 a.m.u. respectively.

[Given : Molal depression constant and density of water are  $1.86 \text{ K kg mol}^{-1}$  and  $1 \text{ g cm}^{-3}$  respectively.

**Answer (372)**



$$i = 1.2$$

$$[\text{CH}_3\text{COOH}] = \frac{5}{60 \times 0.5} = \frac{5}{30} \text{ M}$$

$$\Delta T_f = i K_f m$$

$$1.2 \times 1.86 \times \frac{5}{30} = 0.372^\circ\text{C}$$

$$= 372 \times 10^{-3}^\circ\text{C}$$



57.  $1 \times 10^{-5}$  M  $\text{AgNO}_3$  is added to 1 L of saturated solution of  $\text{AgBr}$ . The conductivity of this solution at 298 K is \_\_\_\_\_  $\times 10^{-8} \text{ S m}^{-1}$ .


[Given :  $K_{\text{SP}}(\text{AgBr}) = 4.9 \times 10^{-3}$  at 298 K

$$\lambda_{\text{Ag}^+}^0 = 6 \times 10^{-3} \text{ S m}^2 \text{ mol}^{-1}$$

$$\lambda_{\text{Br}^-}^0 = 8 \times 10^{-3} \text{ S m}^2 \text{ mol}^{-1}$$

$$\lambda_{\text{NO}_3^-}^0 = 7 \times 10^{-3} \text{ S m}^2 \text{ mol}^{-1}]$$

**Answer (13039.2)**



$$x(x + 10^{-5}) = 4.9 \times 10^{-13}$$

$$x \approx 4.9 \times 10^{-8} \text{ M}$$

$$\lambda_{\text{Ag}^+}^0 = 6 \times 10^{-3} \text{ S cm}^2 \text{ mol}^{-1}$$

$$\lambda_{\text{Br}^-}^0 = 8 \times 10^{-3} \text{ S cm}^2 \text{ mol}^{-1}$$

$$\lambda_{\text{NO}_3^-}^0 = 7 \times 10^{-3} \text{ S cm}^2 \text{ mol}^{-1}$$

$$K_{\text{solution}} = K_{\text{Ag}^+} + K_{\text{Br}^-} + K_{\text{NO}_3^-}$$

$$= 6 \times 10^{-3} \times 10^{-5} \times 10^3 + 8 \times 10^{-3} \times 4.9 \times 10^{-8} \times 10^3 + 7 \times 10^{-3} \times 10^{-5} \times 10^3$$

$$= (6000 + 39.2 + 7000) \times 10^{-8}$$

$$= 13039.2 \times 10^{-8} \text{ Sm}^{-1}$$

58. Among the following, the number of tranquilizer/s is /are \_\_\_\_\_.

- A. Chloroliazepoxide
- B. Veronal
- C. Valium
- D. Salvarsan

**Answer (3)**

**Sol.** Chloroliazepoxide

Veronal

Valium

Salvarsan is an antibiotic

59. The molarity of a 10% (v/v) solution of di-bromine solution in  $\text{CCl}_4$  (carbon tetrachloride) is 'x'.

$$x = \text{_____} \times 10^{-2} \text{ M. (Nearest integer)}$$

[Given : molar mass of  $\text{Br}_2 = 160 \text{ g mol}^{-1}$

atomic mass of C = 12  $\text{g mol}^{-1}$

atomic mass of Cl = 35.5  $\text{g mol}^{-1}$

density of dibromine = 3.2  $\text{g cm}^{-3}$

density of  $\text{CCl}_4 = 1.6 \text{ g cm}^{-3}$ ]

**Answer (139)**

**Sol.** Mass of 10 mL of  $\text{Br}_2 = 10 \times 3.2 = 32 \text{ gm}$

Mass of 90 mL of  $\text{CCl}_4 = 90 \times 1.6 = 144 \text{ gm}$

$$\text{Molality of } \text{Br}_2 \text{ solution in } \text{CCl}_4 = \frac{32 \times 1000}{160 \times 144}$$

$$= 1.39 \text{ M}$$

$$= 139 \times 10^{-2}$$

60. Among following compounds, the number of those present in copper matte is \_\_\_\_\_.

- A.  $\text{CuCO}_3$
- B.  $\text{Cu}_2\text{S}$
- C.  $\text{Cu}_2\text{O}$
- D.  $\text{FeO}$

**Answer (3)**

**Sol.** Copper matte contains

$\text{Cu}_2\text{S}$ ,  $\text{Cu}_2\text{O}$ ,  $\text{FeO}$