



Sol.  $I = \int_{-2}^2 \frac{9x^2}{1+5^x} dx \dots(1)$

$I = \int_{-2}^2 \frac{9x^2}{1+5^{-x}} dx \dots(2)$

Adding (1) and (2)

$$2I = \int_{-2}^2 \left( \frac{9x^2}{1+5^x} + \frac{5^x \cdot 9x^2}{1+5^x} \right) dx$$

$$2I = \int_{-2}^2 \frac{9x^2}{1+5^x} (1+5^x) dx$$

$$2I = \int_{-2}^2 9x^2 dx$$

$$2I = 9 \left( \frac{x^3}{3} \right)^2$$

$$2I = 9 \left( \frac{8}{3} + \frac{8}{3} \right) = 48$$

$I = 24$

5. If the mean and variance of eight observations  $a, b, 8, 12, 10, 6, 4, 15$ , is 9 and 9.25 respectively. Then  $a + b + ab$  is equal to

- (1) 76
- (2) 83
- (3) 79
- (4) 93

Answer (4)

Sol. Mean = 9 =  $\frac{a+b+8+12+10+6+4+15}{8}$

$\Rightarrow a + b + 55 = 72 \Rightarrow a + b = 17$

$\frac{a^2 + b^2 + 64 + 144 + 100 + 36 + 16 + 225}{8} - 9^2 = 9.25$

$a^2 + b^2 + 585 - 8 \cdot 9^2 = 74$

$\Rightarrow a^2 + b^2 = 137$

$\Rightarrow (a + b)^2 - 2ab = 137$

$\Rightarrow 2ab = 289 - 137 \Rightarrow ab = 76$

$\Rightarrow a + b + ab = 17 + 76 = 93$

6.  $4 \int_0^1 \frac{1}{\sqrt{3+x^2} + \sqrt{1+x^2}} dx - 3 \ln \sqrt{3}$  is equal to

- (1)  $3 - \sqrt{2} + \ln(\sqrt{2} + 1)$
- (2)  $2 + \sqrt{2} - \ln(\sqrt{3} + 1)$
- (3)  $2 - \sqrt{2} - \ln(\sqrt{2} + 1)$
- (4)  $2 - \sqrt{3} - \ln(\sqrt{3} + 1)$

Answer (3)

Sol.  $I = 4 \int_0^1 \frac{1}{\sqrt{3+x^2} + \sqrt{1+x^2}} dx$

$= 2 \int_0^1 \sqrt{3+x^2} - \sqrt{1+x^2} dx$

$= 2 \left[ \int_0^1 \sqrt{3+x^2} dx - \int_0^1 \sqrt{1+x^2} dx \right]$

$= 2 \left[ \left( \frac{1}{2} x \sqrt{x^2+3} + \frac{3}{2} \ln \left| \sqrt{3+x^2} + x \right| \right) - \right.$

$\left. \left( \frac{1}{2} x \sqrt{1+x^2} + \frac{1}{2} \ln \left| \sqrt{1+x^2} + x \right| \right) \right]_0^1$

$= 2 \left[ \left( 1 + \frac{3}{2} \ln 3 - \frac{3}{2} \ln \sqrt{3} \right) - \left( \frac{\sqrt{2}}{2} + \frac{1}{2} \ln(\sqrt{2} + 1) \right) \right]$

$= 2 \left( 1 + \frac{3}{4} \ln 3 - \frac{1}{\sqrt{2}} - \frac{1}{2} \ln(\sqrt{2} + 1) \right)$

$= 3 \ln \sqrt{3} + 2 - \sqrt{2} - \ln(\sqrt{2} + 1)$

$I - 3 \ln \sqrt{3} = 2 - \sqrt{2} - \ln(\sqrt{2} + 1)$

7. If  $y = \cos\left(\frac{\pi}{3} + \cos^{-1}\left(\frac{x}{2}\right)\right)$ , then which of the following is true.

- (1)  $x^2 - 2xy + 8y^2 = 2$
- (2)  $x^2 - 2xy + 4y^2 = 3$
- (3)  $x^2 - 3xy + 4y^2 = 3$
- (4)  $x^2 - 5xy + 4y^2 = 8$

Answer (2)

Sol.  $\therefore y = \cos\left(\frac{\pi}{3} + \cos^{-1}\frac{x}{2}\right)$

## THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE  
TOPPERS

70+ 100  
PERCENTILES  
in Physics

1000+ 99 PERCENTILES  
in Chemistry

4000+ 95 PERCENTILES  
in Biology

100  
Percentile  
in Physics  
& Maths



**Shreyas Lohiya**  
PSID: 00003389699

100  
Percentile  
in Physics



**Harsh Jha**  
PSID: 00014863322

100  
Percentile  
in Physics  
& Chemistry



**Devya Rustagi**  
PSID: 00014768785

99.99  
Percentile



**Amogh Bansal**  
PSID: 00014769016

OUR JEE CHAMPIONS



**Chirag Falor**  
4 Year Classroom  
1 AIR  
JEE (Adv.)  
2020



**Tanishka Kabra**  
4 Year Classroom  
1 AIR-16 CRL  
JEE (Adv.)  
2022



**Sanvi Jain**  
4 Year Classroom  
1 AIR-34 CRL  
JEE (Main)  
2024

$$y = \cos \frac{\pi}{3} \cdot \cos \left( \cos^{-1} \frac{x}{2} \right) - \sin \frac{\pi}{3} \cdot \sin \left( \cos^{-1} \frac{x}{2} \right)$$

$$y = \frac{1}{2} \cdot \frac{x}{2} - \frac{\sqrt{3}}{2} \cdot \sqrt{1 - \frac{x^2}{4}}$$

$$4y = x - \sqrt{3} \sqrt{4 - x^2}$$

$$(4y - x)^2 = 3(4 - x^2)$$

$$16y^2 + x^2 - 8xy = 12 - 3x^2$$

$$4x^2 - 8xy + 16y^2 = 12$$

$$\therefore x^2 - 2xy + 4y^2 = 3$$

8. The image of the point (1, 0, 3) about the line passing through  $\vec{a} = 3\hat{i} + 2\hat{j} - \hat{k}$  and whose direction ratios are  $\vec{r} = 4\hat{i} + 2\hat{j} - \hat{k}$  is

(1)  $\left( \frac{-23}{21}, \frac{20}{21}, \frac{-73}{21} \right)$       (2)  $\left( \frac{1}{21}, \frac{-23}{21}, \frac{-31}{21} \right)$


(3)  $\left( \frac{1}{21}, \frac{21}{23}, \frac{-30}{21} \right)$       (4)  $\left( \frac{3}{21}, \frac{7}{21}, \frac{-5}{21} \right)$

**Answer (1)**

**Sol.**  $\vec{a} : 3\hat{i} + 2\hat{j} - \hat{k}$

Dr:  $4\hat{i} + 2\hat{j} - \hat{k}$

$$L: \frac{x-3}{4} = \frac{y-2}{2} = \frac{z+1}{-1}$$



Any point on line L:  $P' (4\lambda + 3, 2\lambda + 2, -\lambda - 1)$

$$PP' \cdot \vec{n} = 0$$

$$\Rightarrow 4(4\lambda + 2) + 2(2\lambda + 2) + (-\lambda - 4)(-1) = 0$$

$$16\lambda + 8 + 4\lambda + 4\lambda + 4 = 0$$

$$21\lambda + 16 = 0$$

$$\lambda = \frac{-16}{21}$$

$$\therefore P' \left( \frac{-1}{21}, \frac{10}{21}, \frac{-5}{21} \right)$$

Let image of point P be  $[a, b, c]$

$$\therefore \frac{a+1}{2} = \frac{-1}{21} \Rightarrow a = \frac{-23}{21}$$

$$\frac{b+0}{2} = \frac{10}{21} \Rightarrow b = \frac{20}{21}$$

$$\frac{c+3}{2} = \frac{-5}{21} \Rightarrow c = \frac{-73}{21}$$

$$\therefore \text{image will be } \left( \frac{-23}{21}, \frac{20}{21}, \frac{-73}{21} \right)$$

9. If the curve  $x^2 = 4y$  intersects the line  $y = 2(x + 6)$  at  $(a, b)$  in 2<sup>nd</sup> quadrant, then  $\int_a^b \frac{x^4}{1+5^x} dx$  is

(1)  $\frac{512}{5}$       (2)  $\frac{1024}{5}$

(3)  $\frac{32}{5}$

(4)  $\frac{16}{5}$

**Answer (2)**

**Sol.**  $x^2 = 4y$

$$y = 2(x + 6)$$

$$x^2 = 8(x + 6)$$

$$x^2 - 8x - 48 = 0$$

$$(x + 4)(x - 12) = 0$$

$$\Rightarrow x = -4 (\because x < 0)$$

$$\therefore y = 4$$

$$\Rightarrow (a, b) \equiv (-4, 4)$$

$$I = \int_a^b \frac{x^4}{1+5^x} dx$$

$$= \int_{-4}^4 \frac{x^4}{1+5^x} dx$$

## THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE  
TOPPERS

70+ 100  
PERCENTILES  
in Physics

1000+ 99  
PERCENTILES  
in Maths

4000+ 95  
PERCENTILES  
in Biology

100  
Percentile  
in Physics  
& Maths



**Shreyas Lohiya**  
PSID: 00003389699

100  
Percentile  
in Physics



**Harsh Jha**  
PSID: 00014863322

100  
Percentile  
in Physics  
& Chemistry



**Devya Rustagi**  
PSID: 00014768785

99.99  
Percentile



**Amogh Bansal**  
PSID: 00014769016

OUR JEE CHAMPIONS



$$= \int_0^4 \left( \frac{x^4}{1+5^x} + \frac{x^4}{1+5^{-x}} \right) dx$$

$$= \int_0^4 x^4 dx = \frac{x^5}{5} = \frac{4^5}{5} = \frac{1024}{5}$$

10. If  $\lim_{x \rightarrow 0} \frac{\cos 2x + a \cos^4 x - b}{x^4} = L$  (finite)  
then  $a + b$  equals to  
(1) -1 (2) 0  
(3) 2 (4) 3

Answer (1)

Sol.  $\lim_{x \rightarrow 0} \frac{\cos 2x + a \cos^4 x - b}{x^4} = L$

$$\lim_{x \rightarrow 0} \frac{2 \cos^2 x - 1 + a \cos^4 x - b}{x^4} = L \dots (1)$$

To get the finite value,

$$1 + a - b = 0$$

$$\Rightarrow a = b - 1 \dots (2)$$

Apply L Hospital

$$\lim_{x \rightarrow 0} \frac{4 \cos x (-\sin x) + 4a \cos^3 x (-\sin x)}{4x^3}$$

$$\lim_{x \rightarrow 0} \frac{4 \cos x + 4a \cos^3 x}{4x^3} \left( \frac{-\sin x}{x} \right)$$

To get the finite value,  $a = -1$

Also from (1)

$$b = 0$$

$$\therefore a + b = -1$$

11. If the sum of series  $\frac{1}{1+4 \cdot 1^4} + \frac{2}{1+4 \cdot 2^4} + \frac{3}{1+4 \cdot 3^4} + \dots + \frac{10}{1+4 \cdot 10^4}$  is  $\frac{m}{n}$ , where  $m$  and  $n$  are natural coprime numbers, then  $(m + n)$  is  
(1) 289 (2) 276  
(3) 225 (4) 389

Answer (2)

Sol.  $T_r = \frac{r}{1+4 \cdot r^4} = \frac{r}{4r^4 + 4r^2 + 1 - 4r^2} =$

$$\frac{r}{(2r^2 + 1)^2 - (2r)^2} = \frac{r}{(2r^2 - 2r + 1)(2r^2 + 2r + 1)}$$

$$T_r = \frac{1}{4} \left[ \frac{(2r^2 + 2r + 1) - (2r^2 - 2r + 1)}{(2r^2 - 2r + 1)(2r^2 + 2r + 1)} \right]$$

$$= \frac{1}{4} \left[ \frac{1}{r^2 + (r-1)^2} - \frac{1}{r^2 + (r+1)^2} \right]$$

$$\sum_{r=1}^{10} T_r = \frac{1}{4} \left[ \frac{1}{0^2 + 1^2} - \frac{1}{1^2 + 2^2} + \frac{1}{1^2 + 2^2} - \frac{1}{2^2 + 3^2} + \frac{1}{2^2 + 3^2} - \frac{1}{3^2 + 4^2} + \dots - \frac{1}{10^2 + 11^2} \right]$$

$$= \frac{1}{4} \left[ \frac{1-1}{221} \right] = \frac{220}{4 \times 221} = \frac{55}{221}$$

12. A bag is Randomly selected, If drawn ball is red, then probability that ball is selected from bag-I is  $p$ . If ball drawn is green then probability that ball is selected from bag-III is  $q$ . Then  $\frac{1}{p} + \frac{1}{q}$  equals to

	Red	Blue	Green
Bag-I	3	3	4
Bag-II	4	3	3
Bag-III	5	2	3

- (1)  $\frac{22}{3}$  (2)  $\frac{22}{5}$   
(3)  $\frac{11}{3}$  (4)  $\frac{11}{5}$

Answer (1)

Sol.  $p(B_1 / R) = \frac{p(B_1) \cdot p(R / B_1)}{p(R)}$

$$= \frac{\frac{1}{3} \times \frac{3}{10}}{\frac{1}{3} \times \frac{3}{10} + \frac{1}{3} \times \frac{4}{10} + \frac{1}{3} \times \frac{5}{10}} = \frac{1}{4} = p$$

## THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths



**Shreyas Lohiya**  
PSID: 00003389699

100 Percentile in Physics



**Harsh Jha**  
PSID: 00014863322

100 Percentile in Physics & Chemistry



**Devya Rustagi**  
PSID: 00014768785

99.99 Percentile



**Amogh Bansal**  
PSID: 00014769016

OUR JEE CHAMPIONS




$$p(B_3 / G) = \frac{p(B_3) \cdot p(G / B_3)}{p(G)}$$

$$= \frac{\frac{1}{3} \times \frac{3}{10}}{\frac{1}{3} \times \frac{3}{10} + \frac{1}{3} \times \frac{3}{10} + \frac{1}{3} \times \frac{4}{10}} = \frac{3}{10} = q$$


$$\frac{1}{p} + \frac{1}{q} = 4 + \frac{10}{3} = \frac{22}{3}$$

13. In the given figure, number of ways to fill  $a, b, c, d$  and  $e$  into boxes such that no row is empty and at most one letter is filled in one box, is



- (1) 5670  
(2) 5760  
(3) 5880  
(4) 720

**Answer (2)**



Let  $x, y, z$  be the number of box which are filled

$$\Rightarrow 1 \leq x \leq 3, 1 \leq y \leq 3, 1 \leq z \leq 2$$

$x$	$y$	$z$	Number of ways
3	1	1	${}^3C_3 \cdot {}^3C_1 \cdot {}^2C_1 = 6$
2	2	1	${}^3C_2 \cdot {}^3C_2 \cdot {}^2C_1 = 18$
1	3	1	${}^3C_1 \cdot {}^3C_3 \cdot {}^2C_1 = 6$
2	1	2	${}^3C_2 \cdot {}^3C_1 \cdot {}^2C_2 = 9$
1	2	2	${}^3C_1 \cdot {}^3C_2 \cdot {}^2C_2 = 9$

Total ways = (48) to fill boxes

Now to arrange  $a, b, c, d$  and  $e$

Number of ways will be  $48 \cdot 5! = 5760$

14.  
15.  
16.  
17.  
18.  
19.  
20.

**SECTION - B**

**Numerical Value Type Questions:** This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

21. Total number of terms in an AP are even. Sum of odd terms is 24 and sum of even terms is 30. Last term exceeds the first term by  $\frac{21}{2}$ . Then the total number of terms is

**Answer (8)**

Sol. Let the number of terms be  $2n$

$$T_1 + T_3 + T_5 \dots T_{2n-1} = 24$$

$$T_2 + T_4 + T_6 \dots T_{2n} = 30$$

$$\frac{T_2 - T_1 + (T_4 - T_3) + \dots + (T_{2n} - T_{2n-1})}{nd = 6}$$

$$nd = 6$$

$$(a + (2n + 1)d) - a = \frac{21}{2}$$

$$\Rightarrow 2nd - d = \frac{21}{2}$$

$$\Rightarrow 12 - \frac{21}{2} = d$$

$$\Rightarrow d = \frac{3}{2}$$

$$\therefore n = 4$$

$$\therefore \text{Total terms} = 8$$

22. If  $\frac{dy}{dx} + 2y \sec^2 x = 2 \sec^2 x + 3 \tan x \sec^2 x$  and

$$f(0) = \frac{5}{4}. \text{ Then the value of } 12 \left( y \left( \frac{\pi}{4} \right) - \frac{1}{e^2} \right) \text{ equal to}$$

**Answer (21)**

**THE LEGACY OF SUCCESS CONTINUES**

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths



**Shreyas Lohiya**  
PSID: 00003389699

100 Percentile in Physics & Chemistry



**Harsh Jha**  
PSID: 00014863322

100 Percentile in Physics & Chemistry



**Devya Rustagi**  
PSID: 00014768785

99.99 Percentile



**Amogh Bansal**  
PSID: 00014769016

OUR JEE CHAMPIONS



Sol.  $\frac{dy}{dx} + 2y\sec^2 x = 2\sec^2 x + 3\tan x \sec^2 x$

I.F. =  $e^{\int 2\sec^2 x dx}$

I.F. =  $e^{2\tan x}$

$y \cdot e^{2\tan x} = \int e^{2\tan x} (2 + 3\tan x) \sec^2 x dx$

Put  $\tan x = u$

$\sec^2 x dx = du$

$y \cdot e^{2u} = \int e^{2u} (2 + 3u) du$

$y \cdot e^{2u} \Rightarrow \frac{2e^{2u}}{2} + 3 \int e^{2u} \cdot u du$

$y \cdot e^{2u} = e^{2u} + 3 \left[ \frac{ue^{2u}}{2} - \int \frac{e^{2u}}{2} \right]$

$ye^{2u} = e^{2u} + 3 \left[ \frac{ue^{2u}}{2} - \frac{e^{2u}}{4} \right] + C$

$ye^{2\tan x} = e^{2\tan x} + 3 \left[ \frac{\tan x e^{2\tan x}}{2} - \frac{e^{2\tan x}}{4} \right] + C$

$F(0) = \frac{5}{4}$

$\frac{5}{4} = 1 - \frac{3}{4} + C$

$\frac{5}{4} - \frac{1}{4} = C$

$1 = C$

$y = 1 + 3 \left( \frac{\tan x}{2} - \frac{1}{4} \right) + 1 \cdot e^{-2\tan x}$

$y \left( \frac{\pi}{4} \right) = 1 + 3 \left( \frac{1}{2} - \frac{1}{4} \right) + \frac{1}{e^2}$

$y \left( \frac{\pi}{4} \right) = \frac{7}{4} + \frac{1}{e^2}$

$12 \left( y \left( \frac{\pi}{4} \right) - \frac{1}{e^2} \right) = 12 \left( \frac{7}{4} + \frac{1}{e^2} - \frac{1}{e^2} \right) = 21$

23. If the non-zero  $3 \times 3$  matrix  $A$  satisfies  $A^2(A - 4I) - 4(A - I) = 0$  and if  $A^5 = \alpha A^2 + \beta A + \gamma I$ , where  $I$  is  $3 \times 3$  identity matrix, then  $\alpha + \beta + \gamma$  is equal to

Answer (76)

Sol.  $A^2(A - 4I) - 4(A - I) = 0$

$A^3 - 4A^2 - 4A + 4I = 0$

Multiple by  $A$

$A^4 = 4A^3 + 4A^2 - 4A$

$= 4(4A^2 + 4A - 4I) + 4A^2 - 4A$

$= 20A^2 + 12A - 16I$

Multiple again by  $A$

$\Rightarrow A^5 = 20A^3 + 12A^2 - 16A$

$= 20(4A^2 + 4A - 4I) + 12A^2 - 16A$


$= 92A^2 + 64A - 80I = \alpha A^2 + \beta A + \gamma I$

$\Rightarrow \alpha = 92, \beta = 64, \gamma = -80 \Rightarrow \alpha + \beta + \gamma = 76$

24. If  $PQ$  be the focal chord of a parabola  $y^2 = 16x$  such that  $P(1, -4)$  and  $\frac{PF}{QF} = \frac{m}{n}$ , ( $F$  is focus) where  $m$  and  $n$  are coprime natural numbers, then  $m^2 + n^2$  is

Answer (17)

Sol.



$y^2 = 16x$

$\Rightarrow 4a = 16 \Rightarrow a = 4$

$Q \equiv (at_2^2, 2at_2)$

$\equiv (4t_2^2, 8t_2)$

## THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths



**Shreyas Lohiya**  
PSID: 00003389699

100 Percentile in Physics



**Harsh Jha**  
PSID: 00014863322

100 Percentile in Physics & Chemistry



**Devya Rustagi**  
PSID: 00014768785

99.99 Percentile



**Amogh Bansal**  
PSID: 00014769016

OUR JEE CHAMPIONS



**Chirag Falor**  
4 Year Classroom  
1 AIR JEE (Adv.) 2020



**Tanishka Kabra**  
4 Year Classroom  
1 AIR-16 CRL JEE (Adv.) 2022  
ALL INDIA FEMALE TOPPER



**Sanvi Jain**  
4 Year Classroom  
1 AIR-34 CRL JEE (Main) 2024  
ALL INDIA FEMALE TOPPER

$$P \equiv (4t_1^2, 8t_1)$$

$$4t_1^2 = 1, 8t_1 = -4 \Rightarrow t_1 = \frac{-1}{2}$$

since  $P$  and  $Q$  are ends points of focal chord

$$t_1 t_2 = 1 \Rightarrow t_2 = 2$$

$$\Rightarrow Q \equiv (16, 16)$$

$$\Rightarrow PF = \sqrt{3^2 + 4^2}, FQ = \sqrt{12^2 + 16^2}$$

$$\Rightarrow \frac{PF}{FQ} = \frac{5}{20} = \frac{1}{4} = \frac{m}{n}$$

$$\Rightarrow m^2 + n^2 = 17$$

25.



## THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE  
TOPPERS

70+ 100  
PERCENTILERS  
PHYSICS & CHEMISTRY

1000+ 99  
PERCENTILERS  
& ABOVE

4000+ 95  
PERCENTILERS  
& ABOVE

100  
Percentile  
in  
Physics  
& Maths



**Shreyas Lohiya**  
PSID: 00003389699

100  
Percentile  
in  
Physics



**Harsh Jha**  
PSID: 00014863322

100  
Percentile  
in  
Physics  
& Chemistry



**Devya Rustagi**  
PSID: 00014768785

99.99  
Percentile



**Amogh Bansal**  
PSID: 00014769016

### OUR JEE CHAMPIONS



INTRODUCING

# FIRST-EVER

## JEE MAIN SCORE CALCULATOR

Paste the sheet URL here

1

**OPEN YOUR  
RESPONSE SHEET  
ON THE JEE MAIN SITE**

2

**COPY THE LINK FROM  
THE BROWSER'S  
ADDRESS BAR**

3

**PASTE IT INTO THE  
REQUIRED FIELD IN  
THE CALCULATOR**

**CHECK NOW FOR FREE**

Curious about your JEE Main score?

Get an instant estimate with our JEE Main Score Calculator -

<https://bit.ly/4jau6gR>

## THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE  
TOPPERS

70+ 100  
PERCENTILERS  
IN PHYSICS

1000+ 99  
PERCENTILERS  
IN ABOVE

4000+ 95  
PERCENTILERS  
IN ABOVE

100  
Percentile  
in Physics  
& Maths



**Shreyas Lohiya**  
PSID: 00003389699

100  
Percentile  
in Physics



**Harsh Jha**  
PSID: 00014863322

100  
Percentile  
in Physics  
& Chemistry



**Devya Rustagi**  
PSID: 00014768785

99.99  
Percentile



**Amogh Bansal**  
PSID: 00014769016

OUR JEE CHAMPIONS



**Chirag Falor**  
4 Year Classroom  
**1** AIR  
JEE (Adv.)  
2020



**Tanishka Kabra**  
4 Year Classroom  
**1** AIR-16 CRL  
JEE (Adv.)  
2022



**Sanvi Jain**  
4 Year Classroom  
**1** AIR-34 CRL  
JEE (Main)  
2024