





$$67.05 \text{ mL weighs } \frac{28}{22400} \times 67.05$$

0.0838 g


$$\% \text{ of N} = \frac{0.0838}{0.4} \times 100 = 20.95$$

4. Which of the following reagent is used to prepare tribromoaniline?



Answer (4)

Sol.



Step (1) involves reduction of nitrobenzene to aniline and step (2) involves bromination of aniline to give 2, 4, 6-tribromoaniline

5. Match the following list-I with list-II :

List-I (Groups)

List-II (Elements)

- | | |
|-----------------|----------|
| (A) Pnictogens | (I) Rn |
| (B) Chalcogens | (II) At |
| (C) Halogens | (III) Te |
| (D) Noble gases | (IV) Bi |

- | | |
|----------------------------|----------------------------|
| (1) A-I, B-II, C-III, D-IV | (2) A-IV, B-III, C-II, D-I |
| (3) A-IV, B-II, C-III, D-I | (4) A-I, B-III, C-II, D-IV |

Answer (2)

Sol. Pnictogens → group-15 ⇒ N, P, As, Sb, Bi

Chalcogens → group-16 ⇒ O, S, Se, Te, Po

Halogens → group-17 ⇒ F, Cl, Br, I, At

Noble gases → group-18 ⇒ He, Ne, Ar, Kr, Xe, Rn

6. Find orbital angular momentum for 2s and 2p energy levels

- | | |
|------------------------------------|--------------------------------|
| (1) $0, \frac{h}{(\sqrt{2})\pi}$ | (2) $0, \frac{h}{\sqrt{2}\pi}$ |
| (3) $\frac{h}{\pi}, \frac{h}{\pi}$ | (4) $0, \frac{h}{2\pi}$ |

Answer (1)

Sol. Orbital angular momentum = $\frac{h}{2\pi} \sqrt{l(l+1)}$

For 2s $l = 0$

Orbital angular momentum = 0

For 2p $l = 1 \Rightarrow \frac{h}{2\pi} \sqrt{l(l+1)}$

7. Which of the following order is correct?

- (A) Electronegativity : B > Tl > In > Ga > Al
 (B) First Ionisation energy : B > Tl > Ga > Al > In
 (C) Density : Tl > In > Ga > Al > B
 (D) Size : B > Al > Ga > In > Al
- (1) (A, B, C) only
 (2) (B, C, D) only
 (3) (A, B, D) only
 (4) (A, B, C, D)

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
IN PHYSICS

1000+ 99
PERCENTILERS
IN CHEMISTRY

4000+ 95
PERCENTILERS
IN MATHS

100
Percentile
in
Physics
& Math



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in
Physics



Harsh Jha
PSID: 00014863322

99.99
Percentile
in
Physics
& Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



Answer (1)

Sol. EN : B > Tl > In > Ga > Al

(2.0) (1.8) (1.7) (1.6) (1.5)

IE₁ : B > Tl > Ga > Al > In


(801) (589) (579) (577) (558) (in kJ/mol)

Density : Tl > In > Ga > Al > B

(11.85) (7.31) (5.9) (2.7) (2.35) (in g/ml)

Size : B < Ga < Al < In < Tl

(85) (135) (143) (167) (170) (in pm)




this reaction is

- (1) Br₂/CCl₄ (2) Br₂/H₂O
(3) HBr/H₂SO₄ (4) Br₂/acetone

Answer (2)

Sol. Br₂ in water causes tribromination



H₂O Promotes formation of Br⁺ in excess

9. Which of the following vitamins are fat-soluble?

B₁₂, C, D, B, E

- (1) B and C
(2) D and E
(3) B and E
(4) C, D and E

Answer (2)

Sol. Fat soluble vitamins are (A, D, E and K)

10. Statement-I : CrO₃ is a strong oxidising agent


Statement-II : Cr⁺⁶ is more stable than Mo⁺⁶

considering the above statements, choose the correct option.

- (1) Both statement-I and Statement-II are correct
(2) Both statement-I and Statement-II are incorrect
(3) Statement-I is correct but Statement-II is incorrect
(4) Statement-I is incorrect but Statement-II is correct

Answer (3)

Sol. CrO₃ is a strong oxidising agent and itself is reduced to Cr³⁺ which is more stable than Cr⁶⁺ because Cr³⁺ has stable electronic configuration.



∴ Statement-I is correct.

Stability of +6 state in group-6 increases down the group. Therefore, Mo⁺⁶ is more stable than Cr⁺⁶. So, statement-II is incorrect.

11. Which of the following compound or complex ions is/are diamagnetic in nature

- (a) CrO₃ (b) [Fe(CN)₆]⁴⁻
(c) [Co(H₂O)₃F₃] (d) [Cr(NH₃)₆]³⁺

- (1) a and b only
(2) a, b and c only
(3) a, b, c and d
(4) c and d only

Answer (2)

THE LEGACY OF SUCCESS CONTINUES

OUR JEE CHAMPIONS

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS
(max. 1 each, 1 below)

1000+ 99 PERCENTILERS
5 ABOVE

4000+ 95 PERCENTILERS
5 ABOVE



100 Percentile in Physics & Math
100 Percentile in Chemistry
100 Percentile in Biology
100 Percentile
Shreyas Lohiya
PSID: 00003389699



100 Percentile in Physics & Chemistry
99.99 Percentile
Harsh Jha
PSID: 00014863322



100 Percentile in Physics & Chemistry
99.99 Percentile
Devyra Rustagi
PSID: 00014768785



100 Percentile in Physics & Chemistry
99.99 Percentile
Amogh Bansal
PSID: 00014769016



Chirag Falor
4 Year Classroom
1 AIR
JEE (Adv.)
2020





Tanishka Kabra
4 Year Classroom
1 AIR-16 CB
JEE (Adv.)
2022



Sanvi Jain
4 Year Classroom
1 AIR-34 CB
JEE (Adv.)
2024

Sol. Species having unpaired electron are paramagnetic in nature



Number of unpaired electron = 0

\Rightarrow diamagnetic



Number of unpaired electrons = 3

\Rightarrow Paramagnetic

12. 20 mL 1M NaOH is mixed with 10 mL 2M HCl which is further diluted to 100 mL. Find concentration of final solution?

(1) 2×10^{-3} M

(2) 0.2 M

(3) 2×10^{-2} M

(4) 0.1 M

Answer (2)



$$20 \times 1 \quad 10 \times 2$$

$$20 \text{ mmol} \quad 20 \text{ mmol}$$

$$0 \quad 0 \quad 20 \text{ mmol}$$

$$\text{So, concentration of NaCl} = \frac{20 \text{ mmol}}{100 \text{ mL}}$$

$$= 0.2 \text{ M}$$

13. Which of the following statement is correct w.r.t. Arrhenius equation?

(1) Dimensions of k and A are same

(2) k decreases with increase in temperature generally

(3) A decreases with increase in temperature always

(4) k increases as value of E_a increase

Answer (1)

$$\text{Sol. } k = A e^{-\frac{E_a}{RT}}$$


$$E_a \uparrow \quad k \downarrow$$

14. Match the column-I with column-II and choose the correct option


Column I

Column II


(P) Finkelstein reaction




(Q) Lucas Test



(3) Wurtz reaction



(4) Gatterman Koch



Reaction

(1) P \rightarrow (4), Q \rightarrow (3), R \rightarrow (2), S \rightarrow (1)

(2) P \rightarrow (1), Q \rightarrow (2), R \rightarrow (3), S \rightarrow (4)

(3) P \rightarrow (1), Q \rightarrow (3), R \rightarrow (2), S \rightarrow (4)

(4) P \rightarrow (3), Q \rightarrow (2), R \rightarrow (1), S \rightarrow (4)

Answer (1)

Sol. P \rightarrow (4), Q \rightarrow (3), R \rightarrow (2), S \rightarrow (1)

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
IN PHYSICS

1000+ 99
PERCENTILERS
IN CHEMISTRY

4000+ 95
PERCENTILERS
IN BIOLOGY

100
Percentile
in
Physics
& Chemistry



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in
Physics



Harsh Jha
PSID: 00014863322

99.99
Percentile
in
Physics
& Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



15. **Statement-I** : When ice and water are at equilibrium, heat is absorbed by system, yet there is no increase in temperature until ice melts completely.

Statement-II : System absorbs the heat to break inter molecular H-bond and there is no increase in kinetic energy.

- (1) Both statement-I and Statement-II are correct
- (2) Statement-I is correct. Statement-II is incorrect
- (3) Statement-I is incorrect. Statement-II is correct
- (4) Both statements-I and II are incorrect

Answer (1)

Sol. Heat absorbed is consumed in conversion of solid ice to liquid water. So no temperature change. Even kinetic energy is same is the transition.

16. **Statement-I** : Wet cotton clothes made up of cellulose based carbohydrates take a comparatively longer time to get dried than wet nylon based clothes.

Statement-II : Both form intermolecular H-bonds with water molecules

- (1) Both statement-I and statement-II are correct
- (2) Both statement-I and statement-II are incorrect
- (3) Statement-I is correct but statement-II is incorrect
- (4) Statement-I is incorrect but statement-II is correct

Answer (1)


Sol. Wet cellulose-based cotton clothes take longer time to dry than wet nylon-based clothes due to more number of H-bonds between cellulose and water molecules. So statement-I is correct. Statement-II is also correct as both the cellulose and nylon form intermolecular H-bonds with water molecules.

- 17.
- 18.
- 19.
- 20.

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

21. Find out magnitude of work done in the process ABCD (in kJ) (1 Latm = 101.3 J)



Answer (304)

Sol. $(W) = 3 \times 1000$
 $= 3000 \text{ atm L}$
 $(W) = 303900 \text{ Joule}$
 $= 303.9 \text{ kJ}$
 $= 304 \text{ kJ}$

22. Amount of magnesium (Mg) (in mg) required to liberate 224 mL of H₂ gas at STP, when reacted with HCl.

Answer (240)

Sol. $\text{Mg} + 2\text{HCl} \rightarrow \text{MgCl}_2 + \text{H}_2$

Moles of H₂ gas at STP = $\frac{224}{22400} \text{ mol}$
 $= 0.01 \text{ mol}$

1 mol H₂ formed by 1 mol of Mg.
 mol of Mg required = 0.01 mol
 mass of Mg required = 0.01×24
 $= 0.24 \text{ g}$
 $= 240 \text{ mg}$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE TOPPERS

70+ 100 PERCENTILERS

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100 Percentile in Physics & Maths



Shreyas Lohiya
PSID: 00003389699

100 Percentile in Physics



Harsh Jha
PSID: 00014863322

100 Percentile in Physics & Chemistry



Devya Rustagi
PSID: 00014768785

99.99 Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS

Chirag Falor
4 Year Classroom
1 AIR-36 (Male) JEE (Main) 2020

Tanishka Kabra
4 Year Classroom
1 AIR-16 CBZ JEE (Main) 2022

Sanvi Jain
4 Year Classroom
1 AIR-34 CBZ JEE (Main) 2024

23. Among Sc, Ti, Mn and Co, calculate the spin only magnetic moment in +2 oxidation state of metal having highest heat of atomisation

Answer (3)

Sol. Enthalpy of atomisation


$$\text{Sc} - 326 \text{ kJ mol}^{-1}$$

$$\text{Ti} - 473 \text{ kJ mol}^{-1}$$

$$\text{Mn} - 281 \text{ kJ mol}^{-1}$$

$$\text{Co} - 425 \text{ kJ mol}^{-1}$$

Ti has highest heat of atomisation



$$\mu_{\text{spin only}} = \sqrt{n(n+2)} \text{ BM}$$

n = number of unpaired electrons

$$= 2$$

$$\mu = \sqrt{2(2+2)}$$

$$= \sqrt{8} \text{ BM}$$

$$= 2.82 \text{ BM}$$


$$\approx 3 \text{ BM}$$

24. C_9H_{12} is a derivative of benzene, how many total structural isomers of molecular formula C_9H_{12} are possible.

Answer (8)


Sol. Degree of unsaturation =

$$\frac{2 \times 9 + 2 - 12}{2} = 4$$



25. 'x' g of nitrobenzene gives 4.2 g 1, 3-dinitrobenzene with 100% yield. Find the value of 'x'.

Answer (3)



$$\text{Molar mass of } \text{C}_6\text{H}_5\text{NO}_2 = 123 \text{ g mol}^{-1}$$

$$\text{Molar mass of } \text{C}_6\text{H}_4(\text{NO}_2)_2 = 168 \text{ g mol}^{-1}$$

$$\text{No. of moles of } \text{C}_6\text{H}_5\text{NO}_2 = \text{No of moles } \text{C}_6\text{H}_4(\text{NO}_2)_2$$

$$\frac{x}{123} = \frac{4.2}{168}$$

$$x = 3.075 \text{ g} \approx 3 \text{ g}$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS
(max. 1 each, 1 below)

1000+ 99
PERCENTILERS
& ABOVE

4000+ 95
PERCENTILERS
& ABOVE

100
Percentile
in
Physics
&
Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in
Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in
Physics
&
Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016



Chirag Falor
4 Year Classroom
AIR-36 (Male)
JEE (Main)-3
2020



Tanishka Kabra
4 Year Classroom
AIR-16 CBL
JEE (Main)-3
2022



Sanvi Jain
4 Year Classroom
AIR-34 CBL
JEE (Main)-3
2024