

CHEMISTRY

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

1. Which of the following is animal starch?

- (1) Glycogen
- (2) Lactose
- (3) Amylopectin
- (4) Amylose

Answer (1)

Sol. Lactose is present in milk.

Amylopectin and amylose are part of starch.

Glycogen is animal starch.

2. **Statement 1** : Correct order of ionic radius for Mg^{2+} ,

Na^+ , O^{2-} , & F^- is $\text{F}^- > \text{O}^{2-} > \text{Na}^+ > \text{Mg}^{2+}$

Statement 2 : Correct order of electron gain enthalpy

for 17th group elements follows order $\text{Cl} > \text{F} > \text{Br} > \text{I}$

(Magnitude only)



- (1) Statement-1 & Statement-2 are correct
- (2) Statement-1 is correct Statement-2 is incorrect
- (3) Statement-1 & Statement-2 are incorrect
- (4) Statement-1 is incorrect Statement-2 is correct

Answer (4)

Sol.: Correct order of ionic radius $\text{O}^{2-} > \text{F}^- > \text{Na}^+ > \text{Mg}^{2+}$


Correct order for electron gain enthalpy (Magnitude)
 $\text{Cl} > \text{F} > \text{Br} > \text{I}$

3. Identify the product formed in the following reaction



Answer (2)

Sol. Aryl halides having strong electron withdrawing group like NO_2 either at the ortho or para position undergo $\text{S}_{\text{N}}\text{Ar}$ reaction easily involving carbanion intermediate



4. Which of the following is steam volatile

- (1) Ortho nitrophenol
- (2) Para nitrophenol
- (3) Para aminophenol
- (4) Para nitroaniline

Answer (1)

Sol. Ortho nitrophenol is steam volatile due to intramolecular H-bonding. It's B.P is less. p-nitrophenol, p-amino phenol, paranitro aniline show intermolecular H-bonding

5. Consider the following complexes

- $[\text{Mn}(\text{CN})_6]^{4-}$ (1) $[\text{Fe}(\text{CN})_6]^{4-}$ (2) $[\text{Fe}(\text{CN})_6]^{3-}$ (3) $[\text{Co}(\text{CN})_6]^{3-}$ (4)

Correct order of CFSE (Δ) will be

- (1) $3 > 4 > 2 > 1$
- (2) $4 > 3 > 2 > 1$
- (3) $4 > 3 > 1 > 2$
- (4) $3 > 4 > 1 > 2$

Answer (2)

Sol. (1) $[\text{Mn}(\text{CN})_6]^{4-}$, Mn^{2+}

(2) $[\text{Fe}(\text{CN})_6]^{4-}$, Fe^{2+}

(3) $[\text{Fe}(\text{CN})_6]^{3+}$, Fe^{3+}

(4) $[\text{Co}(\text{CN})_6]^{3+}$, Co^{3+}

order of CFSE will be $4 > 3 > 2 > 1$



Delivering Champions Consistently



Medical | IIT-JEE | Foundations

JEE (Advanced) 2024



AIR 25

Rishi Shekher Shukla
2 Year Classroom

JEE (Main) 2024



AIR 67

Krishna Sai Shishir
2 Year Classroom

JEE (Advanced) 2024



AIR 78

Abhishek Jain
2 Year Classroom

JEE (Advanced) 2024



AIR 93

Harsh Aggarwal
2 Year Classroom

JEE (Advanced) 2024



AIR 95

Ujjwal Singh
2 Year Classroom

JEE (Advanced) 2024



AIR 98

Rashit Aggarwal
2 Year Classroom

JEE (Main) 2024



AIR 1

Sanvi Jain
2 Year Classroom


JEE (Main) 2024



AIR 15

M Sai Divya Teja Reddy
2 Year Classroom


JEE (Main) 2024




AIR 19

Rishi Shekher Shukla
2 Year Classroom

6. Consider the following reaction



Identify the final product P.




Answer (1)

Sol. Clemmensen's reduction reagent reduces aldehyde and ketone to alkane.

7. What is the value of van't Hoff Factor for A_2B , if 30% of A_2B is dissociated?

- (1) 1.60 (2) 1.30
(3) 1.50 (4) 1.20

Answer (1)




$$i = 1 - \alpha + 2\alpha + \alpha = 1 + 2\alpha$$




$$\alpha = 0.30$$

$$i = 1 + 2 \times 0.30 = 1.60$$

8. Find the order of the reaction



if the mechanism of the reaction is as follows:



- (1) 1 (2) 3
(3) 2 (4) 4

Answer (3)

Sol. Since the slowest step is considered as rate determining step.

$$\text{So, here } r = k[A][B]$$

$$\text{Order} = 2$$

9. Match the following **List-I** with **List-II** and choose the correct option

List-I (Complexes)

List-II (Hybridisation)


- (A) $[\text{Co}(\text{OX})_3]^{3-}$ (i) sp^3d^2
(B) $[\text{FeF}_6]^{3-}$ (ii) d^2sp^3
(C) $[\text{Ni}(\text{CO})_4]$ (iii) dsp^2
(D) $[\text{PtCl}_4]^{2-}$ (iv) sp^3

- (1) A-(i), B-(ii), C-(iii), D-(iv)
(2) A-(ii), B-(i), C-(iii), D-(iv)
(3) A-(i), B-(ii), C-(iv), D-(iii)
(4) A-(ii), B-(i), C-(iv), D-(iii)


Answer (4)

Sol. : $[\text{Co}(\text{OX})_3]^{3-} \Rightarrow \text{Co}^{3+}$, (OX) act as SFL for Co^{3+}


$$\Rightarrow d^6 \Rightarrow t_{2g}^6 e_g^0 \Rightarrow d^2sp^3 \text{ hybridisation}$$



$$\text{Fe}^{3+} \Rightarrow d^5 \Rightarrow t_{2g}^3 e_g^2 \Rightarrow sp^3d^2 \text{ hybridisation.}$$





$$\text{Ni}(0) \Rightarrow s^2d^8 \Rightarrow d^{10} \Rightarrow sp^3 \text{ hybridisation}$$



$$\text{Pt}^{2+} \Rightarrow d^8 \Rightarrow dsp^2 \text{ hybridisation.}$$

10. What is the correct Nernst equation representation for the following cell reaction



$$(1) E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{RT}{2F} \ln \frac{[\text{Mg}^{2+}]}{[\text{Ag}^+]^2}$$

$$(2) E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{RT}{2F} \ln \frac{[\text{Ag}^+]^2}{[\text{Mg}^{2+}]}$$




Delivering Champions Consistently

JEE (Advanced) 2024

JEE (Main) 2024

$$(3) E_{\text{cell}} = E_{\text{cell}}^{\circ} + \frac{RT}{F} \ln \frac{[\text{Mg}^{2+}]}{[\text{Ag}^+]^2}$$

$$(4) E_{\text{cell}} = E_{\text{cell}}^{\circ} + \frac{RT}{2F} \ln \frac{[\text{Ag}^+]^2}{[\text{Mg}^{2+}]}$$

Answer (1)

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{RT}{2F} \ln \frac{[\text{Mg}^{2+}]}{[\text{Ag}^+]^2}$$


11. The correct order of melting point of d-block elements is :

- (1) Fe > Mn (2) Tc > Ru
 (3) Os > Re (4) Ta > W

Answer (1)


Sol. Melting point order is Fe > Mn, Ru > Tc, Re > Os, W > Ta

12. Consider the following reaction



If P is total pressure at equilibrium & K_p is equilibrium constant. Then α in terms of K_p & P is (Assume $\alpha \ll 1$)

- (1) $\sqrt{\frac{K_p}{P}}$ (2) $\sqrt[4]{\frac{K_p}{P}}$
 (3) $\sqrt{\frac{2K_p}{P}}$ (4) $\sqrt[3]{\frac{2K_p}{P}}$

Answer (4)

$$t = 0 \quad p_0$$

$$t = t_{\text{eq}} \quad p_0(1 - \alpha) \quad p_0\alpha \quad p_0\frac{\alpha}{2}$$

$$P = p_0 + p_0\frac{\alpha}{2}$$

$$P = p_0\left(1 + \frac{\alpha}{2}\right) \quad (P \approx p_0)$$

$$\text{At equilibrium } K_p = \frac{(p_{\text{A}_2})\left(\frac{p_{\text{B}_2}}{2}\right)}{(p_{\text{A}_2\text{B}})} = (\alpha \ll 1)$$

$$K_p = \frac{(p_0\alpha)\left(p_0\frac{\alpha}{2}\right)^{\frac{1}{2}}}{p_0(1 - \alpha)} = K_p = \alpha\left(p\frac{\alpha}{2}\right)^{\frac{1}{2}}$$

$$\frac{K_p}{\frac{1}{p^2}} = \frac{\alpha^{3/2}}{2^{1/2}}$$

$$\frac{2K_p^2}{P} = \alpha^3$$


$$\sqrt[3]{\frac{2K_p^2}{P}} = \alpha$$

13. $\wedge_m$ is linearly dependent to $\sqrt{c}$ for an electrolyte, then molar conductance for the same electrolyte at infinite dilution shows

- (1) Small increase (2) Small decrease
 (3) Sharp increase (4) Sharp decrease

Answer (1)

Sol. $\wedge_m$ decreases linearly with $\sqrt{c}$ for strong electrolytes having small -ve slope. It can be extrapolated to $\wedge_m^{\infty}$ as $c \rightarrow 0$.



The molar conductance of the same electrolyte at infinite dilution or as $c \rightarrow 0$ shows small increase.

14. Given ionisation enthalpy of element E(g) is 300 kJ/mol and electron gain enthalpy of A, B, C and D gaseous atoms are -320 kJ/mol, -340 kJ/mol, -200 kJ/mol and -250 kJ/mol, then what will be the correct order of ionic nature of compounds?

- (1) EB > EA > ED > EC (2) EB > EA > EC > ED
 (3) EC > ED > EA > EB (4) EC > ED > EB > EA

Answer (1)



Delivering Champions Consistently



Medical | IIT-JEE | Foundations

JEE (Advanced) 2024



AIR 25

Rishi Shekhar Shukla
2 Year Classroom

JEE (Main) 2024



AIR 67

Krishna Sai Shishir
2 Year Classroom

JEE (Advanced) 2024



AIR 78

Abhishek Jain
2 Year Classroom

JEE (Advanced) 2024



AIR 93

Harsh Aggarwal
2 Year Classroom

JEE (Advanced) 2024



AIR 95

Ujjwal Singh
2 Year Classroom

JEE (Advanced) 2024



AIR 98

Rashit Aggarwal
2 Year Classroom

JEE (Main) 2024



AIR 1

Sanvi Jain
2 Year Classroom

JEE (Main) 2024



AIR 15

M Sai Divya Teja Reddy
2 Year Classroom

JEE (Main) 2024




AIR 19

Rishi Shekhar Shukla
2 Year Classroom

Sol. Since ionic strength depends on IE of electropositive atom; E.G.E. of electronegative element and lattice energy, more the negative value of electron gain enthalpy, more will be ionic nature.

15. Graph between de Broglie wavelength (λ_D) and kinetic energy (K) of an electron is



Answer (2)

Sol. de Broglie wavelength (λ_D) of an electron of mass (m), moving with velocity (v) is given by

$$\lambda_D = \frac{h}{mv}$$

Where h is planck's constant.


$$\text{Kinetic energy (K)} = \frac{1}{2}mv^2$$

$$mv = \sqrt{2mK}$$

$$\lambda_D = \frac{h}{\sqrt{2mK}}$$

$$\frac{1}{K} = \frac{2m\lambda_D^2}{h^2}$$

Plot of $\frac{1}{K}$ vs λ_D is



16. Which of the following ions is strongest oxidising agent

$$\text{Given : } E_{\text{Al}^{3+}/\text{Al}}^\circ = -2.7\text{V}$$

$$E_{\text{Cu}^{2+}/\text{Cu}}^\circ = 0.34\text{V}$$

$$E_{\text{Pb}^{4+}/\text{Pb}^{2+}}^\circ = 1.8\text{V}$$


$$E_{\text{Ti}^{3+}/\text{Ti}^{2+}}^\circ = -0.37\text{V}$$

- (1) Al^{3+} (2) Cu^{2+}
(3) Pb^{4+} (4) Ti^{3+}

Answer (3)

Sol. Reduction potential of $\text{Pb}^{4+} \rightarrow \text{Pb}^{2+}$ is most positive, Hence Pb^{4+} is strongest oxidising agent.

17. Total number of nucleophiles among the following are Ph-SH , OH^- , $\text{CH}_2=\text{CH}_2$, >N-CH_3 , H_3O^+ ,



- (1) 5 (2) 6
(3) 7 (4) 4

Answer (2)

Sol. Species having atom containing lone pair available for donation can act as nucleophile

18. Radius of 1st orbit of hydrogen atom is $a_0 \text{ \AA}$, then find de-Broglie wavelength of 2nd orbit of hydrogen atom.

- (1) $4\pi a_0$ (2) $\frac{4}{\pi a_0}$
(3) $8\pi a_0$ (4) $2\pi a_0$

Answer (1)

Sol. $r_n = a_0 \frac{n^2}{Z}$
for $n = 1, Z = 1$

$$r_1 = a_0$$

$$r_2 = a_0 \frac{4}{1} = 4a_0$$

$$2\pi r_n = n\lambda$$

$$\lambda = \frac{2\pi r_2}{2} = \frac{2\pi \times 4a_0}{2} = 4\pi a_0$$

19.

20.

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.



Delivering Champions Consistently


JEE (Advanced) 2024

AIR 25 Rishi Shekher Shukla 2 Year Classroom	AIR 67 Krishna Sai Shishir 2 Year Classroom	AIR 78 Abhishek Jain 2 Year Classroom	AIR 93 Harsh Aggarwal 2 Year Classroom	AIR 95 Ujjwal Singh 2 Year Classroom	AIR 98 Rishabh Aggarwal 2 Year Classroom
---	--	--	---	---	---


JEE (Main) 2024

Karnataka Topper AIR 1 Sanvi Jain 2 Year Classroom	Telangana Topper AIR 15 M Sai Divya Teja Reddy 2 Year Classroom	Telangana Topper AIR 19 Rishi Shekher Shukla 2 Year Classroom
---	--	--

21. Calculate the total number of sigma and π -bonds in the given molecule?



Answer (15)



Sol.

Number of sigma bonds = 11 σ

Number of π -bonds = 4 π


Total = 15

22. Chromite ore + $\text{Na}_2\text{CO}_3 + \text{O}_2 \longrightarrow$ Insoluble product

Calculate the molar mass of insoluble product formed.

(Given : Molar mass of Cr = 52 g/mol, Na = 23 g/mol, Fe = 56 g/mol, O = 16 g/mol)

Answer (160)




Molar mass of Fe_2O_3

$$\Rightarrow 2(56) + 3(16)$$


$$\Rightarrow 160$$


23. Consider the following amines



1 gram of most basic compound reacts with x mg of HCl, calculate value of x.

Answer (341)

Sol. Most basic compound is 




$$\frac{1}{107} \text{ mol} \quad \frac{1}{107} \text{ mol}$$

mass of HCl required to react with Benzyl amine

$$= \frac{1}{107} \times 36.5 \text{ g}$$


$$= 0.341 \text{ g} = 341 \text{ mg}$$

24. Consider the following reaction



Find the mass of final product(D) formed in g

Answer (13)



Molar mass of D = 130 g mol^{-1}

Mass of 0.1 mol of (D) formed = 13g

25.



Delivering Champions Consistently



Medical | IIT-JEE | Foundations

JEE (Advanced) 2024



AIR 25
Rishi Shekher Shukla
2 Year Classroom



AIR 67
Krishna Sai Shishir
2 Year Classroom



AIR 78
Abhishek Jain
2 Year Classroom



AIR 93
Hardik Aggarwal
2 Year Classroom



AIR 95
Ujjwal Singh
2 Year Classroom



AIR 98
Rashit Aggarwal
2 Year Classroom

JEE (Main) 2024



AIR 1
Sanvi Jain
2 Year Classroom

Telangana Topper



AIR 15
M Sai Divya Teja Reddy
2 Year Classroom

Telangana Topper



AIR 19
Rishi Shekher Shukla
2 Year Classroom