

Answer & Solutions *for* JEE MAIN 2026

Don't guess your JEE Main Score.

Calculate it Instantly

with our

**JEE Main Score
Calculator.**

HOW TO
GET YOUR
SCORES

01

Open your response
sheet on the JEE Main Site

02

Copy the link from the
browser's Address bar

03

Paste it into the required
field in the Calculator

 jee-marks-calculator.aakash.ac.in/



Scan the QR code
to know more.

ADMISSIONS OPEN FOR SESSION 2026-27

Get up to

90% Total Scholarship*

Appear for Instant Admission cum Scholarship Test
Register now for **FREE**, visit: iacst.aakash.ac.in

SCAN TO REGISTER



*Terms & Conditions Apply

TRUST Aakash

Stellar Ranks in JEE (Main) 2025

2 Ranks in
Top 10 AIR*

12 Ranks in
Top 100 AIR*

39 Ranks in
Top 500 AIR*

*Includes students of classroom, digital & distance across all categories.



Harsh
Jha
AIR
23



Kushagra
Baingaha
AIR
7



Shreyas
Lohiya
AIR
6



Harssh
A Gupta
AIR
15



Devya
Rustagi
AIR
28



Shreyas
Lohiya
AIR
68



Rujul
Garg
AIR
41



Advay
Mayank
AIR
36



Arush
Anand
AIR
64



Kotha
D Reddy
AIR
74

Top Ranks in JEE (Advanced) 2025

13 Ranks in
Top 100 AIR*

51 Ranks in
Top 500 AIR*

103 Ranks in
Top 1000 AIR*

*Includes students of classroom, digital & distance across all categories.



Scan the QR Code
to access JEE
Cut-off 2025

Aakashians Create History in International Olympiads

Our Olympiads Results

378 Classroom
Students
Aakashians Qualified
in NSEs 2025-26

777 Classroom
Students
Aakashians Qualified
in IOQM 2025

134 Classroom
Students
Aakashians Qualified
in RMO 2025



Aarav Gupta
Gold Medalist

66th International
Mathematical Olympiad
(IMO) 2025



Yug Gandhi
Gold Medalist

Singapore Math
Olympiad 2025



Arjun Tyagi
Gold Medalist

International Olympiad
in Artificial Intelligence
(IOAI) 2025

HOME OF PROBLEM SOLVERS

MATHEMATICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer :

1. If the line $\alpha x + 4y = \sqrt{7}$, where $\alpha \in \mathbb{R}$, touches the ellipse $3x^2 + 4y^2 = 1$ at the point P in the first quadrant, then one of the focal distances of P is:

- (1) $\frac{1}{\sqrt{3}} - \frac{1}{2\sqrt{5}}$
- (2) $\frac{1}{\sqrt{3}} + \frac{1}{2\sqrt{5}}$
- (3) $\frac{1}{\sqrt{3}} + \frac{1}{2\sqrt{7}}$
- (4) $\frac{1}{\sqrt{3}} - \frac{1}{2\sqrt{11}}$

Answer (3)

Sol. There is exactly one intersection point of the line

$$\alpha x + 4y = \sqrt{7} \text{ and the ellipse } 3x^2 + 4y^2 = 1$$

$$\Rightarrow 3x^2 + 4\left(\frac{\sqrt{7} - \alpha x}{4}\right)^2 = 1$$

$$12x^2 + 7 + \alpha^2 x^2 - 2\sqrt{7}\alpha x - 4 = 0$$

$$(12 + \alpha^2)x^2 - 2\sqrt{7}\alpha x + 3 = 0$$

$$\Rightarrow D = 0$$

$$28\alpha^2 = 12(12 + \alpha^2) \Rightarrow \alpha = \pm 3$$

\therefore Line touches Ellipse in first quadrant

\Rightarrow Slope of line is negative.

$$\Rightarrow \alpha = 3$$

$$\Rightarrow x^2 - \frac{2}{\sqrt{7}}x + \frac{1}{7} = 0 \Rightarrow x = \frac{1}{\sqrt{7}}$$

$$\Rightarrow y = \frac{\sqrt{7} - \alpha x}{4} = \frac{\sqrt{7} - 3}{4\sqrt{7}} = \frac{1}{\sqrt{7}}$$

$$\text{Point } P \equiv \left(\frac{1}{\sqrt{7}}, \frac{1}{\sqrt{7}}\right)$$

$$E: \frac{x^2}{\frac{1}{3}} + \frac{y^2}{\frac{1}{4}} = 1 \Rightarrow \frac{1}{4} = \frac{1}{3}(1 - e^2) \Rightarrow e = \frac{1}{2}$$

$$\text{Equation of directrices are } x = \pm \frac{2}{\sqrt{3}}$$

$$\Rightarrow SP = ePM = \frac{1}{2}\left(\frac{2}{\sqrt{3}} - \frac{1}{\sqrt{7}}\right) = \frac{1}{\sqrt{3}} - \frac{1}{2\sqrt{7}}$$

$$S'P = ePM' = \frac{1}{2}\left(\frac{2}{\sqrt{3}} + \frac{1}{\sqrt{7}}\right) = \frac{1}{\sqrt{3}} + \frac{1}{2\sqrt{7}}$$

\Rightarrow Option (3) is Correct.

2. Let $A = \{x : |x^2 - 10| \leq 6\}$ and $B = \{x : |x - 2| > 1\}$.

Then

- (1) $A - B = [2, 3]$
- (2) $A \cup B = (-\infty, 1] \cup (2, \infty)$
- (3) $B - A = (-\infty, -4) \cup (-2, 1) \cup (4, \infty)$
- (4) $A \cap B = [-4, -2] \cup [3, 4]$

Answer (3)

Sol. $|x^2 - 10| \leq 6$

$$\Rightarrow -6 \leq x^2 - 10 \leq 6$$

$$\Rightarrow 4 \leq x^2 \leq 16$$

$$\Rightarrow 2 \leq |x| \leq 4$$

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



(5, 2), (7, 2), (9, 2), (5, 3), (7, 3), (9, 3), (9, 5)

$$\Rightarrow m = 7$$

$$\Rightarrow l + m = 25$$

Option (3) is correct

6. A random variable X takes values 0,1,2,3 with probabilities $\frac{2a+1}{30}, \frac{8a-1}{30}, \frac{4a+1}{30}, b$ respectively, where $a, b \in \mathbb{R}$. Let μ and σ respectively be the mean and standard deviation of X such that

$$\sigma^2 + \mu^2 = 2. \text{ Then } \frac{a}{b} \text{ is equal to :}$$

- (1) 60 (2) 3
(3) 30 (4) 12

Answer (1)

Sol. $\sigma^2 + \mu^2 = 2 = 0^2 \times \left(\frac{2a+1}{30}\right)$

$$+ 1^2 \left(\frac{8a-1}{30}\right) + 2^2 \left(\frac{4a+1}{30}\right) + 3b$$

After solving:

$$24a + 270b = 57 \quad \dots(i)$$

$$\sum xp(x) = 1$$

$$\Rightarrow \frac{2a+1}{30} + \frac{8a-1}{30} + \frac{4a+1}{30} + b = 1$$

$$\Rightarrow b = \frac{2a-14a}{30} \quad \dots(ii)$$

solving (i) and (ii)

$$a = 2, b = \frac{1}{30}$$


$$\frac{a}{b} = 60$$

7. Let z be the complex number satisfying $|z-5| \leq 3$ and having maximum positive principal argument.

Then $34 \left| \frac{5z-12}{5z+16} \right|^2$ is equal to :

- (1) 20
(2) 12
(3) 16
(4) 26

Answer (1)



Sol.

$$\Rightarrow \arg(z) = \sin^{-1}\left(\frac{3}{5}\right) = \tan^{-1}\left(\frac{3}{4}\right).$$

$$\Rightarrow z = |z|e^{i\theta} = 4 \times (\cos\theta + i\sin\theta)$$

$$= 4 \times \left(\frac{4}{5} + i\frac{3}{5}\right)$$

$$= \frac{16}{5} + \frac{12i}{5}$$

$$\Rightarrow 5z = 16 + 12i$$

$$5zi = 16i - 12$$

$$\left(\frac{5z-12}{5zi+16}\right) = \frac{(4+12i)}{(4+16i)} = \frac{(1+3i)}{(1+4i)}$$

$$\Rightarrow \left| \frac{5z-12}{5zi+16} \right| = \frac{\sqrt{10}}{\sqrt{17}}$$

$$34 \left| \frac{5z-12}{5zi+16} \right|^2 = 34 \times \frac{10}{17} = 20.$$

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



10. For the matrices $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$ and $B = \begin{bmatrix} -29 & 49 \\ -13 & 18 \end{bmatrix}$,

if $(A^{15} + B) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$, then among the following

which one is true?

- (1) $x = 18, y = 11$ (2) $x = 11, y = 2$
 (3) $x = 16, y = 3$ (4) $x = 5, y = 7$

Answer (2)

Sol. $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$

$$A^2 = \begin{bmatrix} 5 & -8 \\ 2 & -3 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 7 & -12 \\ 3 & -5 \end{bmatrix}$$

$$A^n = \begin{bmatrix} 2n+1 & -4n \\ n & -(2n-1) \end{bmatrix}$$

$$\Rightarrow A^{15} = \begin{bmatrix} 31 & -60 \\ 15 & -29 \end{bmatrix}$$

$$A^{15} + B = \begin{bmatrix} 2 & -11 \\ 2 & -11 \end{bmatrix}$$

$$(A^{15} + B) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -11 \\ 2 & -11 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$2x - 11y = 0$$

$$\therefore x = 11, y = 2$$

11. Let $f(x) = x^3 + x^2 f'(1) + 2x f''(2) + f'''(3)$, $x \in \mathbf{R}$.

Then the value of $f'(5)$ is

- (1) $\frac{62}{5}$ (2) $\frac{657}{5}$
 (3) $\frac{2}{5}$ (4) $\frac{117}{5}$

Answer (4)

Sol. Let $f(1) = a$

$$f'(2) = b$$

$$f''(3) = c$$

$$f(x) = x^3 + ax^2 + bx + c$$

$$f'(x) = 3x^2 + 2ax + b$$

$$f'(1) = a = 3 + 2a + b \Rightarrow a + b = 3 \dots (1)$$

$$f''(x) = 6x + 2a$$

$$\Rightarrow f''(2) = 12 + 2a = \frac{b}{2} \Rightarrow 4a - b = -24$$

$$\Rightarrow f''(x) = 6$$

$$\Rightarrow f''(3) = c = 6$$

$$\Rightarrow a = \frac{-27}{5}, b = \frac{12}{5}$$

$$f'(5) = 75 + 10a + b$$

$$= 75 - 54 + \frac{12}{5}$$

$$= 21 + \frac{12}{5} = \frac{117}{5}$$

12. Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a twice differentiable function such that $f''(x) > 0$ for all $x \in \mathbf{R}$ and $f'(a-1) = 0$ where a is a real number. Let $g(x) = f(\tan^2 x - 2\tan x + a)$,

$0 < x < \frac{\pi}{2}$. Consider the following statements:

(I) g is increasing in $\left(0, \frac{\pi}{4}\right)$

(II) g is decreasing in $\left(\frac{\pi}{2}, \frac{\pi}{4}\right)$

Then,

- (1) Only (I) is true
 (2) Both (I) and (II) are true
 (3) Only (II) is true
 (4) Neither (I) nor (II) is true

Answer (4)

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



Sol. $g(x) = f(\tan^2 x - 2\tan x + a)$

$$g'(x) = f'(\tan^2 x - 2\tan x + a)(2\tan x - 2)\sec^2 x$$

$$= f'((\tan x - 1)^2 + a - 1)(2\tan x - 2)\sec^2 x$$

$f(x)$ is increasing

$$f(a - 1) = 0$$

$$f(x) < 0 \text{ for } x < a - 1$$

$$f(x) > 0 \text{ for } x > a - 1$$

$$(\tan x - 1)^2 + (a - 1) > a - 1$$

$$\Rightarrow f'((\tan x - 1)^2 + (a - 1)) > f'(a - 1)$$

$$\Rightarrow f'((\tan x - 1)^2 + (a - 1)) > 0$$

\therefore For $x \in \left(0, \frac{\pi}{4}\right)$, $g'(x) < 0 \Rightarrow g(x)$ is decreasing

For $x \in \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$, $g'(x) > 0 \Rightarrow g(x)$ is increasing

13. Let $y = y(x)$ be the solution of the differential equation $\sec x \frac{dy}{dx} - 2y = 2 + 3\sin x$, $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$,

$y(0) = -\frac{7}{4}$. Then $y\left(\frac{\pi}{6}\right)$ is equal to

(1) $-\frac{5}{4}$

(2) $-\frac{5}{2}$

(3) $-3\sqrt{2} - 7$

(4) $-3\sqrt{3} - 7$

Answer (2)

Sol. $(\sec x) \frac{dy}{dx} - 2y = 2 + 3\sin x$

$$\Rightarrow \frac{dy}{dx} - 2\cos xy = 2\cos x + 3\sin x \cdot \cos x$$

IF = $e^{-2\sin x}$

$$y \cdot e^{-2\sin x} = \int e^{-2\sin x} \cdot \cos x(2 + 3\sin x) dx$$

$$\sin x = t \Rightarrow \cos x dx = dt$$

$$y \cdot e^{-2\sin x} = \int e^{-2t}(2 + 3t) dt$$

$$= 2 \frac{e^{-2t}}{-2} + 3 \int t \cdot e^{-2t} dt$$

$$= -e^{-2t} + 3 \left(\frac{t \cdot e^{-2t}}{-2} + \frac{1}{2} \int e^{-2t} dt \right)$$

$$-e^{-2t} + 3 \left(-\frac{te^{-2t}}{2} - \frac{1}{4} e^{-2t} \right) + c$$

$$y = -1 - \frac{3}{2}(\sin x) - \frac{3}{4} + ce^{2\sin x}$$

$$= -\frac{7}{4} - \frac{3}{2}\sin x + c \cdot e^{2\sin x}$$

$$y(0) = -\frac{7}{4} \Rightarrow c = 0$$

$$y = -\frac{7}{4} - \frac{3}{2}\sin x$$

$$y(\pi/6) = -\frac{7}{4} - \frac{3}{4} = -\frac{10}{4} = -\frac{5}{2}$$

14. Let $a_1, \frac{a_2}{2}, \frac{a_3}{2^2}, \dots, \frac{a_{10}}{2^9}$ be a G.P. of common ratio

$\frac{1}{\sqrt{2}}$. If $a_1 + a_2 + \dots + a_{10} = 62$, then a_1 is equal to

(1) $\sqrt{2} - 1$

(2) $2(2 - \sqrt{2})$

(3) $2(\sqrt{2} - 1)$

(4) $2 - \sqrt{2}$

Answer (3)

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



Sol. $\frac{a_2}{2} = a_1 r \Rightarrow a_2 = 2a_1 r = \sqrt{2}a_1$

$\frac{a_3}{2^2} = a_1 r^2 \Rightarrow a_3 = 4a_1 r^2 = 2a_1$

$\frac{a_4}{2^3} = a_1 r^3 \Rightarrow a_4 = 8a_1 r^3 = 2\sqrt{2}a_1$

$a_1 + a_2 + \dots + a_{10} = 62$

$= a_1 + \sqrt{2}a_1 + 2a_1 + \dots = 62$

$= \frac{a_1((\sqrt{2})^{10} - 1)}{\sqrt{2} - 1} = 62$

$= a_1 = 2(\sqrt{2} - 1)$

15. The positive integer n , for which the solutions of the equation $x(x+2) + (x+2)(x+4) + \dots + (x+2n-2)(x+2n) = \frac{8n}{3}$ are two consecutive even integers,

is:

(1) 3 (2) 12

(3) 6 (4) 9

Answer (1)

Sol. $x(x+2) + (x+2)(x+4) + \dots + (x+2n-2)(x+2n) = \frac{8n}{3}$

$T_r = (x+2r-2)(x+2r), r = 1, 2, \dots, n$

$\sum T_r = \sum x^2 + 2x \sum (2r-1) + \sum (r^2 - r)$

$= nx^2 + 2x(n^2) + 4\left(\frac{(n)(n+1)(2n+1)}{6} - \frac{n(n+1)}{2}\right)$

$\Rightarrow nx^2 + 2xn^2 + \frac{4n(n^2-1)}{3} = \frac{8n}{3}$

$\Rightarrow x^2 + 2nx + \frac{4(n^2-1)}{3} - \frac{8}{3} = 0$

$\Rightarrow 3x^2 + 6nx + 4n^2 - 12 = 0$

$|\alpha - \beta| = 2$

$\Rightarrow (\alpha + \beta)^2 - 4\alpha\beta = 4$

$\Rightarrow 4n^2 - 4\left(\frac{4n^2 - 12}{3}\right) = 4$

$\Rightarrow 12n^2 - 16n^2 + 48 - 12 = 0$

$\Rightarrow 4n^2 = 36$

$\Rightarrow n^2 = 9$

$\Rightarrow n = 3$


16. Let $y^2 = 12x$ be the parabola with its vertex at O . Let P be a point on the parabola and A be a point on the x -axis such that $\angle OPA = 90^\circ$. Then the locus of the centroid of such triangles OPA is :

(1) $y^2 - 9x + 6 = 0$ (2) $y^2 - 4x + 8 = 0$

(3) $y^2 - 6x + 4 = 0$ (4) $y^2 - 2x + 8 = 0$

Answer (4)

Sol.



$M_{OP} \cdot M_{PA} = -1$

$\frac{2}{t} \cdot \frac{6t}{3t^2 - x_A} = -1$

$12 = x_A - 3t^2$

$x_A = 3t^2 = -12$

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY MAYANK
AIR 36



RUJUL GARG
AIR 41



ARUSH ANAND
AIR 64



SHREYAS LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH A GUPTA
AIR 15
Telangana Topper
100 Overall



$$h = \frac{x_A + 3t^2}{3}, k = \frac{6t}{3}$$

$$3h = 3t + 3t^2 - 12 \quad k = 2t$$

$$h = 2t^2 - 4$$

$$h = 2\left(\frac{k}{2}\right)^2 - 4$$

$$2h = k^2 - 8$$

$$h \rightarrow x \quad k \rightarrow y$$

$$2x = y^2 - 8$$

$$y^2 - 2x - 8 = 0$$

17. For a triangle ABC , let $\vec{p} = \overline{BC}$, $\vec{q} = \overline{CA}$ and

$\vec{r} = \overline{BA}$. If $|\vec{p}| = 2\sqrt{3}$, $|\vec{q}| = 2$ and $\cos\theta = \frac{1}{\sqrt{3}}$, where

θ is the angle between \vec{p} and \vec{q} , then

$|\vec{p} \times (\vec{q} - 3\vec{r})|^2 + 3|\vec{r}|^2$ is equal to :

(1) 200


(2) 220

(3) 410

(4) 340

Answer (1)

Sol.



$$\therefore \vec{p} + \vec{q} = \vec{r} \dots (1)$$

Squaring both sides we get

$$|\vec{p}|^2 + |\vec{q}|^2 + 2\vec{p} \cdot \vec{q} = |\vec{r}|^2$$

$$12 + 4 + 8\sqrt{3} \cdot \frac{1}{\sqrt{3}} = |\vec{r}|^2$$

$$\therefore |\vec{r}| = \sqrt{24}$$

from equation (1) : $\vec{p} \times (\vec{p} + \vec{q}) = \vec{p} \times \vec{r}$.

$$\vec{p} \times \vec{q} = \vec{p} \times \vec{r}$$

$$\therefore \vec{p} \times (\vec{q} - 3\vec{r}) = \vec{p} \times \vec{q} - 3\vec{p} \times \vec{r}$$

$$= -2\vec{p} \times \vec{q}$$

$$\therefore |\vec{p} \times (\vec{q} - 3\vec{r})|^2 + 3|\vec{r}|^2$$

$$= 4.|\vec{p} \times \vec{q}|^2 + 3 \times 24.$$

$$= 4(|\vec{p}|^2 |\vec{q}|^2 - (\vec{p} \cdot \vec{q})^2) + 3 \times 24$$

$$= 4(12 \cdot 4 - 16) + 72 = 200$$

18. Let α and β be the roots of the equation

$$x^2 + 2ax + (3a + 10) = 0$$

such that $\alpha < 1 < \beta$. Then the set of all possible values of a is:

(1) $\left(-\infty, -\frac{11}{5}\right)$


(2) $(-\infty, -3)$

(3) $\left(-\infty, -\frac{11}{5}\right) \cup (5, \infty)$

(4) $(-\infty, -2) \cup (5, \infty)$

Answer (1)

Sol. $x^2 + 2ax + (3a + 10) = 0$



$$\Rightarrow f(1) < 0$$

$$\Rightarrow 1 + 2a + 3a + 10 < 0$$

$$5a + 11 < 0$$

$$\boxed{a < -\frac{11}{5}}$$

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



19. If the area of the region $\{(x, y) : 1 - 2x \leq y \leq 4 - x^2, x \geq 0, y \geq 0\}$ is


$\frac{\alpha}{\beta}, \alpha, \beta \in \mathbf{N}, \text{gcd}(\alpha, \beta) = 1$, then the value of $(\alpha + \beta)$

is :

- (1) 67 (2) 73
(3) 85 (4) 91

Answer (2)

Sol. $1 - 2x \leq y \leq 4 - x^2$



$$\text{Area} = \int_0^2 (4 - x^2) dx - \frac{1}{2} \times \frac{1}{2} \times 1$$

$$4x - \frac{x^3}{3} \Big|_0^2 - \frac{1}{4}$$

$$8 - \frac{8}{3} - \frac{1}{4}$$

$$\frac{61}{12} = \frac{\alpha}{\beta} \therefore \alpha + \beta = 73$$

20. Let the line L pass through the point $(-3, 5, 2)$ and make equal angles with the positive coordinate axes. If the distance of L from the point $(-2, r, 1)$ is

$\sqrt{\frac{14}{3}}$, then the sum of all possible values of r is :

- (1) 16 (2) 10
(3) 12 (4) 6

Answer (2)

Sol. $A = (-3, 5, 2), P(-2, r, 1)$

$$L : \vec{r} = \langle -3, 5, 2 \rangle + t \langle 1, 1, 1 \rangle$$

$$\text{dist} = \frac{|AP \times \vec{d}|}{|\vec{d}|}$$

$$\vec{AP} = \langle 1, r - 5, -1 \rangle \quad \vec{d} = \langle 1, 1, 1 \rangle$$

$$|AP \times \vec{d}|^2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & r - 5 & -1 \\ 1 & 1 & 1 \end{vmatrix} = \langle r - 4, -2, 6 - r \rangle$$

$$\Rightarrow \sqrt{\frac{14}{3}} = \frac{\sqrt{(r - 4)^2 + 4 + (6 - r)^2}}{\sqrt{3}}$$

SOBS

$$14 = r^2 + 16 - 8r + 4 + 36 + r^2 - 12r$$

$$2r^2 - 20r + 56 = 14$$

$$2r^2 - 20r + 42 = 0$$

$$r^2 - 10r + 21 = 0$$

$$r = 3 \text{ or } 7$$

$$\text{Sum} = 7 + 3 = 10$$

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

21. If P is a point on the circle $x^2 + y^2 = 4$, Q is a point on the straight line $5x + y + 2 = 0$ and $x - y + 1 = 0$ is the perpendicular bisector of PQ , then 13 times the sum of abscissa of all such points P is

Answer (02.00)

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall




KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



Sol.



$$\Rightarrow \frac{x-a}{1} = \frac{y-b}{-1} = -2 \frac{(a-b+1)}{2}$$

$$= -a + b - 1$$

$$\Rightarrow x = b-1, y = a+1$$

$$\Rightarrow ((b-1), (a+1)) \text{ lie on } 5x + y + 2 = 0$$

$$\Rightarrow 5(b-1) + (a+1) + 2 = 0$$

$$a + 5b - 2 = 0$$

and also $a^2 + b^2 = 4$

$$\Rightarrow a^2 + \left(\frac{2-a}{5}\right)^2 = 4$$

$$\Rightarrow 25a^2 + (a^2 - 4a + 4) = 100$$

$$\Rightarrow 26a^2 - 4a - 96 = 0$$

$$\text{Sum of all } a = \frac{2}{13}$$

$$\Rightarrow \frac{13 \times 2}{13} = 2$$

22. If $\int_0^1 4 \cot^{-1}(1-2x+4x^2) dx = a \tan^{-1}(2) - b \log_e(5)$,

where $a, b \in \mathbb{N}$, then $(2a+b)$ is equal to _____.

Answer (9)

Sol. $I = 4 \int_0^1 \cot^{-1}(1-2x+4x^2) dx$

Since $1-2x+4x^2 = \left(2x - \frac{1}{2}\right)^2 + \frac{3}{4} > 0$

$$\Rightarrow I = 4 \int_0^1 \tan^{-1}\left(\frac{1}{1+2x(2x-1)}\right) dx$$

$$I = 4 \int_0^1 (\tan^{-1}(2x) - \tan^{-1}(2x-1)) dx$$

$$= 4 \int_0^1 \tan^{-1}(2x) dx - 4 \int_0^1 \tan^{-1}(2x-1) dx$$

for $\int_0^1 \tan^{-1}(2x-1) dx = \int_{-1}^1 \frac{\tan^{-1}(t)}{2} dt = 0$

$$\Rightarrow I = 4 \int_0^1 \tan^{-1}(2x) dx$$

$$= 4(x \tan^{-1} 2x) \Big|_0^1 - 4 \int_0^1 \frac{x}{1+4x^2} \times 2 dx$$

$$= 4 \tan^{-1} 2 - 4 \frac{\ln(1+4x^2)}{4} \Big|_0^1$$

$$= 4 \tan^{-1} 2 - \ln 5 \Rightarrow a = 4, b = 1$$

$$\Rightarrow 2(4) + 1 = 9$$

23. If $\left(\frac{1}{{}^{15}C_0} + \frac{1}{{}^{15}C_1}\right) \left(\frac{1}{{}^{15}C_1} + \frac{1}{{}^{15}C_2}\right) \dots \left(\frac{1}{{}^{15}C_{12}} + \frac{1}{{}^{15}C_{13}}\right)$,

$$= \frac{\alpha^{13}}{{}^{14}C_0 {}^{14}C_1 \dots {}^{14}C_{12}}$$

then 30α is equal to _____

Answer (32)

Sol. Notice that

$$\frac{1}{{}^nC_r} + \frac{1}{{}^nC_{r+1}} = \frac{{}^nC_{r+1} + {}^nC_r}{{}^nC_r \cdot {}^nC_{r+1}} = \frac{{}^{n+1}C_{r+1}}{{}^nC_r \cdot {}^nC_{r+1}}$$

$$= \frac{{}^{n+1}C_r + {}^nC_r}{{}^nC_r \cdot {}^nC_{r+1}} = \frac{(n+1)}{(r+1) \cdot \frac{n}{r+1} \cdot {}^{n-1}C_r} = \frac{n+1}{n \cdot {}^{n+1}C_r}$$

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



$$\begin{aligned} \therefore & \left(\frac{1}{{}^{15}C_0} + \frac{1}{{}^{15}C_1} \right) \left(\frac{1}{{}^{15}C_1} + \frac{1}{{}^{15}C_2} \right) \dots \left(\frac{1}{{}^{15}C_{12}} + \frac{1}{{}^{15}C_{13}} \right) \\ &= \frac{16}{15 \cdot {}^{14}C_0} \cdot \frac{16}{15 \cdot {}^{14}C_1} \dots \frac{16}{15 \cdot {}^{14}C_{12}} \\ &= \frac{\left(\frac{16}{15} \right)^{13}}{{}^{14}C_0 \cdot {}^{14}C_1 \cdot {}^{14}C_2 \dots {}^{14}C_{12}} \end{aligned}$$

$$\therefore \alpha = \frac{16}{15}$$

$$\therefore 30\alpha = 32.$$

24. Let the maximum value of $(\sin^{-1}x)^2 + (\cos^{-1}x)^2$ for $x \in \left[-\frac{\sqrt{3}}{2}, \frac{1}{\sqrt{2}} \right]$ be $\frac{m}{n}\pi^2$, where $\gcd(m, n) = 1$.

Then $m + n$ is equal to _____.

Answer (65)

Sol.

$$\begin{aligned} & (\sin^{-1}x)^2 + (\cos^{-1}x)^2 \\ &= (\sin^{-1}x + \cos^{-1}x)^2 - 2\sin^{-1}x \left(\frac{\pi}{2} - \sin^{-1}x \right) \\ &= \frac{\pi^2}{4} + 2(\sin^{-1}x)^2 - \pi\sin^{-1}x \\ &= \frac{\pi^2}{4} + 2 \left((\sin^{-1}x) - \frac{\pi}{4} \right)^2 - 2 \times \frac{\pi^2}{16} \\ &= \frac{\pi^2}{8} + 2(\sin^{-1}x - \frac{\pi}{4})^2 \end{aligned}$$

For maximum,

Since,

$$\sin^{-1}x \in \left[-\frac{\pi}{3}, \frac{\pi}{4} \right] \text{ at}$$

$$\sin^{-1}x = -\frac{\pi}{3}$$

$$\Rightarrow \text{maximum value} = \frac{29}{36}\pi^2$$

$$\Rightarrow \frac{m}{n} = \frac{29}{36} \Rightarrow m + n = 65$$

25. Let $[.]$ denote the greatest integer function and

$f(x) = \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{k=1}^n \left[\frac{k^2}{3^x} \right]$. Then $12 \sum_{j=1}^{\infty} f(j)$ is equal to _____.

Answer (02.00)

Sol.

$$\begin{aligned} f(x) &= \lim_{n \rightarrow \infty} \left(\frac{1}{n^3} \sum_{k=1}^n \left[\frac{k^2}{3^x} \right] \right) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{k=1}^n \left(\frac{k^2}{3^x} \right) - \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{k=1}^n \left\{ \frac{k^2}{3^x} \right\} \\ f(x) &= \lim_{n \rightarrow \infty} \frac{1}{n^3} \times \frac{n(n+1)(2n+1)}{6 \times 3^x} - 0 \\ \Rightarrow f(x) &= \frac{1}{3^{x+1}} \\ 12 \sum_{j=1}^{\infty} f(j) &= 12 \left(\frac{1}{3^2} + \frac{1}{3^3} + \frac{1}{3^4} + \dots \right) \\ &= 12 \left(\frac{\frac{1}{9}}{1 - \frac{1}{3}} \right) = \frac{12}{9-3} = 2 \end{aligned}$$

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall




PHYSICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer :

26. An infinitely long straight wire carrying current I is bent in a planner shape as shown in the diagram. The radius of the circular part is r . The magnetic field at the centre O of the circular loop is :




- (1) $-\frac{\mu_0 I}{2\pi r}(\pi+1)\hat{i}$ (2) $-\frac{\mu_0 I}{2\pi r}(\pi-1)\hat{i}$
 (3) $\frac{\mu_0 I}{2\pi r}(\pi+1)\hat{i}$ (4) $\frac{\mu_0 I}{2\pi r}(\pi-1)\hat{i}$

Answer (2)

Sol. $\frac{\mu_0 I}{2\pi r}\hat{i} - \frac{\mu_0 I}{2r}\hat{i}$
 $\Rightarrow \vec{B} = \frac{-\mu_0 I}{2\pi r}(\pi-1)\hat{i}$

27. The pulley shown in figure is made using a thin rim and two rods of length equal to diameter of the rim. The rim and each rod have a mass of M . Two blocks of mass of M and m are attached to two ends of a light string passing over the pulley, which is hinged to rotate freely in vertical plane about its center. The magnitudes of the acceleration experienced by the blocks is _____
 (assume no slipping of string on pulley).



- (1) $\frac{(M-m)g}{2M+m}$ (2) $\frac{(M-m)g}{\left[\left(\frac{13}{6}\right)M+m\right]}$
 (3) $\frac{(M-m)g}{M+m}$ (4) $\frac{(M-m)g}{\left[\left(\frac{8}{3}\right)M+m\right]}$

Answer (4)

Sol. $\tau = (M-m)gR$
 $\alpha = \frac{(M-m)gR}{MR^2 + mR^2 + MR^2 + \frac{2M4R^2}{12}}$
 $\Rightarrow \alpha = \frac{(M-m)gR}{\left(\frac{8M}{3} + m\right)R}$
 $\Rightarrow a = \alpha R = \frac{(M-m)g}{\left(\frac{8}{3}M + m\right)}$

28. The energy of an electron in an orbit of the Bohr's atom is $-0.04E_0$ eV where E_0 is the ground state energy. If L is the angular momentum of the electron in this orbit and h is the Planck's constant, then $\frac{2\pi L}{h}$ is _____.
 (1) 2 (2) 5
 (3) 4 (4) 6

Answer (2)

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY MAYANK AIR 36		RUJUL GARG AIR 41		ARUSH ANAND AIR 64	
-------------------------------	--	-----------------------------	--	------------------------------	--

JEE (MAIN)

SHREYAS LOHIYA AIR 6 Uttar Pradesh Topper 100		KUSHAGRA BAINGAHA AIR 7 Uttar Pradesh Topper 100		HARSSH A GUPTA AIR 15 Telangana Topper 100	
---	--	--	--	--	--

Sol. $E = \frac{-E_0}{n^2} = \frac{-E_0}{25}$

$\Rightarrow n = 5$

So $\frac{h5}{2\pi} = L$

$\Rightarrow \frac{2\pi L}{h} = 5$

29. Consider two identical metallic spheres of radius R each having charge Q and mass m . Their centers have an initial separation of $4R$. Both the spheres are given an initial speed of u towards each other. The minimum value of u , so that they can just touch each other is :

(Take $k = \frac{1}{4\pi\epsilon_0}$ and assume $kQ^2 > Gm^2$ where G

is the Gravitational constant)

(1) $\sqrt{\frac{kQ^2}{2mR} \left(1 - \frac{Gm^2}{2kQ^2}\right)}$ (2) $\sqrt{\frac{kQ^2}{2mR} \left(1 - \frac{Gm^2}{kQ^2}\right)}$

(3) $\sqrt{\frac{kQ^2}{4mR} \left(1 - \frac{Gm^2}{kQ^2}\right)}$ (4) $\sqrt{\frac{kQ^2}{4mR} \left(1 + \frac{Gm^2}{kQ^2}\right)}$


Answer (3)

Sol. $\frac{KQ^2}{4R} - \frac{Gm^2}{4R} + \frac{1}{2}mu^2 \times 2 = \frac{KQ^2}{2R} - \frac{Gm^2}{2R}$

$\Rightarrow mu^2 = \frac{KQ^2}{4R} - \frac{Gm^2}{4R}$

$\Rightarrow u = \sqrt{\frac{KQ^2}{4mR} \left(1 - \frac{Gm^2}{KQ^2}\right)}$

30. A large drum having radius R is spinning around its axis with angular velocity ω , as shown in figure. The minimum value of ω so that a body of mass M remains stuck to the inner wall of the drum, taking the coefficient of friction between the drum surface and mass M as μ , is :



(1) $\sqrt{\frac{2g}{\mu R}}$

(2) $\sqrt{\frac{\mu g}{R}}$

(3) $\sqrt{\frac{g}{\mu R}}$

(4) $\sqrt{\frac{g}{2\mu R}}$

Answer (3)

Sol. $N = m\omega^2 R$

$mg \leq \mu m\omega^2 R$

so $\omega \geq \sqrt{\frac{g}{\mu R}}$

31. The kinetic energy of a simple harmonic oscillator is oscillating with angular frequency of 176rad/s. The frequency of this simple harmonic oscillator is

_____ Hz. [take $\pi = \frac{22}{7}$]

(1) 28

(2) 88

(3) 176

(4) 14

Answer (4)

Sol. We know that energy term oscillates with twice the frequency of SHM.

so $\omega_0 = \frac{176}{2}$

$\Rightarrow f = \frac{88 \times 7}{2 \times 22}$

$\Rightarrow f = 14$

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY MAYANK
AIR 36



RUJUL GARG
AIR 41



ARUSH ANAND
AIR 64



SHREYAS LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH A GUPTA
AIR 15
Telangana Topper
100 Overall



32. A river of width 200 m is flowing from west to east with a speed of 18 km/h. A boat, moving with speed of 36 km/h in still water, is made to travel one-round trip (bank to bank of the river). Minimum time taken by the boat for this journey and also the displacement along the river bank are ____ and ____ respectively.

- (1) 40 s and 200 m
- (2) 20 s and 100 m
- (3) 40 s and 100 m
- (4) 40 s and 0 m

Answer (1)

Sol. $\Delta t = \frac{400}{10} = 40 \text{ sec}$

and $\Delta x = 5 \times 40 = 200 \text{ m}$

33. A capacitor C is first charged fully with potential difference of V_0 and disconnected from the battery. The charged capacitor is connected across an inductor having inductance L . In t s 25% of the initial energy in the capacitor is transferred to the inductor. The value of t is ____ s.

- (1) $\frac{\pi\sqrt{LC}}{3}$
- (2) $\frac{\pi\sqrt{LC}}{6}$
- (3) $\frac{\pi\sqrt{LC}}{2}$
- (4) $\pi\sqrt{\frac{LC}{2}}$


Answer (2)

Sol. $q = q_0 \cos(\omega t)$

$$q_1^2 = \frac{75}{100} q_0^2 \Rightarrow q_1 = \frac{\sqrt{3}}{2} q_0$$

So, $\omega t = \frac{\pi}{6} \Rightarrow t = \frac{\pi\sqrt{LC}}{6}$

34. The charge stored by the capacitor C in the given circuit in the steady state is ____ μC



- (1) 5
- (2) 12.5
- (3) 10
- (4) 7.5

Answer (3)

Sol. $I = \frac{2.5}{5} = \frac{1}{2} \text{ A}$

$$\Delta V = 4 \times \frac{1}{2} \text{ volts}$$

$$q = C\Delta V = 10 \mu\text{F}$$

35. Keeping the significant figures in view, the sum of the physical quantities 52.01 m, 153.2 m and 0.123 m is

- (1) 205 m
- (2) 205.3 m
- (3) 205.333 m
- (4) 205.33 m

Answer (2)

Sol. After decimal least number of digits to be considered.

36. A battery with EMF E and internal resistance r is connected across a resistance R . The power consumption in R will be maximum when

- (1) $R = r$
- (2) $R = 2r$
- (3) $R = r/2$
- (4) $R = \sqrt{2}r$

Answer (1)

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



Sol. $I = \frac{E}{R+r}$

$$P = \frac{E^2}{(R+r)} \cdot R$$

For $P_{\max} \Rightarrow R = r$

37. Surface tension of two liquids (having same densities), T_1 and T_2 , are measured using capillary rise method utilizing two tubes with inner radii of r_1 and r_2 where $r_1 > r_2$. The measured liquid heights in these tubes are h_1 and h_2 respectively. [Ignore the weight of the liquid about the lowest point of meniscus]. The heights h_1 and h_2 and surface tensions T_1 and T_2 satisfy the relation:

- (1) $h_1 = h_2$ and $T_1 = T_2$ (2) $h_1 > h_2$ and $T_1 < T_2$
 (3) $h_1 > h_2$ and $T_1 = T_2$ (4) $h_1 < h_2$ and $T_1 = T_2$

Answer (4)

Sol. $h = \frac{2T \cos \theta}{\rho g r}$

$$\text{So, } \frac{h_1}{h_2} = \frac{T_1 r_2}{T_2 r_1}$$

$$\Rightarrow \frac{r_1}{r_2} = \frac{T_1 h_2}{T_2 h_1}$$

Since $r_1 > r_2$

So, for $T_1 = T_2$

$$h_1 < h_2$$

38. The r.m.s. speed of oxygen molecules at 47°C is equal to that of the hydrogen molecules kept at _____ $^\circ\text{C}$. (Mass of oxygen molecule/mass of hydrogen molecule = 32/2)

- (1) -253 (2) -100
 (3) -235 (4) -20

Answer (1)

Sol. $V_{\text{r.m.s.}} = \sqrt{\frac{3KT}{m}}$

$$\Rightarrow V_{\text{O}_2} = \sqrt{\frac{3K \times 320}{32 m_0}}$$

And $V_{\text{N}_2} = \sqrt{\frac{3KT}{2m_0}}$

$$\Rightarrow T = 20\text{ K} = -253^\circ\text{C}$$

39. Two cars A and B each of mass 10^3 kg are moving on parallel tracks separated by a distance of 10 m, in same direction with speeds 72 km/h and 36 km/h. The magnitude of angular momentum of car A with respect to car B is _____ J.s.


- (1) 3×10^5 (2) 3.6×10^5
 (3) 10^5 (4) 2×10^5

Answer (3)

Sol. $V_A = 20$ m/s; $V_B = 10$ m/s

So $L_{A/B} = 10^3 \times 10 \times 10 = 10^5$ J.s.

40. Two known resistances of $R \Omega$ and $2R \Omega$ and one unknown resistance $X \Omega$ are connected in a circuit as shown in the figure. If the equivalent resistance between points A and B in the circuit is $X \Omega$, then the value of X is _____ Ω .



- (1) $(\sqrt{3}-1)R$ (2) $2(\sqrt{3}-1)R$
 (3) $(\sqrt{3}+1)R$ (4) R

Answer (1)

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY MAYANK
AIR 36



RUJUL GARG
AIR 41



ARUSH ANAND
AIR 64



SHREYAS LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



JEE (MAIN)

KUSHAGRA BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH A GUPTA
AIR 15
Telangana Topper
100 Overall



Sol. $\frac{1}{R} + \frac{1}{2R+x} = \frac{1}{x}$
 $\Rightarrow x^2 + 2Rx - 2R^2 = 0$
 $\Rightarrow x = R(\sqrt{3} - 1)$

41. Given below are two statements:

Statement-I: In a Young's double slit experiment, the angular separation of fringes will increase as the screen is moved away from the plane of the slits.

Statement-II: In a Young's double slit experiment, the angular separation of fringes will increase when monochromatic source is replaced by another monochromatic source of higher wavelength.

In the light of the above statements, choose the correct answer from the options given below :


- (1) Statement-I is false but statement-II is true
- (2) Both Statement-I and statement-II are false
- (3) Both Statement-I and statement-II are true
- (4) Statement-I is true but statement-II is false

Answer (1)

Sol. $\omega = \frac{\lambda D}{d}$

Angular fringe width = $\frac{\lambda}{d}$.

42. The total length of potentiometer wire AB is 50 cm in the arrangement as shown in figure. If P is the point where the galvanometer shows zero reading then the length AP is _____ cm.



- (1) 30
- (2) 25
- (3) 20
- (4) 15

Answer (1)

Sol. ∴ galvanometer shows zero

So, $6 \times R_{PB} = 4R_{AP}$

$\frac{I_{AP}}{50 - I_{AP}} = \frac{3}{2}$

$I_{AP} = 30$ cm

43. A body of mass 2 kg is moving along x-direction such that its displacement as function of time is given by $x(t) = \alpha t^2 + \beta t + \gamma$ m, where $\alpha = 1 \text{ m/s}^2$, $\beta = 1 \text{ m/s}$ and $\gamma = 1 \text{ m}$. The work done on the body during the time interval $t = 2 \text{ s}$ to $t = 3 \text{ s}$, is _____ J.

- (1) 49
- (2) 42
- (3) 12
- (4) 24

Answer (4)

Sol. $x = \alpha t^2 + \beta t + \gamma$

$v = 2\alpha t + \beta$

$v(3) = 6\alpha + \beta = 7 \text{ m/s}$

$v(2) = 4\alpha + \beta = 5 \text{ m/s}$

$\omega = \Delta kt$
 $= \frac{1}{2} \cdot 2 \cdot (7^2 - 5^2)$

$= 24$

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall




KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall




44. As shown in the diagram, when the incident ray is parallel to base of the prism, the emergent ray grazes along the second surface.



If refractive index of the material of prism is $\sqrt{2}$, the angle θ of prism is.

- (1) 75°
- (2) 60°
- (3) 45°
- (4) 90°

Answer (2)



$$\sin 45^\circ = \sqrt{2} \sin r_1$$

$$r_1 = 30^\circ$$

$$\sqrt{2} \sin r_2 = 1$$

$$r_2 = 45^\circ$$

$$A = r_1 + r_2$$

$$= 75^\circ$$

$$\theta = 180^\circ - 45^\circ - 75^\circ$$

$$= 60^\circ$$

45. A spherical body of radius r and density σ falls freely through a viscous liquid having density ρ and viscosity η and attains a terminal velocity v_0 . Estimated maximum error in the quantity η is :

(Ignore errors associated with σ , ρ and g , gravitational acceleration)

(1) $\frac{2\Delta r}{r} + \frac{\Delta v_0}{v_0}$

(2) $2\frac{\Delta r}{r} - \frac{\Delta v_0}{v_0}$

(3) $2\left[\frac{\Delta r}{r} - \frac{\Delta v_0}{v_0}\right]$

(4) $2\left[\frac{\Delta r}{r} + \frac{\Delta v_0}{v_0}\right]$

Answer (1)

Sol. $V_0 = \frac{2r^2g}{9\eta}(\sigma - \rho)$

$$\frac{V_0\eta}{r^2} = \text{constant}$$

$$\frac{\Delta\eta}{\eta} = \frac{2\Delta r}{r} + \frac{\Delta V_0}{V_0}$$

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

46. The terminal velocity of a metallic ball of radius 6 mm in a viscous fluid is 20 cm/s. The terminal velocity of another ball of same material and having radius 3 mm in the same fluid will be _____ cm/s.

Answer (5)

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY MAYANK
AIR 36



RUJUL GARG
AIR 41



ARUSH ANAND
AIR 64



JEE (MAIN)

SHREYAS LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH A GUPTA
AIR 15
Telangana Topper
100 Overall



Sol. $V_T \propto r^2$

$$V_T = \left(\frac{3}{6}\right)^2 \cdot 20$$

$$= 5 \text{ cm/s}$$

47. A diatomic gas ($\gamma = 1.4$) does 100 J of work when it is expanded isobarically. Then the heat given to the gas _____ J.

Answer (350)

Sol. $Q = \Delta V + W$

$$nC_p \Delta T = nC_v \Delta T + nR \Delta T$$

$$Q = \frac{7}{2} W = 350$$

48. In a Young's double slit experiment set up, the two slits are kept 0.4 mm apart and screen is placed at 1 m from slits. If a thin transparent sheet of thickness 20 μm is introduced in front of one of the slits then center bright fringe shifts by 20 mm on the screen. The refractive index of transparent sheet is given by $\frac{\alpha}{10}$, where α is _____.

Answer (14)

Sol. $d = 0.4 \text{ mm}$ $D = 1 \text{ m}$

$$(\mu - 1)t = \frac{yd}{D}$$

$$\mu = 1 + \frac{yd}{tD}$$

$$= 1.4 = \frac{14}{10}$$

49. A particle having electric charge $3 \times 10^{-19} \text{ C}$ and mass $6 \times 10^{-27} \text{ kg}$ is accelerated by applying an electric potential of 1.21 V. Wavelength of the matter wave associated with the particle is $\alpha \times 10^{-12} \text{ m}$. The value of α is _____.

(Take Planck's constant = $6.6 \times 10^{-34} \text{ J.s}$)

Answer (10)

Sol. $T = \frac{h}{\sqrt{2mkE}}$

$$= \frac{6.6 \times 10^{-34}}{\sqrt{2 \times 6 \times 10^{-27} \times 3 \times 10^{-19} \times 1.21}}$$

$$= \frac{6.6}{6.6} \times 10^{-11}$$

$$= 10 \times 10^{-12}$$

50. An electromagnetic wave of frequency 100 MHz propagates through a medium of conductivity, $\sigma = 10 \text{ mho/m}$. The ratio of maximum conduction current density to maximum displacement current density is _____.

[Take $\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ Nm}^2 / \text{C}^2$]

Answer (1800)

Sol. $\frac{J_c}{J_d} = \frac{\sigma}{\omega\epsilon_0}$

$$= \frac{2\sigma}{f \cdot 4\pi\epsilon_0}$$

$$= \frac{2 \times 10 \times 9 \times 10^9}{10^8}$$

$$= 1800$$

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



CHEMISTRY

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer:

51. Given below are some of the statements about Mn and Mn_2O_7 . Identify the correct statements.

- A. Mn forms the oxide Mn_2O_7 , in which Mn is in its highest oxidation state.
- B. Oxygen stabilizes the Mn in higher oxidation states by forming multiple bonds with Mn.
- C. Mn_2O_7 is an ionic oxide.
- D. The structure of Mn_2O_7 consists of one bridged oxygen.


Choose the **correct** answer from the options given below :

- (1) A, B and D only
- (2) A, B and C only
- (3) A, C and D only
- (4) A, B, C and D

Answer (1)

Sol. (A) In Mn_2O_7 Mn is in +7 highest oxidation state.

- (B) Correct
- (C) Mn_2O_7 is covalent



52. Given below are two statements :

Statement I: Crystal Field Stabilization Energy (CFSE) of $[Cr(H_2O)_6]^{2+}$ is greater than that of $[Mn(H_2O)_6]^{2+}$.

Statement II: Potassium ferricyanide has a greater spin-only magnetic moment than sodium ferrocyanide.

In the light of the above statements, choose the **correct** answer from the options given below :

- (1) Statement I is false but Statement II is true
- (2) Both Statement I and Statement II are true
- (3) Statement I is true but Statement II is false
- (4) Both Statement I and Statement II are false

Answer (2)

Sol. $[Cr(H_2O)_6]^{2+} \Rightarrow t_{2g}^3 e_g^1$

$$\begin{aligned} \text{CFSE} &= (-0.4 \times 3 + 0.6 \times 1) \Delta_0 \\ &= (-1.2 + 0.6) \Delta_0 \\ &= -0.6 \Delta_0 \end{aligned}$$

$[Mn(H_2O)_6]^{2+} \Rightarrow t_{2g}^3 e_g^2$

$$\begin{aligned} \text{CFSE} &= (-0.4 \times 3 + 0.6 \times 2) \Delta_0 \\ &= 0 \end{aligned}$$

S – I is correct

I. $K_3[Fe(CN)_6] \Rightarrow Fe^{3+} \Rightarrow 3d^5$

Unpaired e^- (n) = 1

II. $Na_4[Fe(CN)_6] \Rightarrow Fe^{2+} \Rightarrow 3d^6$

n = 0

$\mu_I > \mu_{II}$

S – II is correct

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100




HARSSH
A GUPTA
AIR 15
Telangana Topper
100




53. Given below are two statements :

Statement I: Compound (X), shown below, dissolves in NaHCO_3 solution and has two chiral carbon atoms




Statement II: Compound (Y), shown below, has two carbons with sp^3 hybridization, one carbon with sp^2 and one carbon with sp hybridization



In the light of the above statements, choose the **correct** answer from the options given below :

- (1) Both Statement I and Statement II are false
- (2) Statement I is false but Statement II is true
- (3) Statement I is true but Statement II is false
- (4) Both Statement I and Statement II are true

Answer (4)



NaHCO_3 solution due to presence of $-\text{COOH}$ group.

It has 2-chiral centre

S - I is correct



Number of sp^3 carbon = 2

Number of sp^2 carbon = 1


Number of sp carbon = 1

54. Aqueous HCl reacts with $\text{MnO}_2(\text{s})$ to form $\text{MnCl}_2(\text{aq})$, $\text{Cl}_2(\text{g})$ and $\text{H}_2\text{O}(\text{l})$. What is the weight (in g) of Cl_2 liberated when 8.7 g of $\text{MnO}_2(\text{s})$ is reacted with excess aqueous HCl solution?

(Given Molar mass in g mol^{-1} $\text{Mn} = 55$, $\text{Cl} = 35.5$, $\text{O} = 16$, $\text{H} = 1$)

- (1) 7.1
- (2) 14.2
- (3) 21.3
- (4) 71

Answer (1)



$$n_{\text{MnO}_2} = \frac{8.7}{87} = 0.1 \text{ mol}$$

mole of $\text{MnO}_2 = \text{mol of Cl}_2$

$$\begin{aligned} \text{Mass of Cl}_2 &= 0.1 \times 71 \text{ g} \\ &= 7.1 \text{ g} \end{aligned}$$

55. Consider the following data:

$$\Delta_f H^\ominus (\text{methane, g}) = -X \text{ kJmol}^{-1}$$

$$\text{Enthalpy of sublimation of graphite} = Y \text{ kJmol}^{-1}$$

$$\text{Dissociation enthalpy of H}_2 = Z \text{ kJmol}^{-1}$$

The bond enthalpy of $\text{C}-\text{H}$ bond is given by :

- (1) $\frac{-X + Y + Z}{4}$
- (2) $\frac{X + Y + 4Z}{2}$
- (3) $\frac{X + Y + 2Z}{4}$
- (4) $X + Y + Z$

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100



HARSSH
A GUPTA
AIR 15
Telangana Topper
100



Answer (3)







$$Y + 2 \times Z - 4BE_{C-H} = -X$$

$$X + Y + 2Z = 4BE_{C-H}$$

$$BE_{CH} = \frac{X + Y + 2Z}{4}$$

56. For a closed circuit Daniell cell, which of the following plots is the accurate one at a given temperature?



Answer (3)

Sol. Standard potential of cell remains constant with time.





57. Given below are four compounds :

- (a) n-propyl chloride
- (b) iso-propyl chloride
- (c) sec-butyl chloride
- (d) neo-pentyl chloride

Percentage of carbon in the one which exhibits optical isomerism is :

- (1) 46
- (2) 52
- (3) 56
- (4) 40

Answer (2)



(c) can show optical isomerism

$$\% C = \frac{12 \times 4}{92.5} \times 100 = 51.89 \approx 52\%$$

58. Given below are two statements :

Statement I : The correct order in terms of bond dissociation enthalpy is $Cl_2 > Br_2 > F_2 > I_2$.

Statement II : The correct trend in the covalent character of the metal halides is $[SnCl_4 > SnCl_2]$, $[PbCl_4 > PbCl_2]$ and $[UF_4 > UF_6]$.

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100% in Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100% in Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100% in Overall



In the light of the above statements, choose the **correct** answer from the options given below :

- (1) Statement I is true but Statement II is false
- (2) Both Statement I and Statement II are false
- (3) Statement I is false but Statement II is true
- (4) Both Statement I and Statement II are true

Answer (1)

Sol. BDE of $F_2 = 158.8 \text{ kJ/mol}$

$Cl_2 = 242.6 \text{ kJ/mol}$

$Br_2 = 192.8 \text{ kJ/mol}$

$I_2 = 151.1 \text{ kJ/mol}$

Order = $Cl_2 > Br_2 > F_2 > I_2$

Statement I is correct

More the charge density on cation more will be covalent character


$SnCl_4(Sn^{4+}) > SnCl_2(Sn^{2+})$

$PbCl_4(Pb^{4+}) > PbCl_2(Pb^{2+})$

$UF_4(U^{4+}) < UF_6(U^{6+})$

Statement II is false.

59. The correct order of reactivity of the following benzyl halides towards reaction with KCN is :



- (1) $b > a > d > c$
- (2) $a > b > d > c$
- (3) $b > a > c > d$
- (4) $a > b > c > d$

Answer (1)

Sol. As only KCN is given so it will follow S_N2 mechanism. Electron donor group via resonance will stabilise transition state.

The rate of S_N2 is:


$b > a > d > c$

60. On heating a mixture of common salt and $K_2Cr_2O_7$ in equal amount along with concentrated H_2SO_4 in a test tube, a gas is evolved. Formula of the gas evolved and oxidation state of the central metal atom in the gas respectively are:

- (1) $Cr_2O_2Cl_2$ and +6
- (2) CrO_2Cl_2 and +6
- (3) $Cr_2O_2Cl_2$ and +3
- (4) CrO_2Cl_2 and +5

Answer (2)

Sol. $K_2Cr_2O_7 + 4NaCl + 6H_2SO_4 \rightarrow$



CrO_2Cl_2 gas is evolved in which Cr is in +6 oxidation state.

61. Match List-I with List-II.

	List-I Pair of Compounds		List-II Type of Isomers
A.	2-Methylpropene and but-1-ene	I.	Stereoisomers
B.	Cis-but-2-ene and trans-but-2-ene	II.	Position isomers
C.	2-Butanol and diethyl ether	III.	Chain isomers
D.	But-1-ene and but-2-ene	IV.	Functional group isomers

Choose the **correct** answer from the options given below :

- (1) A-III, B-I, C-II, D-IV
- (2) A-III, B-I, C-IV, D-II
- (3) A-I, B-IV, C-III, D-II
- (4) A-II, B-I, C-IV, D-III

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100% in Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100% in Overall






HARSSH
A GUPTA
AIR 15
Telangana Topper
100% in Overall



Answer (2)

Sol.



62. The correct increasing order of C-H(A), C-O(B), C=O(C) and C \equiv N(D) bonds in terms of covalent bond length is :

- (1) D < C < B < A (2) A < D < C < B
(3) A < B < C < D (4) D < C < A < B

Answer (2)

Sol. Bond length Values (pm)

C - H	107
C - O	143
C = O	121
C \equiv N	116

Order A < D < C < B

63. Consider the following spectral lines for atomic hydrogen:

- A. First line of Paschen series
B. Second line of Balmer series
C. Third line of Paschen series
D. Fourth line of Bracket series

The correct arrangement of the above lines in ascending order of energy is

- (1) D < A < C < B (2) A < B < C < D
(3) C < D < B < A (4) D < C < A < B

Answer (1)

Sol. 1st line of Paschen series

$$E \propto \left(\frac{1}{3^2} - \frac{1}{4^2} \right) \propto \frac{7}{144}$$

Second line of Balmer series

$$E \propto \left[\frac{1}{2^2} - \frac{1}{4^2} \right] = \frac{12}{64} = \frac{3}{16}$$

Third line of Paschen series

$$E \propto \left[\frac{1}{3^2} - \frac{1}{6^2} \right] \propto \frac{3}{36} \propto \frac{1}{12}$$

Fourth line of Bracket series

$$E \propto \left[\frac{1}{4^2} - \frac{1}{8^2} \right] \propto \frac{3}{64}$$

Order of energy

B > C > A > D

64. Given below are two statements:

Statement I: The correct order in terms of atomic/ionic radii is Al > Mg > Mg²⁺ > Al³⁺.

Statement II: The correct order in terms of the magnitude of electron gain enthalpy is Cl > Br > S > O.

In the light of the above statements, choose the **correct** answer from the options given below.

- (1) Statement I is true but statement II is false
(2) Statement I is false but statement II is true
(3) Both statement I and statement II are false
(4) Both statement I and statement II are true

Answer (2)

Sol. For Al, Mg, Mg²⁺, Al³⁺

Al < Mg

Al³⁺ < Mg²⁺

Statement I false

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100




KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100



HARSSH
A GUPTA
AIR 15
Telangana Topper
100






Consider the above sequence of reactions. The number of bromine atom(s) in the final product (P) will be :

- (1) 3 (2) 1
(3) 6 (4) 5

Answer (4)



70. The **correct** statements are :

- A. Activation energy for enzyme catalysed hydrolysis of sucrose is lower than that of acid catalysed hydrolysis.
B. During denaturation, secondary and tertiary structures of a protein are destroyed but primary structure remains intact.
C. Nucleotides are joined together by glycosidic linkage between C₁ and C₄ carbons of the pentose sugar.

D. Quaternary structure of proteins represents overall folding of the polypeptide chain.

Choose the **correct** answer from the options given below:

- (1) A, C and D Only (2) B and C Only
(3) A and B Only (4) A, B and D Only

Answer (3)

Sol. (E_a)_{acid catalysed} ≈ 6 kJ/mol

(E_a)_{enzyme catalysed} ≈ 2.15 kJ/mol

A correct

Nucleotide are joined together by phosphodiester linkage.

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

71. The first and second ionization constants of H₂X are 2.5 × 10⁻⁸ and 1.0 × 10⁻¹³ respectively. The concentration of X²⁻ in 0.1 M H₂X solution is _____ × 10⁻¹⁵ M. (Nearest Integer)

Answer (100)

Sol. Value of K_{a1} >> K_{a2}

$$\therefore [A^{2-}] = K_{a2} = 10^{-13} = 100 \times 10^{-15}$$

72. The osmotic pressure of a living cell is 12 atm at 300 K. The strength of sodium chloride solution that is isotonic with the living cell at this temperature is _____ g L⁻¹. (Nearest integer)

Given : R = 0.08 L atm K⁻¹ mol⁻¹

Assume complete dissociation of NaCl

(Given : Molar mass of Na and Cl are 23 and 35.5 g mol⁻¹ respectively.)

Answer (15)

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100



HARSSH
A GUPTA
AIR 15
Telangana Topper
100



Sol. $\pi_{\text{NaCl}} = iCRT$

$$12 = 2 \times \frac{n}{V} \times 0.08 \times 300$$

$$12 = 2 \times \frac{W}{58.5 \times V} \times 0.08 \times 300$$

$$14.625 = \frac{W}{V}$$

73. Identify the metal ions among Co^{2+} , Ni^{2+} , Fe^{2+} , V^{3+} and Ti^{2+} having a spin-only magnetic moment value more than 3.0 BM. The sum of unpaired electrons present in the high spin octahedral complexes formed by those metal ions is _____.

Answer (7)

Sol. $\mu > 3.0 \text{ BM}$

$$\sqrt{n(n+2)} > 3$$

$$n \geq 3$$

$$\text{For } \text{Co}^{2+}(\text{d}^7) \quad n = 3$$

$$\text{For } \text{Ni}^{2+}(\text{d}^8) \quad n = 2$$

$$\text{For } \text{Fe}^{2+}(\text{d}^6) \quad n = 4$$

$$\text{For } \text{V}^{3+}(\text{d}^2) \quad n = 2$$

$$\text{For } \text{Ti}^{2+}(\text{d}^2) \quad n = 2$$

Sum of unpaired e^- in high spin octahedral complex = 7

74. A substance 'X' (1.5 g) dissolved in 150 g of a solvent 'Y' (molar mass = 300 g mol⁻¹) led to an elevation of the boiling point by 0.5 K. The relative lowering in the vapour pressure of the solvent 'Y' is _____ $\times 10^{-2}$. (nearest integer)

[Given : K_b of the solvent = 5.0 K kg mol⁻¹]

Assume the solution to be dilute and no association or dissociation of X takes place in solution.

Answer (3)

Sol. $\Delta T_b = K_b m$

$$0.5 = 5 \times \frac{1.5 \times 1000}{M_0 \times 150}$$


$$M_0 = 100 \text{ g}$$

$$\text{RLVP} = X_{\text{solute}}$$

$$\approx \frac{1.5}{100 \times 150} \times 300$$

$$\approx 3 \times 10^{-2}$$

75. MX is a sparingly soluble salt that follows the given solubility equilibrium at 298 K.



If the standard reduction potential for $\text{M}^+(\text{aq}) \xrightarrow{+e^-} \text{M}(\text{s})$ is $(E_{\text{M}^+/\text{M}}^\ominus) = 0.79 \text{ V}$, then the value of the standard reduction potential for the metal/metal insoluble salt electrode $E_{\text{X}^-/\text{MX}(\text{s})/\text{M}}^\ominus$

is _____ mV. (nearest integer)

[Given : $\frac{2.303 RT}{F} = 0.059 \text{ V}$]

Answer (200)

Sol. Cell reaction: $\text{MX} + e^- \longrightarrow \text{M}(\text{s}) + \text{X}^-$

$$E_{\text{X}^-/\text{MX}/\text{M}}^\ominus = E_{\text{M}^+/\text{M}}^\ominus + \frac{0.059}{n} \log K_{\text{sp}}$$

$$E_{\text{X}^-/\text{MX}/\text{M}}^\ominus = 0.79 + \frac{0.059}{1} \log 10^{-10}$$

$$E^\ominus = 0.79 - 0.59$$

$$= 0.2 \text{ V}$$

$$= 200 \text{ mV}$$



Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100% in Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100% in Overall



HARSSH
A GUPTA
AIR 15
Telangana Topper
100% in Overall

