

Date: 23/11/2025

63



Corporate Office: AESL, 3rd Floor, Incuspaze Campus-2, Plot-13, Sector-18,
Udyog Vihar, Gurugram, Haryana-122015

Answers & Solutions

Time : 120 Minute

for

Max. Marks : 216

Indian Association of Physics Teachers (IAPT) National Standard Examination in PHYSICS (NSEP)-2025

INSTRUCTIONS TO CANDIDATES

- (1) There are 60 questions in this paper. Attempt all the 60 Questions.
- (2) Question paper has two parts. In **Part A-1** (Q. No. 1 to 48) each question has four alternatives, out of which **only one** is correct. Choose the correct alternative and fill the appropriate bubble, as shown.

Q. No. 22 a b c d

In **Part A-2** (Q. No. 49 to 60) each question has four alternatives, out of which **any number of alternative (s)** (1, 2, 3 or 4) may be correct. You have to choose ALL correct alternative(s) and fill the appropriate bubble(s), as shown.

Q. No. 54 a b c d

- (3) For **Part A-1**, each correct answer carries **3 marks** whereas 1 mark will be deducted for each wrong answer. In **Part A-2**, you get **6 marks** if all the correct alternatives are marked. No negative marks in this part.

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

Answer (b)

Sol. $P_5 = P_6 = 7.2$

$i = 1.2 \text{ A}$

$V_5 = 6 \text{ V}$


$i_{6,9} = \frac{6}{15} = 0.4 \text{ A}$

$i_4 = 1.6 \text{ A}$

$V_{AB} = 1.6 \times 4 + 6 = 12.4 \text{ V} \Rightarrow \text{correct}$

$P_6 = (0.4)^2 \times 6 = 0.96 \text{ W} \Rightarrow \text{incorrect}$

3. A transparent and homogeneous sphere of glass of radius r is immersed in water (refractive indices of glass and water being ${}_a\mu_g = \frac{3}{2}$ and ${}_a\mu_w = \frac{4}{3}$). The image of a point object O , located at distance d on its axis in front of the sphere, is formed at point I at the same distance d from the sphere on the opposite side as shown.




The distance d is equal to

- (a) $2r$
- (b) $3r$
- (c) $6r$
- (d) $8r$

Answer (d)

Sol. Ray must be parallel to principal axis while passing through sphere.



$$\frac{\mu_2}{v} - \frac{\mu_1}{u} = \frac{\mu_2 - \mu_1}{R}$$

$$1.5 - \frac{4}{-d} = \frac{2 - 3}{r}$$

$$\frac{4}{3d} = \frac{1}{6r}$$

$d = 8r$

1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

4. A certain substance, with a dielectric constant $k = 2.5$ and the dielectric strength $E = 1.8 \times 10^7$ N/C, completely fills the space between the plates of a parallel plate capacitor (with circular plates) of capacitance $C = 72.0$ nF. The minimum diameter of the circular plates, to ensure that the capacitor can withstand a potential difference of $V = 4.0$ kV, is
- (a) 12 cm (b) 24 cm
(c) 48 cm (d) 96 cm

Answer (d)

Sol. Given, $E_{\max} = 1.8 \times 10^7$ N/C

Also, $V = Ed$


$$\Rightarrow d = \frac{4 \times 10^3}{1.8 \times 10^7} \text{m}$$

And, $C = \frac{K\epsilon_0 A}{d}$

$$\Rightarrow 72 \times 10^{-9} = \frac{2.5 \times 8.85 \times 10^{-12}}{4 \times 10^3} \times \frac{\pi d^2}{4} \times 1.8 \times 10^7$$

$$d = 0.96 \text{ m}$$

5. A uniform solid sphere of radius R rolls without slipping on a rough horizontal surface with a forward velocity v of its centre. On its way, it suddenly encounters a small step of height $0.2 R$ as shown. The angular velocity of the sphere just after the impact is [given that the sphere does not bounce back, rather it goes ahead up the step]



- (a) $\frac{v}{7R}$ (b) $\frac{3v}{7R}$
(c) $\frac{6v}{7R}$ (d) $\frac{v}{R}$

Answer (b)

Sol. Conserving angular momentum about point P . Let's ω be angular speed of sphere assuming no slipping at P .

$$mv_{\text{cm}} r + I_{\text{cm}} \omega = I_P \omega$$

$$mv \times 0.8 R + 0.4 mR^2 \frac{v}{R} = 1.4 mR^2 \omega$$

$$\frac{v}{R} \left(\frac{0.8 + 0.4}{1.4} \right) = \omega$$

$$\frac{6v}{7R} = \omega$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

6. The magnetic field (B) produced by the current i flowing through the sides of a square loop of side l , at a point P at distance x from the center of the square, on the axis perpendicular to the plane of the square loop and passing through its center, is

(a) $B = \frac{\mu_0 i}{4\pi} \frac{2\sqrt{2} l^2}{(4x^2 + l^2)\sqrt{2x^2 + l^2}}$

(b) $B = \frac{\mu_0 i}{4\pi} \frac{4\sqrt{2} lx}{(x^2 + l^2)\sqrt{2x^2 + l^2}}$

(c) $B = \frac{\mu_0 i}{4\pi} \frac{4 \times 2\sqrt{2} l^2}{(4x^2 + l^2)\sqrt{2x^2 + l^2}}$

(d) $B = \frac{\mu_0 i}{4\pi} \frac{4\sqrt{2} lx}{(4x^2 + l^2)\sqrt{x^2 + l^2}}$

Answer (c)



Sol. $B_i = \frac{\mu_0 i}{4\pi r} \{\cos \alpha + \cos \beta\}$

$$B_i = \frac{\mu_0 i}{4\pi \sqrt{\left(\frac{l}{2}\right)^2 + x^2}} (2) \frac{\frac{l}{2}}{\sqrt{\left(\frac{l}{2}\right)^2 + x^2}}$$


$B_{\text{net}} = 4B_1 \cos \phi$

$$= 4 \frac{\mu_0 i l}{4\pi \sqrt{l^2 + 4x^2} \left(\frac{1}{2}\right) \sqrt{l^2 + 2x^2} \left(\frac{1}{\sqrt{2}}\right) \sqrt{\left(\frac{l}{2}\right)^2 + x^2}}$$

$$B = \frac{\mu_0 i l^2 (2\sqrt{2}) 4}{(4)\pi (l^2 + 4x^2) \sqrt{l^2 + 2x^2}}$$



7. A linear positive charge distribution, with linear charge density λ coulomb per meter, extends along $+x$ -axis from $x = 0$ to $x = \infty$.



The electric field \vec{E} at any point $P(0, y)$ on the y -axis.

- (a) is proportional to $\frac{\lambda}{y^2}$ irrespective of whether y is positive or negative.
- (b) is always directed away and perpendicular to the line of charge.
- (c) has a vanishing component parallel to the line of charge.
- (d) is directed along a straight line of slope $m = -1$ if y is positive but along a line of slope $m = +1$ if y is negative.

1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA


(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

Answer (d)




$$E_x = \frac{\lambda}{4\pi\epsilon_0 r} (\cos \alpha + \cos \beta)$$

$$E_y = \frac{\lambda}{4\pi\epsilon_0 r} |\sin \alpha - \sin \beta|$$

$$\alpha = 90^\circ, \beta = 0$$

$$\Rightarrow E_x = E_y = \frac{\lambda}{2\pi\epsilon_0 r}$$



8. Imagine a situation, in which an infinite sheet with positive charge $+\sigma$ per unit area lies in the xy -plane and a second infinite sheet with negative charge $-\sigma$ per unit area lies in the yz -plane. The net electric field E at any point (x, y, z) [that does not lie on either of these planes xy or yz] can be expressed as

(a) $\vec{E} = \frac{\sigma}{2\epsilon_0} (-\hat{i} + \hat{k})$

(b) $\vec{E} = \frac{\sigma}{2\epsilon_0} \hat{j}$

(c) $\vec{E} = \frac{\sigma}{2\epsilon_0} \left[-\frac{x}{|x|} \hat{i} + \frac{z}{|z|} \hat{k} \right]$

(d) $\vec{E} = \frac{\sigma}{\epsilon_0} \left[\frac{x}{|x|} \hat{i} - \frac{z}{|z|} \hat{k} \right]$

Answer (c)

Sol. Component of field will be in \hat{k} and \hat{i} direction either with + or - sign depending on quadrant.


If $x > 0$, E_x is -ve.

If $z > 0$, E_z is +ve.

Let's write field in first quadrant.

$$\vec{E} = \frac{\sigma}{2\epsilon_0} (-\hat{i} + \hat{k})$$

In general, $E = \frac{\sigma}{2\epsilon_0} \left(-\frac{x}{|x|} \hat{i} + \frac{z}{|z|} \hat{k} \right)$



1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

9. For the electric field E , in a region of space where a non-uniform, but spherically symmetric distribution of charge has a charge density $\rho(r)$ as $\rho(r) = \rho_0 \left(1 - \frac{r}{R}\right)$ for $r \leq R$, one can say that
- $$\rho(r) = 0 \quad \text{for } r \geq R,$$
- (a) $E = 0$: both at $r = 0$ and $r = R$
- (b) $E \propto r$ for $r < R$ and $E \propto \frac{1}{r^2}$ for $r \geq R$
- (c) The magnitude of E increases with r and reaches its maximum at $r = \frac{2R}{3}$
- (d) The maximum electric field produced by the given charge distribution is $E_{\max} = \frac{\rho_0 R}{3\epsilon_0}$

Answer (c)

Sol. Using Gauss's law

$$\oint \vec{E} \cdot d\vec{A} = \frac{\int \rho dv}{\epsilon_0}$$

$$\theta = \int_{\beta}^{\gamma} \rho_0 \left(1 - \frac{r}{R}\right) 4\pi r^2 dr$$

$$\theta = 4\pi\rho_0 \left\{ \frac{r^3}{3} - \frac{r^4}{4R} \right\}$$

$$E = K 4\pi\rho_0 \left\{ \frac{r}{3} - \frac{r^2}{4R} \right\}$$

Now we can see,

$E \propto r$ for $r < R$

\Rightarrow (a), (b) is wrong

$E \neq 0$ at $r = R$

\Rightarrow (a) is wrong

$$\text{Now } \frac{dE}{dr} = C \left\{ \frac{1}{3} - \frac{R}{2r} \right\}$$

$$\frac{dE}{dr} \text{ is zero at } r = \frac{2R}{3}$$

$$\begin{cases} \text{for } r < \frac{2R}{3} \frac{dE}{dr} > 0 \\ \text{for } r > \frac{2R}{3} \frac{dE}{dr} < 0 \end{cases}$$

Hence E is maximized at $r = \frac{2R}{3}$

$$E_{\max} = \frac{\rho_0}{\epsilon_0} \left\{ \frac{2R}{9} - \frac{4R^2}{36R} \right\} = \frac{\rho_0 R}{9\epsilon_0}$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)


70
NSEC

(Group A & B)

38
NSEP

34
NSEJS


10. A typical network of resistances R_1 and R_2 shown below extends to infinity towards the right. The total resistance $R_{\text{effective}}$ of this network between points A and B is



- (a) $R_{\text{effective}} = R_1 + \sqrt{R_1^2 + 2R_1R_2}$ (b) $R_{\text{effective}} = R_2 + \sqrt{R_1^2 + 2R_1R_2}$
 (c) $R_{\text{effective}} = R_1 + \sqrt{3R_1R_2}$ (d) $R_{\text{effective}} = R_1 + \sqrt{R_2^2 + 2R_1R_2}$

Answer (a)

Sol. Infinite circuit can be reduced to following circuit, where X



$$\frac{R_2 X}{R_2 + X} + 2R_1 = X$$

$$R_2 X = R_2 X + X^2 - 2R_1 R_2 - 2R_1 X$$

$$X^2 - 2R_1 X - 2R_1 R_2 = 0$$

$$X = \frac{2R_1 \pm \sqrt{4R_1^2 + 8R_1 R_2}}{2}$$

$$X = R_1 + \sqrt{R_1^2 + 2R_1 R_2}$$

11. A cylindrical cavity of diameter 'a' exists inside a long solid cylinder of diameter '2a' as shown in figure. Both the cylinder and the cavity are taken to be infinitely long. The axis of the cavity is parallel to the axis of the cylinder and is at a distance $\frac{a}{2}$ from it. A uniform current of current density J (Am^{-2}) flows through the cylinder along its length and not through the cavity. The magnitude of the magnetic field at a point P on the surface of the cylinder lying farthest from the axis of the cavity, is

420

Classroom Students
Qualified in
NSEs 2024-25

1484 Students Scored Above MAS

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)


70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

- (a) $B = \frac{3 \mu_0 J}{8 a}$
- (b) $B = \frac{3}{4} \mu_0 J a$
- (c) $B = \frac{3}{8} \mu_0 J a$
- (d) $B = \frac{5}{12} \mu_0 J a$



Answer (d)

Sol. Let fill the cavity with current density $+J$ and $-J$

$$B_p = B_a - B_a$$

Using Ampere's law, we know

$$B = \frac{\mu_0 J r}{2} \text{ for } r < R$$

$$B = \frac{\mu_0 J R^2}{2r} \text{ } r \geq R$$

$$B_p = \frac{\mu_0 J a}{2} - \frac{\mu_0 J \left(\frac{a}{2}\right)^2}{2 \left(\frac{3a}{2}\right)}$$


$$= \mu_0 J a \left\{ \frac{1}{2} - \frac{1}{12} \right\}$$

$$B = \mu_0 J a \frac{5}{12}$$

12. A thin uniform rod, of length $\ell = 0.200$ m with negligible mass, is attached to the floor by a frictionless hinge at a fixed-point P . A horizontal spring connects the other end of the rod to a vertical wall. The rod is in a uniform magnetic field $B = 0.500$ tesla directed into the plane of paper. There is a current $I = 10.0$ A in the rod in the direction shown. Force constant of the spring is 5.00 N/m. The rod is in equilibrium at $\theta = \tan^{-1} \frac{4}{3}$

Statement (1): Torque on the rod due to magnetic force is 0.1 Nm clockwise

Statement (2): In equilibrium the energy stored in the spring is 0.039 J



1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)


38
NSEP

34
NSEJS

- (a) Statement (1) is correct but statement (2) is wrong
- (b) Statement (1) is wrong but statement (2) is correct
- (c) Both statements (1) and (2) are wrong
- (d) Both statements (1) and (2) are correct

Answer (d)

Sol. Force diagram of rod and balancing torque



We get

$$kx \ell \sin \theta = i \ell B \frac{\ell}{2}$$

$$5 \times x \times \frac{4}{5} = 10 \times \frac{0.2}{2} \times 0.5$$

$$x = \frac{1}{8}$$

$$\text{Potential energy : } U_S = \frac{1}{2} kx^2 = \frac{1}{2} \times 5 \times \frac{1}{64} = 0.039$$

$$\text{Torque : } \tau_B = 10 \times 0.2 \times 0.5 \times \frac{0.2}{2} = \frac{0.2}{2} = 0.1 \text{ Nm}$$

(This torque is about hinge)

13. The electric flux through a certain area of a dielectric medium is $\phi = (8.00 \times 10^3)t^4$ in SI units. The displacement current through that area is 12.5 pA at a time $t = 20.0$ ms. The dielectric constant of the dielectric medium is
- (a) 22.1
 - (b) 5.52
 - (c) 55.2
 - (d) 2.76

Answer (b)

Sol. Displacement current : $i_d = K \epsilon_0 \frac{d\phi_E}{dt}$

$$12.5 \times 10^{-12} = K \times 8.85 \times 10^{-12} \times 8 \times 10^3 \times 4 \times 8000 \times 10^{-9}$$


$$K = 5.52$$



1484 Students Scored Above MAS




14. A thin equi-convex lens of flint glass (refractive index μ_1) is kept coaxially in contact with another thin equi-concave lens of crown glass (refractive index μ_2). The system is completely immersed in water (${}_a\mu_w = \frac{4}{3}$)



Parallel rays of light incident parallel to the principal axis in water are focused by this system at a distance of 24 cm beyond the system. The thickness of the system is negligible. If the radius of curvature of each surface is $R = 20$ cm, the difference ($\mu_1 - \mu_2$) is

- (a) $\frac{2}{9}$ (b) $\frac{3}{9}$
(c) $\frac{4}{9}$ (d) $\frac{5}{9}$

Answer (d)



Sol. Equivalent system :

$$\text{Now, } \frac{1}{f_1} = \left(\frac{\mu_1}{\mu_w} - 1 \right) \left\{ \frac{1}{R} - \left(-\frac{1}{R} \right) \right\}$$

$$\frac{1}{f_2} = \left(\frac{\mu_2}{\mu_w} - 1 \right) \left\{ -\frac{1}{R} - \frac{1}{R} \right\}$$

$$\frac{1}{f_1} = \frac{1}{f_1} + \frac{1}{f_2} = \frac{2}{R} \left\{ \frac{\mu_1 - \mu_2}{\mu_w} \right\} = \frac{1}{24}$$

$$\mu_1 - \mu_2 = \frac{4}{3} \times \frac{1}{24} \times \frac{20}{2} = \frac{80}{144} = \frac{5}{9}$$

15. Two identical large thin metal plates carrying charges $+q_1$ and $+q_2$ ($q_1 > q_2$), respectively, are kept close at a distance d apart and parallel to each other to form a parallel plate capacitor of capacitance C . The potential difference between the plates is

- (a) $\frac{q_1 - q_2}{C}$ (b) $\frac{q_1 - q_2}{2C}$
(c) $\frac{q_1 - q_2}{4C}$ (d) $\frac{q_1 + q_2}{2C}$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

Answer (b)

Sol. Charge distribution shall be

$$\frac{q_1 + q_2}{2} \left| \begin{array}{c} \frac{q_1 - q_2}{2} \\ -\frac{q_1 - q_2}{2} \end{array} \right| \frac{q_1 + q_2}{2}$$

Using, $q = CV$

$$\frac{q_1 - q_2}{2} = CV$$

16. A point mass m moves in a straight line under a retardation kv^2 [where k is a positive constant and v is the instantaneous velocity]. The initial velocity of the point mass is u . The displacement of the point mass at time t is

(a) $\frac{1}{k} \ln(1 + kut)$

(b) $\frac{1}{k} \ln kut$

(c) $k \ln kut$

(d) $\frac{1}{k} \ln(1 - kut)$

Answer (a)

Sol. Using definition of velocity and acceleration

$$\frac{dv}{dt} = -kv^2$$


Integrating we get

$$v = \frac{4}{1 + 4kt}$$

Integrating we get

$$s = \frac{1}{k} \ln(1 + 4kt)$$

17. In the arrangement shown in figure, 'a' represents the magnitude of acceleration of small blocks A and B while 'T' is the tension in the massless string passing over the frictionless and massless pulley. The sum of the masses of blocks A and B is constant. For this system, a linear relationship can be obtained between



(a) a and $\frac{1}{T}$

(b) a and T

(c) a and T^2

(d) T and a^2

Answer (d)

1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA


(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

Sol.




$$a = \frac{(M - m)g - mg}{M}$$

$$m = \frac{M}{2} \left(1 - \frac{a}{g} \right)$$

$$T = \frac{2(M - m).m.g}{M}$$

$$T = \frac{Mg}{2} \left(1 - \frac{a^2}{g^2} \right) g$$

18. A thin uniform circular ring of mass m is rolling without slipping down an inclined plane of inclination 30° with the horizontal. The coefficient of friction between the ring and the surface is μ . The correct statement is



- (a) Linear acceleration of the center of the ring along the plane is $a = \frac{g}{2}$
- (b) Force of friction between the ring and the inclined plane is $F_{\text{friction}} = \frac{mg}{4}$
- (c) The ring keeps rolling for all values of the coefficient of friction $\mu \geq \frac{1}{4}$
- (d) Linear acceleration of the center of the ring along the plane is $a = \frac{g}{3}$

Answer (b)

1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA


(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

Sol.



$$\frac{mg}{2} - f = ma$$

$$f \cdot R = mR^2 \cdot \frac{a}{R}$$

$$\frac{mg}{2} = 2ma$$

$$a = \frac{g}{4}$$


$$f = \frac{mg}{4}$$

$$\mu mg \frac{\sqrt{3}}{2} \geq \frac{mg}{4}$$

$$\mu \geq \frac{1}{2\sqrt{3}}$$

Rolling for all values of $\mu \geq \frac{1}{2\sqrt{3}}$

19. A bullet of mass m can penetrate a target (a heavy block of mass M) up to a distance S , when the target M is held stationary by a stopper P (shown in figure). Up to what distance S' the bullet will penetrate if the block of mass M is free to move (i.e. when the stopper P is removed) on the frictionless surface T .



(a) $S' = S$

(b) $S' = \frac{m}{M} S$

(c) $S' = \frac{m}{m+M} S$

(d) $S' = \frac{M}{M+m} S$

Answer (d)

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

Sol. With stopper, final speed of m will be zero.

Let average friction is f

$$fs = \frac{1}{2}mv_0^2$$

Without stopper



$$mv_0 = (m + M)v$$

$$fs' = \frac{1}{2}mv_0^2 - \frac{1}{2}(m + M)\left(\frac{mv_0}{m + M}\right)^2$$

$$fs' = \frac{1}{2} \frac{mM}{m + M} v_0^2$$

$$S' = \frac{M}{m + M} S$$

20. Knowing that the atomic masses of Al and Mg are respectively ${}^{25}_{13}\text{Al} = 24.990432 u$ and ${}^{25}_{12}\text{Mg} = 24.985839 u$ while electron mass is often expressed as $m_e = 0.511 \text{ MeV}$, the Q value (energy liberated) of the β decay nuclear reaction ${}^{25}\text{Al} \rightarrow {}^{25}\text{Mg} + e^+ + \nu$ in MeV is


- (a) 4.278 (b) 3.767
(c) 3.256 (d) 931.478

Answer (c)

Sol. ${}^{25}\text{Al} \longrightarrow {}^{25}\text{Mg} + e^+ + \nu$

$$\begin{aligned} Q &= (m_{\text{Al}} - m_{\text{Mg}} - 2m_e)c^2 \\ &= (m_{\text{Al}} - m_{\text{Mg}})c^2 - 2m_e c^2 \\ &= 0.004593 \times 931 - 2 \times 0.511 \\ &= 3.25 \text{ MeV} \end{aligned}$$

21. A block of mass m , lying on a rough horizontal plane, is acted upon by a horizontal force P and simultaneously by another force Q acting at an angle θ from the vertical as shown. The block will remain in equilibrium if the coefficient of friction between the block and the surface S is



1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS


(a) At least $\frac{P + Q \sin \theta}{mg + Q \cos \theta}$

(b) At least $\frac{P + Q \cos \theta}{mg + Q \sin \theta}$

(c) Equal to $\frac{P + Q \sin \theta}{mg - Q \cos \theta}$

(d) Equal to $\frac{P + Q \cos \theta}{mg - Q \sin \theta}$

Answer (a)



block will remain in equilibrium if

$$(f_s)_{\max} \geq P + Q \sin \theta$$

$$\mu(mg + Q \cos \theta) \geq P + Q \sin \theta$$

$$\mu \geq \frac{P + Q \sin \theta}{mg + Q \cos \theta}$$

22. Knowing that the acceleration due to gravity on the Earth surface is g and the radius of the Earth is R , a small body of mass m falls on the Earth from a height $h = \frac{R}{5}$ above the Earth's surface. During the freefall, the potential energy of the falling body decreases by


(a) mgh

(b) $\frac{4}{5} mgh$

(c) $\frac{5}{6} mgh$

(d) $\frac{6}{7} mgh$

Answer (c)



$$PE_i = \frac{-GMm}{R + \frac{R}{5}} = \frac{-5mgR}{6}$$

$$PE_f = \frac{-GMm}{R}$$

$$\Delta PE = \frac{mgR}{6} = \frac{5mg}{6} \cdot \left(\frac{R}{5}\right) = \frac{5mgh}{6}$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC


(Group A & B)

38
NSEP

34
NSEJS

23. At some instant, a motor car is moving on a circular path of radius 600 m, with a speed $u = 30 \text{ ms}^{-1}$. If its speed is increased at a rate of 2 ms^{-2} , the magnitude of the acceleration of the car at that instant is
- (a) 2.0 ms^{-2} (b) 2.5 ms^{-2}
 (c) 3.5 ms^{-2} (d) 1.5 ms^{-2}

Answer (b)



Sol.

$$a_t = 2 \text{ m/s}^2$$

$$a_c = \frac{u^2}{R} = \frac{30 \times 30}{600} = \frac{3}{2} \text{ m/s}^2$$

$$a = \sqrt{a_c^2 + a_t^2}$$

$$= \frac{5}{2} \text{ m/s}^2$$

24. A cricket ball, thrown across a field, is at heights of h_1 and h_2 above the point of projection, at time t_1 and time t_2 after the throw, respectively. It is then caught by the wicket keeper at the same height as that from which it was thrown. The time of flight (T) of the ball is

(a) $T = \frac{h_1 t_2^2 - h_2 t_1^2}{h_1 t_2 - h_2 t_1}$


(b) $T = \frac{h_1 t_2^2 + h_2 t_1^2}{h_2 t_1 + h_1 t_2}$

(c) $T = \frac{h_1 t_1^2 - h_2 t_2^2}{h_1 t_1 - h_2 t_2}$

(d) $T = \frac{h_1 t_1^2 + h_2 t_2^2}{h_1 t_1 + h_2 t_2}$

Answer (a)

Sol.



$$h_1 = u_y t_1 - \frac{1}{2} g t_1^2$$

$$h_2 = u_y t_2 - \frac{1}{2} g t_2^2$$

$$h_1 t_2^2 = u_y t_1 t_2^2 - \frac{1}{2} g t_1^2 t_2^2$$

1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

$$h_2 t_1^2 = u_y t_2 t_1^2 - \frac{1}{2} g t_1^2 t_2^2$$

$$h_1 t_2^2 - h_2 t_1^2 = u_y t_1 t_2 (t_2 - t_1) \quad \dots(1)$$

$$h_1 t_2 = u_y t_1 t_2 - \frac{1}{2} g t_1^2 t_2$$


$$h_2 t_1 = u_y t_1 t_2 - \frac{1}{2} g t_1 t_2^2$$

$$h_1 t_2 - h_2 t_1 = \frac{1}{2} g t_1 t_2 (t_2 - t_1) \quad \dots(2)$$

(1)/(2)

$$\frac{h_1 t_2^2 - h_2 t_1^2}{h_1 t_2 - h_2 t_1} = \frac{2u_y}{g} = T$$

25. A plate of mass M is placed on a horizontal frictionless surface S . A block of mass m is placed on the plate. The coefficient of dynamic friction between the block and the plate is μ . If a horizontal force $F = 2\mu mg$ is applied to the block (as shown), the acceleration of the plate will be



(a) $\frac{\mu mg}{M}$

(b) $\frac{\mu mg}{m + M}$

(c) $\frac{2\mu mg}{M}$

(d) $\frac{2\mu mg}{m + M}$

Answer (a or d)

Sol. If both moves together

$$a_c = \frac{2\mu mg}{M + m}$$

For plate $\mu mg > M \cdot \frac{2\mu mg}{m + M}$

$$m + M > 2m$$

$$m > M$$

i.e. If $M < m$, $a_{\text{plate}} = \frac{2\mu mg}{m + M}$

But If $M > m$, $a_{\text{plate}} = \frac{\mu mg}{M}$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49

NSEA

(Group A & B)

229

NSEB

(Group A & B)

70

NSEC

(Group A & B)

38

NSEP


34

NSEJS

26. A simple pendulum, with a bob of mass m , oscillates in a vertical plane, with an angular amplitude θ_0 . The tension in its string when it passes through the mean position is $2mg$. Neglecting the effect of air friction and the viscosity of air, the angular amplitude θ_0 is
- (a) 30° (b) 60°
 (c) 90° (d) 120°

Answer (b)

Sol.



$$T - mg = \frac{mv_0^2}{l}$$


$$v_0^2 = gl$$

$$\frac{1}{2}mv_0^2 = mgl(1 - \cos\theta_0)$$

$$1 - \cos\theta_0 = \frac{1}{2}$$

$$\theta_0 = 60^\circ$$

27. Because of their mutual gravitational attraction, four identical planets each of mass m are orbiting in a circular path of radius r in the same sense (angular direction). The magnitude of the velocity of each planet is



(a) $\left[\frac{Gm}{r} \left(\frac{1+2\sqrt{2}}{4} \right) \right]^{\frac{1}{2}}$

(b) $3\sqrt{\frac{Gm}{r}}$

(c) $\sqrt{\frac{Gm}{r}(1+2\sqrt{2})}$


(d) $\left[\frac{1}{2} \frac{Gm}{r} \left(\frac{1+\sqrt{2}}{2} \right) \right]^{\frac{1}{2}}$

Answer (a)

1484 Students Scored Above MAS

<div style="border: 1px solid black; border-radius: 50%; width: 60px; height: 60px; margin: 0 auto; display: flex; flex-direction: column; align-items: center; justify-content: center;"> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">420</div> <div style="font-size: 8px; font-weight: normal; margin-bottom: 5px;">Classroom Students</div> <div style="font-size: 8px; font-weight: normal;">Qualified in NSEs 2024-25</div> </div>	<div style="border: 1px solid black; border-radius: 50%; width: 60px; height: 60px; margin: 0 auto; display: flex; flex-direction: column; align-items: center; justify-content: center;"> <div style="font-size: 8px; font-weight: normal; margin-bottom: 5px;">(Group A & B)</div> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">49</div> <div style="font-size: 8px; font-weight: normal;">NSEA</div> </div>	<div style="border: 1px solid black; border-radius: 50%; width: 60px; height: 60px; margin: 0 auto; display: flex; flex-direction: column; align-items: center; justify-content: center;"> <div style="font-size: 8px; font-weight: normal; margin-bottom: 5px;">(Group A & B)</div> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">229</div> <div style="font-size: 8px; font-weight: normal;">NSEB</div> </div>	<div style="border: 1px solid black; border-radius: 50%; width: 60px; height: 60px; margin: 0 auto; display: flex; flex-direction: column; align-items: center; justify-content: center;"> <div style="font-size: 8px; font-weight: normal; margin-bottom: 5px;">(Group A & B)</div> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">70</div> <div style="font-size: 8px; font-weight: normal;">NSEC</div> </div>	<div style="border: 1px solid black; border-radius: 50%; width: 60px; height: 60px; margin: 0 auto; display: flex; flex-direction: column; align-items: center; justify-content: center;"> <div style="font-size: 8px; font-weight: normal; margin-bottom: 5px;">(Group A & B)</div> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">38</div> <div style="font-size: 8px; font-weight: normal;">NSEP</div> </div>	<div style="border: 1px solid black; border-radius: 50%; width: 60px; height: 60px; margin: 0 auto; display: flex; flex-direction: column; align-items: center; justify-content: center;"> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">34</div> <div style="font-size: 8px; font-weight: normal;">NSEJS</div> </div>
---	--	---	--	--	---

Sol.




$$F_1 + F_2 \cos 45 + F_3 \cos 45 = \frac{mv^2}{r}$$

$$\frac{Gm^2}{4r^2} + \frac{Gm^2}{2r^2} \cdot \frac{1}{\sqrt{2}} + \frac{Gm^2}{2r^2} \cdot \frac{1}{\sqrt{2}} = \frac{mv^2}{r}$$

$$\frac{Gm^2}{4r^2} (1 + 2\sqrt{2}) = \frac{mv^2}{r}$$

$$v = \sqrt{\frac{Gm}{r} \left(\frac{1 + 2\sqrt{2}}{4} \right)}$$

28. A rigid square sheet of size 2 m × 2 m is hinged at the middle of the vertical edges to serve as a door which can turn about the horizontal axis OO'. A fluid of density ρ fills the space to the left of the sheet up to its top. The horizontal force F required (to be applied at the lower edge) to hold the sheet vertical is



(a) $\frac{2}{3} \rho g$


(b) $\frac{4}{3} \rho g$

(c) $\frac{8}{3} \rho g$

(d) $\frac{1}{3} \rho g$

Answer (b)

Sol. Torque due to fluid on the sheet about OO'



$$\int_{-1}^1 \rho(1+x)g \cdot 2dx \cdot x$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

$$\rho g \cdot 2 \cdot \frac{2}{3} = \frac{4\rho g}{3}$$

$$F \times 1 = \frac{4\rho g}{3}$$

$$F = \frac{4\rho g}{3}$$

29. A major artery in human body, with radius 0.4 cm, carries blood at a flow rate of 5.0 cubic centimeters per second. The pressure difference of blood per meter length of the artery is nearly [Given that the coefficient of viscosity (η) of blood at body temperature is 4.0×10^{-3} Pa.s and the density of mercury is 13.6 g/cm^3]
- (a) 9.6 mm of Hg (b) 3.2 mm of Hg
 (c) 1.5 mm of Hg (d) 6.0 mm of Hg

Answer (c)

Sol. $\Delta P = \frac{8\eta LQ}{\pi R^4}$

$$= \frac{8 \times 4 \times 10^{-3} \times 1 \times 5 \times 10^{-6}}{3.14 \times (4 \times 10^{-3})^4} = 0.199 \times 10^3 \text{ Pa}$$

$$h = \frac{0.199 \times 10^3}{13.6 \times 10^3 \times 10} \times 1000 \text{ mm}$$

$$= 1.467 \text{ mm} \approx 1.5 \text{ mm of Hg}$$

30. If P represents radiation pressure, E represents radiation energy striking per unit area per unit time and c represents speed of light then the possible values of non-zero integers x , y and z such that $P^x E^y c^z$ is dimensionless, may be
- (a) $x = 1, y = 1, z = 1$ (b) $x = -1, y = 1, z = 1$
 (c) $x = 1, y = -1, z = 1$ (d) $x = 1, y = 1, z = -1$

Answer (c)

Sol. $[P] = \text{ML}^{-1}\text{T}^{-2}$
 $[E] = \text{MT}^{-3}$
 $[C] = \text{LT}^{-1}$
 $[P]^1 [E]^{-1} [C]^1 = \text{M}^0 \text{L}^0 \text{T}^0$

31. A large tank, open at the top, has two small holes in the vertical wall. One is a square hole of side ' s ' at a depth h below the top and the other is a circular hole of radius r at a depth $4h$ below the top (given that $s \ll h; r \ll h$). When the tank is completely filled up to the brim with water, the quantity of water flowing out per second from each hole is the same, then r is equal to
- (a) $2\pi s$ (b) $\frac{s}{2\pi}$
 (c) $\frac{s}{\sqrt{2\pi}}$ (d) $\frac{s}{2\sqrt{\pi}}$

Answer (c)

1484 Students Scored Above MAS

420

Classroom Students
 Qualified in
 NSEs 2024-25

(Group A & B)

49
 NSEA

(Group A & B)

229
 NSEB

(Group A & B)

70
 NSEC

(Group A & B)

38
 NSEP

34
 NSEJS


Sol. Rate of water flow = $AV = A\sqrt{2gh}$

$$S^2 \cdot \sqrt{2gh} = \pi r^2 \cdot \sqrt{2g4h}$$

$$S^2 = 2\pi r^2$$

$$r = \frac{S}{\sqrt{2\pi}}$$

32. A pendulum consists of a heavy but very small bob of mass M suspended at the end of a rigid rod of mass m and length L . The time period of small oscillations in the vertical plane, about a horizontal axis through the upper end of the rod is



(a) $2\pi\sqrt{\left(\frac{m+2M}{m+3M}\right) \times \left(\frac{3L}{2g}\right)}$


(b) $2\pi\sqrt{\left(\frac{m+3M}{m+2M}\right) \times \left(\frac{2L}{3g}\right)}$

(c) $2\pi\sqrt{\frac{3L}{2g}}$

(d) $2\pi\sqrt{\left(\frac{2m+M}{3m+M}\right) \times \left(\frac{3L}{2g}\right)}$

Answer (b)

Sol.



$$\tau = -\left(Mgl \sin\theta + Mg \cdot \frac{l}{2} \sin\theta\right)$$

$$\tau = -\left(M + \frac{m}{2}\right)gl \sin\theta$$

$$\tau = -\left(\frac{2M+m}{2}\right)gl\theta$$

$$I = Ml^2 + \frac{Ml^2}{3}$$

$$T = 2\pi\sqrt{\frac{(3M+m)\frac{l^2}{3}}{(2M+m)\frac{gl}{2}}}$$

$$= 2\pi\sqrt{\left(\frac{3M+m}{2M+m}\right) \cdot \left(\frac{2l}{3g}\right)}$$

420
Classroom Students
Qualified in
NSEs 2024-25

1484 Students Scored Above MAS

(Group A & B)
49
NSEA


(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

33. A transverse wave is travelling along a long stretched string from left to right (along +ve x direction). The snapshot of a small part of the string at any moment t is shown in the figure. At this particular instant



- (a) A and E are at rest for a moment while C and G have maximum speed
- (b) B and D have upward velocity whereas F and H have downward
- (c) D, E, F are moving downward at that moment
- (d) B and H are moving downward at that moment

Answer (d)


Sol. $v_{\text{particle}} = \left(-\frac{\partial y}{\partial x}\right)v_{\text{wave}}$

Where $\frac{\partial y}{\partial x}$ = slope

\therefore Here only B and H are having positive slope.

\therefore They move downward.

34. Two tuning forks, with natural frequency 700 Hz each, move relative to a stationary observer. Fork (1) moves towards the observer while the fork (2) moves away from the observer. Both the forks move with same velocity v on the same line. The observer, standing between the two forks, hears 4 beats per sec. Using the speed of sound in air as $v_s = 350 \text{ ms}^{-1}$, the speed of each tuning fork is



- (a) 2.0 ms^{-1}
- (b) 1.5 ms^{-1}
- (c) 1.0 ms^{-1}
- (d) 0.5 ms^{-1}

Answer (c)

1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

Answer (d)

Sol. $m_1 = 1$

$$\gamma_1 = \frac{5}{3}$$


$m_2 = 2$

$$\gamma_2 = \frac{7}{5}$$

$$\gamma_{\text{mixture}} = \frac{m_1\gamma_1 + m_2\gamma_2}{m_1 + m_2} = \frac{1 \times \frac{5}{3} + 2 \times \frac{7}{5}}{3} = 1.48$$

$$V_s = \sqrt{\frac{\gamma RT}{m}} = \sqrt{\frac{1.48 \times 8.31 \times 300 \times 1000 \times 3}{68}} \approx 401 \text{ m/s}$$

37. A thin uniform circular disc of radius 'a' is placed in XY plane with its center at origin (0, 0). A small circular disc of radius b with center at (c, 0) is cut and taken out to create a hole. The center of mass of the remaining disc is at



(a) $-\frac{b^2}{a^2}c, 0$

(b) $-\frac{b^2}{a^2 - c^2}c, 0$


(c) $-\frac{b^2}{a^2 + b^2}c, 0$

(d) $-\frac{b^2}{a^2 - b^2}c, 0$


Answer (d)

Sol. Let mass per unit area be σ

\therefore Mass of original complete disc before removal of small disc = $\sigma\pi a^2$



Let COM of remaining disc be at $(-x, 0)$



1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

$$m_1 = \sigma\pi(a^2 - b^2)$$

$$m_2 = \sigma\pi b^2$$

\therefore COM of complete disc would lie at (0, 0)

$$m_1x = m_2c$$

$$\Rightarrow \sigma\pi(a^2 - b^2)x = \sigma\pi b^2c$$

$$\Rightarrow x = \frac{cb^2}{a^2 - b^2}$$

$$\therefore \text{Answer} \left(-\frac{b^2c}{a^2 - b^2}, 0 \right)$$

38. One mole of an ideal monoatomic gas, contained in a cylinder fitted with movable piston, is originally at P_1 , V_1 and $T_1 = 27^\circ\text{C}$. The gas is slowly heated. Initially 8.31 watt-hour of energy is added to it; at the same time it is allowed to expand at constant pressure to a new state P_1 , V_2 and T_2 .

The correct option is

- (a) Value of T_2 is 1740°C
- (b) Work done by the gas is 2160 R joule
- (c) Internal energy of the gas increases by 1440 R joule
- (d) $\frac{V_2}{V_1} = 5.8$

Answer (d)

Sol. $n = 1$

$$T_1 = 27^\circ\text{C} \text{ or } 300\text{ K}$$

$$\text{Energy given to gas is } \Delta Q = 8.31 \times 3600\text{ J} = 29916\text{ J}$$

According to first law of thermodynamics

$$\Delta Q = \Delta U + w$$

$$\Rightarrow 29916 = nC_v\Delta T + P_1\Delta V$$

$$\Rightarrow 29916 = n \frac{3}{2} R\Delta T + nR\Delta T$$

$$\Rightarrow 29916 = \frac{5}{2} R(T_2 - T_1)$$

$$\Rightarrow T_2 = 1740\text{ K}$$

$$w = P_1\Delta V = nR\Delta T$$

$$\Rightarrow w = 1440R$$

$$\Delta U = nC_v\Delta T = \frac{3}{2} R \times 1440$$

$$\Rightarrow \Delta U = 2160R$$

$$\text{and } \frac{V_1}{V_2} = \frac{T_1}{T_2} = \frac{300}{1740}$$

$$\therefore \frac{V_2}{V_1} = \frac{174}{30} = 5.8$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)


70
NSEC

(Group A & B)

38
NSEP


34
NSEJS

39. A non-conducting solid sphere, of radius R with its center at A , has a spherical cavity of diameter R with center at B as shown. There is no charge in the cavity while the solid part has a uniform volume charge density ρ .
 Electric potential at the center of the sphere (at point A) is $V = \frac{k\rho R^2}{12\epsilon_0}$ (in SI units) where the value of k is



- (a) 3
 (c) 7
 (b) 5
 (d) 9

Answer (b)




Potential at centre of complete sphere is $\frac{\rho R^2}{2\epsilon_0}$

As a part of sphere is removed, potential due to it will be subtracted.

$$\therefore V_A = \frac{\rho R^2}{2\epsilon_0} - \frac{\rho \left(\frac{R}{2}\right)^2}{3\epsilon_0} = \frac{5\rho R^2}{12\epsilon_0}$$

$$\therefore k = 5$$

40. The figure below depicts the voltage wave forms of binary input signals A and B and the output signal C of a certain logic gate.



The logic gate is

- (a) AND
 (c) OR
 (b) NAND
 (d) XOR

Answer (a)

Sol. From the diagram shown in figure it is clear that if both inputs are high only then output is high.

\therefore AND gate

1484 Students Scored Above MAS

<div style="border: 2px solid black; border-radius: 50%; width: 60px; height: 60px; display: flex; flex-direction: column; justify-content: center; align-items: center; margin: 0 auto;"> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">420</div> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">Classroom Students</div> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">Qualified in</div> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">NSEs 2024-25</div> </div>	<div style="border: 2px solid black; border-radius: 50%; width: 60px; height: 60px; display: flex; flex-direction: column; justify-content: center; align-items: center; margin: 0 auto;"> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">(Group A & B)</div> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">49</div> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">NSEA</div> </div>	<div style="border: 2px solid black; border-radius: 50%; width: 60px; height: 60px; display: flex; flex-direction: column; justify-content: center; align-items: center; margin: 0 auto;"> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">(Group A & B)</div> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">229</div> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">NSEB</div> </div>	<div style="border: 2px solid black; border-radius: 50%; width: 60px; height: 60px; display: flex; flex-direction: column; justify-content: center; align-items: center; margin: 0 auto;"> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">(Group A & B)</div> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">70</div> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">NSEC</div> </div>	<div style="border: 2px solid black; border-radius: 50%; width: 60px; height: 60px; display: flex; flex-direction: column; justify-content: center; align-items: center; margin: 0 auto;"> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">(Group A & B)</div> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">38</div> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">NSEP</div> </div>	<div style="border: 2px solid black; border-radius: 50%; width: 60px; height: 60px; display: flex; flex-direction: column; justify-content: center; align-items: center; margin: 0 auto;"> <div style="font-size: 24px; font-weight: bold; margin-bottom: 5px;">34</div> <div style="font-size: 8px; font-weight: bold; margin-bottom: 5px;">NSEJS</div> </div>
---	--	---	--	--	---

41. Energy from the Sun falls on the Earth surface at the rate of 1400 W/m^2 , which is known as solar constant. The respective rms values E_{rms} and B_{rms} of electric and magnetic fields in the sunlight (electromagnetic radiation) reaching Earth surface are (Take speed of light $c = 3 \times 10^8 \text{ ms}^{-1}$)
- (a) $E_{\text{rms}} = 726.5 \text{ V/m}$, $B_{\text{rms}} = 2.42 \mu\text{T}$ (b) $E_{\text{rms}} = 7260 \text{ V/m}$, $B_{\text{rms}} = 242 \text{ nT}$
 (c) $E_{\text{rms}} = 1030 \text{ V/m}$, $B_{\text{rms}} = 3.42 \mu\text{T}$ (d) $E_{\text{rms}} = 10300 \text{ V/m}$, $B_{\text{rms}} = 342 \text{ nT}$

Answer (a)

Sol. $I = 1400 \text{ W/m}^2$

4 we know that $I = \frac{1}{2} C \epsilon_0 E_0^2$

$\therefore \frac{1}{2} C \epsilon_0 E_0^2 = 1400$


$\Rightarrow E_0 \sqrt{\frac{2800}{C \epsilon_0}} = 1027.41$

$E_{\text{rms}} = \frac{E_0}{\sqrt{2}} = 726.5 \text{ V/m}$

$\therefore B_{\text{rms}} = 2.42 \mu\text{T}$

42. Two electric charges, $+q$ at the origin $O(0, 0)$ and $-2q$ at the point $A(6, 0)$ are placed on x axis. The locus of the point P in x - y plane where the potential vanishes ($V = 0$) is
- (a) a straight line perpendicular to x axis and passing through $(2, 0)$
 (b) only the point $(2, 0)$
 (c) a circle with center at $(-2, 0)$ and radius 4
 (d) an ellipse with foci at O and A

Answer (c)



If we need to find locus of points with O electric potential then Electric potential due to both charges should be equal and opposite let P be such a point.

$\frac{Kq}{r_1} = \frac{K_2q}{r_2}$

$\Rightarrow \frac{1}{\sqrt{x^2 + y^2}} = \frac{2}{\sqrt{(6-x)^2 + y^2}}$

$\Rightarrow 4(x^2 + y^2) = (6-x)^2 + y^2$

$\Rightarrow 4x^2 + 4y^2 = 36 + x^2 - 12x + y^2$

$\Rightarrow 3x^2 + 3y^2 + 12x - 36 = 0$

1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP


34
NSEJS

$$\Rightarrow x^2 + y^2 + 4x - 12 = 0$$

$$\Rightarrow (x + 2)^2 + y^2 = (4)^2$$

\therefore locus is a circle with center at $(-2, 0)$ and radius 4.

43. In the circuit shown, the Zener diode is an ideal one with breakdown voltage of 5.0 volt. The values of the resistances are $R_S = 10\text{ k}\Omega$ and $R_L = 1\text{ k}\Omega$. The current through the resistances, when the supply voltage is 11.0 V, is




- (a) 0.6 mA through R_S and 5.0 mA through R_L (b) 1.0 mA through R_S and 1.0 mA through R_L
 (c) 1.1 mA through R_S and no current through R_L (d) no current through R_S and 11 mA through R_L

Answer (b)

Sol. $V_z = 5\text{V}$

$$R_S = 10\text{ k}\Omega \quad R_L = 1\text{ k}\Omega$$




\therefore Zener diode is in reverse bias & if no current passes through it, voltage across Zener diode would be

$$\frac{11}{R_S + R_L} \times R_L = 1\text{V}$$

\therefore No current through Zener diode will loss.

$$i = \frac{11}{10000 + 1000} = 1\text{ mA through both resistances.}$$

44. In an accelerator the electrons are accelerated up to an energy of 50 MeV. The electrons do not emerge continuously from the accelerator rather they come in pulses at time interval of 5.0 milliseconds. Each pulse has a much shorter duration of 200 nanoseconds. Electron current during the pulse is 100 mA, while the current is zero between the two successive pulses (see figure), then



1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

- (a) the average current per pulse is 4 mA
- (b) the peak value of power delivered by the electron beam is 50 MW
- (c) the average power delivered by the electron beam is 200 W
- (d) the average power delivered by the electron beam is 2 MW

Answer (c)

Sol. Current per pulse is given as 100 mA and time duration of pulse is 200 ms

$$\therefore i = \frac{q}{t} \Rightarrow q = it$$

$$\Rightarrow q = 100 \times 10^{-3} \times 200 \times 10^{-9}$$

$$\Rightarrow q = 2 \times 10^{-8} \text{ C} = me$$

$$n = \frac{2 \times 10^{-8}}{1.6 \times 10^{-19}} = 1.25 \times 10^{11} \text{ electrons pass during this period.}$$


Each electron has energy 50 MeV

$$\therefore \text{Peak power would be } \frac{50 \times 10^6 \times 1.6 \times 10^{-19} \times 2 \times 10^{-8}}{200 \times 10^{-9} \times 1.6 \times 10^{-9}}$$

$$\Rightarrow 5 \text{ Mw}$$

$$\text{Average power} = \frac{5 \times 10^6 \times 200 \times 10^{-9}}{5 \times 10^{-3}} = 200 \text{ W}$$

45. Two infinitely long straight parallel wires perpendicular to the plane of the paper are 5 m apart. One of the wires, *P* carries current *I* out of the plane of the paper and the other, *Q* carries the current *I* into the plane of paper. The magnetic field *B* at the origin *O* of the coordinate system with *x* and *y* axes as perpendicular and parallel to *PQ*, respectively, is [Given *OP* = 4 m and *OQ* = 3 m]



(a) $\frac{\mu_0 I}{2\pi} \left(\hat{i} - \frac{3}{5} \hat{j} \right)$

(b) $\frac{\mu_0 I}{5\pi} \left(\hat{i} - \frac{7}{24} \hat{j} \right)$

(c) $\frac{\mu_0 I}{5\pi} \left(-\hat{i} + \frac{3}{8} \hat{j} \right)$

(d) $\frac{\mu_0 I}{24\pi} (2\hat{i} + 3\hat{j})$

Answer (b)

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB


(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS



$$B_Q = \frac{\mu_0 I}{2\pi \times 3}$$

$$\text{or } \vec{B}_Q = \frac{\mu_0 I}{2\pi \times 3} (\cos 53^\circ \hat{i} - \sin 53^\circ \hat{j})$$

$$\Rightarrow \vec{B}_Q = \frac{\mu_0 I}{6\pi} \left(\frac{3}{5} \hat{i} - \frac{4}{5} \hat{j} \right) = \frac{\mu_0 I}{30\pi} (3\hat{i} - 4\hat{j})$$

$$\text{Similarly, } \vec{B}_P = \frac{\mu_0 I}{2\pi \times 4} \left(\frac{4}{5} \hat{i} + \frac{3}{5} \hat{j} \right) = \frac{\mu_0 I}{40\pi} (4\hat{i} + 3\hat{j})$$

$$\vec{B}_O = \vec{B}_P + \vec{B}_Q = \frac{\mu_0 I}{10\pi} \left(\hat{i} + \frac{3}{4} \hat{j} + \hat{i} - \frac{4}{3} \hat{j} \right)$$

$$\Rightarrow \frac{\mu_0 I}{10\pi} \left(2\hat{i} - \frac{7}{12} \hat{j} \right)$$

$$\Rightarrow \frac{\mu_0 I}{5\pi} \left(\hat{i} - \frac{7}{24} \hat{j} \right)$$

46. Charge q is uniformly distributed over the surface of a thin non-conducting annular disc of inner radius R_1 and outer radius R_2 . The disc is made to rotate with constant frequency f , about an axis passing through the center of the annular disc and perpendicular to its plane. The magnetic moment of the disc is

(a) $\pi f q \frac{R_2^2 + R_1^2}{2}$

(b) $\pi f q \frac{R_2^2 - R_1^2}{2}$

(c) $\pi f q \frac{R_2^2 - R_1^2}{4}$

(d) $2\pi f q (R_2^2 - R_1^2)$

Answer (a)

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB


(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS



$$\omega = 2\pi f$$

$$\sigma = \frac{q}{\pi(R_2^2 - R_1^2)}$$

Small magnetic moment due to element shown in figure $dM = diA$

$$\Rightarrow dM = \frac{dq}{T} A$$

$$\Rightarrow dM = \sigma 2\pi r dr f \times \pi r^2$$

$$\Rightarrow dM = \sigma 2\pi^2 r^3 dr$$


$$\therefore M = \int dM = \int \sigma 2\pi^2 r^3 dr$$

$$\Rightarrow M = \frac{\sigma 2\pi^2 f}{4} (R_2^4 - R_1^4)$$

$$\Rightarrow M = \frac{qf\pi^2}{2\pi(R_2^2 - R_1^2)} (R_2^2 + R_1^2)(R_2^2 - R_1^2)$$

$$M = \frac{fq\pi(R_2^2 + R_1^2)}{2}$$

47. For a resistance R and capacitance C in series, the impedance is twice that of a parallel combination of the same elements when used with an AC voltage of frequency f . The frequency f of the applied emf is



(a) $f = 2\pi RC$

(b) $f = \frac{1}{2\pi RC}$

(c) $f = \frac{2\pi}{RC}$

(d) $f = \frac{1}{2\pi\sqrt{R^2 + C^2}}$

Answer (b)

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

Sol. $X_C = -\frac{j}{\omega C}$

For series combination

$$Z_1 = R - \frac{j}{\omega C}$$

$$|Z_1| = \sqrt{R^2 + \frac{1}{(\omega C)^2}}$$

For parallel combination

$$Z_2 = \frac{R \times \frac{-j}{\omega C}}{R - \frac{j}{\omega C}}$$

$$|Z_2| = \frac{1}{\sqrt{R^2 + (\omega C)^2}}$$


$$|Z_1| = 2|Z_2|$$

$$\Rightarrow \sqrt{R^2 + \left(\frac{1}{\omega C}\right)^2} = 2 \sqrt{\frac{1}{\left(\frac{1}{R}\right)^2 + (\omega C)^2}}$$

$$\Rightarrow R^2 + \omega^2 C^2 R^4 + \frac{1}{\omega^2 C^2} + R^2 = 4 R^2$$

$$\Rightarrow f = \frac{1}{2\pi RC}$$

48. In a certain electrical network, the three nodes A, B and C are each at a potential of 1.0 volt while the node D is at a potential 2.0 volt. The potential at the Node O in volt is



1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

- (a) $\frac{3}{2}$
(c) $\frac{5}{4}$

- (b) $\frac{4}{3}$
(d) $\frac{6}{5}$

Answer (c)

Sol. Let node O is at x V.

Using KCL

$$\frac{1-x}{10} + \frac{2-x}{10} + \frac{1-x}{10} + \frac{1-x}{10} = 0$$


$$\Rightarrow \boxed{x = \frac{5}{4} \text{ V}}$$

A - 2

ANY NUMBER OF OPTIONS (4, 3, 2 or 1) MAY BE CORRECT

MARKS WILL BE AWARDED ONLY IF ALL THE CORRECT OPTIONS ARE BUBBLED AND NO INCORRECT


49. Two ideal inductors $L_1 = L_2 = L$ and three identical resistors $R_1 = R_2 = R_3 = R$ have been connected to a DC source of emf E as shown in the circuit. When the key K is kept pressed (closed) for a long time, the current through the resistance R_1 on the extreme right is measured to be I . Immediately after releasing (switching off) the key, the current through the resistors is



- (a) I downwards in R_1
(b) I downwards in R_2
(c) $2I$ upwards in R_3
(d) Zero in each R_1, R_2 and R_3


Answer (a, b, c)

Sol. In steady state (Just before reopening switch)



Now current through inductor doesn't change instantaneously but in resistor it can.

After opening switch, using KCL



1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

50. A particle of mass m moves along x -axis with its potential energy as $U(x) = \frac{\alpha}{x^2} - \frac{\beta}{x}$ where α and β are positive constants. The particle is released from rest at $x_0 = \frac{\alpha}{\beta}$. Then

(a) $U(x)$ can be expressed as $U(x) = \frac{\alpha}{x_0^2} \left[\left(\frac{x_0}{x} \right)^2 - \frac{x_0}{x} \right]$

(b) Velocity of the particle $v(x)$ as a function of x can be expressed as $v(x) = \left[\frac{2\alpha}{mx_0^2} \left\{ \frac{x_0}{x} - \left(\frac{x_0}{x} \right)^2 \right\} \right]^{\frac{1}{2}}$

(c) The maximum speed of the particle is $v_{\max} = \sqrt{\frac{\alpha}{2mx_0^2}}$

(d) The total energy of the particle $KE(x) + U(x)$ is zero.

Answer (a, b, c, d)

Sol. $U(x) = \frac{\alpha}{x^2} - \frac{\beta}{x} = \left\{ \frac{\alpha x_0^2}{x^2} - \frac{\beta x_0^2}{x} \right\} \frac{1}{x_0^2}$

$$= \frac{\alpha}{x_0^2} \left\{ \frac{x_0^2}{x^2} - \frac{\beta}{\alpha} \frac{x_0^2}{x} \right\}$$

Here $x_0 = \frac{\alpha}{\beta}$

$$\Rightarrow U = \frac{\alpha}{x_0^2} \left(\left(\frac{x_0}{x} \right)^2 - \frac{x_0}{x} \right)$$

\Rightarrow (a)

Conservative system $\Rightarrow M = \text{constant}$, M at $x_0 = M$ at x .

$$\frac{1}{2} m(0)^2 + U(x_0) = \frac{\alpha}{x_0^2} \left(\left(\frac{x_0}{x} \right)^2 - \frac{x_0}{x} \right) + \frac{1}{2} mv^2$$

$$\Rightarrow \frac{1}{2} mv^2 = \frac{\alpha}{x_0^2} \left(\frac{x_0}{x} - \left(\frac{x_0}{x} \right)^2 \right)$$

\Rightarrow (b)

As M is constant, $M = k + U$, then k is max where U is min.

$$\frac{dv}{dx} = 0 \Rightarrow 2 \left\{ \frac{x_0}{x} \right\} \frac{x_0}{x^2} = \frac{x_0}{x^2} \quad (x \neq 0)$$

$$x = 2x_0$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS


$$\Rightarrow 0 = U(2x_0) + \frac{1}{2}mv_{\max}^2$$

$$\frac{1}{2}mv_{\max}^2 = \frac{\alpha}{x_0^2} \left\{ \frac{1}{2} - \frac{1}{4} \right\}$$

$$v_{\max}^2 = \frac{\alpha}{2mx_0^2}$$

\Rightarrow (c) also (d)

51. Two blocks A and B, of masses M and $2M$, respectively, are connected by a massless spring of natural length L_0 and spring constant K . The blocks are initially at rest on a smooth horizontal floor with spring at its natural length L_0 . A third block C of mass M , identical to that of block A, moves on the floor with speed v along the line joining A and B and collides with A elastically. In the subsequent motion.



- (a) The spring will be compressed to a maximum when at a length of $v\sqrt{\frac{M}{3K}}$
- (b) The kinetic energy of A and B together, when the spring is compressed to the maximum, is $\frac{Mv^2}{6}$
- (c) The blocks A and B stop for a moment when the spring is at the maximum compression
- (d) The time required to reach the maximum compression from the normal length is $\frac{\pi}{2}\sqrt{\frac{2M}{3K}}$

Answer (b, d)

Sol. In elastic collision between identical masses, velocity interchanges after collision.



\Rightarrow At maximum compression, $v_{\text{rel}} = 0$

$$\frac{1}{2}\mu v_0^2 + \frac{1}{2}(3M)v_{\text{cm}}^2 + \frac{1}{2}K(0)^2 = \frac{1}{2}\mu(0)^2 + \frac{1}{2}mv_{\text{cm}}^2 + \frac{1}{2}kx^2$$

$$x = v_0\sqrt{\frac{\mu}{k}} = v_0\sqrt{\frac{2M}{3k}}$$

$\Rightarrow M = \text{constant}$, KE at maximum compression $KE = M - U_{\max}$

$$= \frac{1}{2}mv_0^2 - \frac{1}{2}kx^2 = \frac{2M}{3k}$$

$$= \frac{1}{6}mv_0^2 \Rightarrow b$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

Option (c) violates law conservation of momentum

In frame of centre of mass

$$A \frac{2\vec{v}_0}{3} \cdot \frac{\vec{v}_0}{3} B \xrightarrow{t} A \cdot \dot{C} \cdot \dot{O} \cdot M \cdot \dot{B}$$


That is quarter oscillation of reduced mass system

$$t = \frac{1}{4} 2\pi \sqrt{\frac{2M/3}{K}} = \frac{\pi}{2} \sqrt{\frac{2M}{3K}}$$

52. A thin uniform metallic rod, of length $l = 1.0$ m and area of cross section $A = 2 \text{ mm}^2$, is made to rotate with angular velocity $\omega = 400$ rad/s in a horizontal plane about a vertical axis through one of its ends. The density and the Young's modulus of the material of the rod are $\rho = 10^4 \text{ kg m}^{-3}$ and $Y = 2.0 \times 10^{11} \text{ Nm}^{-2}$. Taking r as the distance of a point on the rod from the axis of rotation, the

- (a) Tension at midpoint of the rod is $T = 1200$ N.
- (b) Tension in the rod varies with distance r from the axis of rotation as $T = 1600 r^2$ N
- (c) Stress in the rod at $r = 0.5$ m is $3.0 \times 10^8 \text{ Nm}^{-2}$
- (d) elongation of the rod is $\frac{8}{3}$ mm

Answer (a, d)



Sol.

$$T(r) = (ma_{\text{COM}})_{r \text{ to } R} \quad a_{\text{COM}} = \omega^2 \left(\frac{L+r}{2} \right)$$

$$= \left\{ \frac{\rho AL}{L} \right\} (L-r) \left(\frac{L+r}{2} \right) \omega^2$$

$$T(r) = \frac{\rho A}{2} (L^2 - r^2) \omega^2$$

$$= \frac{10^4 \times 2 \times 10^{-6}}{2} \times 4 \times 4 \times 10^4$$

$$= 12 \times 10^2 = 1200 \text{ N} \quad \Rightarrow \text{(a)}$$

Tension can't be same at both end points

$$\sigma = \frac{T}{A} = \frac{12 \times 10^6}{2 \times 10^{-6}} = 6 \times 10^{12}$$

1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS


$$\Delta l = \frac{F\ell}{YA} \Rightarrow \Delta l = \int \frac{\rho A \omega^2}{Y2A} (L^2 - r^2) dr$$

$$= \frac{\rho \omega^2}{2Y} \left\{ L^3 - \frac{L^3}{3} \right\}$$

$$= \frac{10^4 \times 16 \times 10^4}{2 \times 2 \times 10^{11}} \times \frac{2}{3}$$

$$= \frac{8}{3} \times 10^{-3} \text{ m}$$

53. One end of a long and thin rope, stretched horizontally with a tension $T = 8 \text{ N}$, along x axis, is supporting a weight after passing over a pulley fixed on a vertical pole (see figure). At the other end, a simple harmonic oscillator (a clamped iron rod along the axis of a solenoid fed with AC voltage and oscillating between north and south poles) at $x = 0$, generates a transverse wave of frequency 100 Hz and an amplitude of 2 cm , in the rope. The wave propagates along the rope. The mass per unit length of the rope is 20 g/m . Ignoring the effect of gravity (on the rope), the correct option(s) is/are



- (a) Wavelength of the transverse wave is 20 cm
 (b) Maximum magnitude of transverse acceleration of any point on the rope is nearly 800 ms^{-2}
 (c) If the oscillator produces maximum negative displacement at $x = 0$ at time $t = 0$, the equation of the wave can be expressed as $y(x, t) = -0.02 \sin[10\pi x - 100\pi t]$ in SI units.
 (d) Tension in the given rope remaining unchanged, if a harmonic oscillator of frequency 200 Hz is used (instead of earlier frequency 100 Hz), the wavelength will be 10 cm .

Answer (a, d)

Sol. $\mu = 20 \text{ g/m}$ $A = 2 \text{ cm}$ $f = 100 \text{ Hz}$

$$v = \sqrt{\frac{8 \times 1000}{20}} = 20 \text{ m/s}$$

(a) $\lambda = \frac{20}{100} = 20 \text{ cm}$

(b) $a_{\text{max}} = A\omega^2$
 $= 2 \times 10^{-2} \cdot 4\pi^2 \cdot 100 \times 100$
 $= 7887 \text{ m/s}^2$

(c) $\omega = 2\pi \times 100 = 200\pi$

(d) $v = f\lambda = 200 \times 0.1 = 20 \text{ m/s}$

V same T same

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

54. Nuclei of a radioactive element A are being produced at a constant rate α . The element A has a decay constant λ . If there are N_0 nuclei at $t = 0$, then

(a) Number of nuclei $N(t)$, at time t , is $N(t) = \frac{1}{\lambda}(\alpha - \lambda N_0)e^{-\lambda t}$

(b) If $\alpha = \lambda N_0$, the number of nuclei $N(t)$ at any time t will remain constant

(c) If $\alpha = 2\lambda N_0$ then $N(t) = 2N_0$ as $t \rightarrow \infty$

(d) If $\alpha = 2\lambda N_0$, the number of nuclei $N(t)$ after one half-life of A is $N\left(\frac{T}{2}\right) = \frac{3}{2}N_0$

Answer (b, c, d)

Sol. $\xrightarrow{\alpha} A \xrightarrow{\lambda}$

$$\frac{dN}{dt} = \alpha - \lambda N$$

$$\int_{N_0}^N \frac{dN}{\alpha - \lambda N} = \int_{t=0}^t dt$$

$$\alpha - \lambda N = (\alpha - \lambda N_0)e^{-\lambda t}$$

$$N = \frac{\alpha}{\lambda} - \left(\frac{\alpha}{\lambda} - N_0\right)e^{-\lambda t}$$

If $\alpha = \lambda N_0$

$$N(t) = \frac{\alpha}{\lambda} \text{ constant}$$

$$\alpha = 2\lambda N_0$$

$$N(t) = 2N_0 \text{ as } t \rightarrow \infty$$

$$N = 2N_0 - (N_0)e^{-\lambda \cdot \frac{\ln 2}{\lambda}}$$

$$= \frac{3N_0}{2}$$

55. In Young's double slit experiment, a fine beam of coherent monochromatic light of wavelength $\lambda = 600$ nm is incident on identical slits S_1 and S_2 at separation d . The intensity at the central maximum formed at O is I_{\max} and the angular fringe width is $\beta = 0.1^\circ$. When a thin transparent film is placed in front of the slit S_2 , the intensity at O changes. It is found that the smallest thickness of the film, for which the intensity at O becomes half the maximum intensity (i.e. $\frac{I_{\max}}{2}$), is 250 nm. Neglecting the absorption of light by the film, the zero order fringe earlier at O now forms at O' where $OO' = 0.5$ mm.

420

Classroom Students
Qualified in
NSEs 2024-25

1484 Students Scored Above MAS

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)


70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

Choose correct option(s)



- (a) The refractive index of the film is 1.6
- (b) The fringe width near O is 2 mm
- (c) On the screen, O' is above O
- (d) The distance D of the screen from the double slit is nearly 1.15 m

Answer (a, b, d)

Sol. $I = I_{\max} \cos^2\left(\frac{\Delta d}{2}\right) = \frac{I_{\max}}{2}$

$$\Delta d = \frac{\pi}{2}$$

$$\Delta x = \frac{\lambda}{4}$$

$$(\mu - 1)t = \frac{\lambda}{4}$$

$$(\mu - 1) = \frac{600}{4 \times 250}$$

$$\mu = 1.6$$

$$\frac{\lambda}{d} = 0.1^\circ = \frac{\pi}{1800} \text{ rad}$$

$$d = \frac{600 \times 1800}{3.14} \text{ nm} = 343949 \text{ nm}$$

Effective path difference at O' is zero Now

$$d \cdot \frac{y}{D} = (\mu - 1)t$$

$$D = \frac{343949 \text{ nm} \times 0.5 \text{ mm}}{150 \text{ nm}}$$

$$= 2992 \times 0.5 \text{ mm}$$

$$D \approx 1.15 \text{ m}$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)


70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

56. An insulated non-conducting solid sphere of radius 'a', carrying a positive charge +4Q uniformly distributed throughout its volume, is surrounded by a concentric thick conducting spherical shell of inner radius b and outer radius c. This thick shell carries a negative charge -2Q (see figure). The correct option(s) is/are




- (a) Electric field strength at distance $r (r < a)$ from the center is $\vec{E} = \frac{1}{4\pi\epsilon} \frac{4Q}{a^3} \vec{r}$
- (b) Charge on the inner surface of the conducting spherical shell is + 2Q
- (c) Charge on the outer surface of the conducting spherical shell is + 2Q
- (d) Electrical energy stored in region $0 < r < a$ [i.e. in the inner sphere] is $\frac{2Q^2}{5\pi\epsilon_0 a}$

Answer (a, c, d)

Sol. At $r < a$ field is due to solid sphere only

$$\vec{E} = \frac{k \cdot 4Q}{a^3} \vec{r} = \frac{1}{4\pi\epsilon_0} \cdot \frac{4Q}{a^3} \cdot \vec{r}$$



$$\frac{dE}{dV} = \frac{1}{2} \epsilon_0 \cdot \left(\frac{Qr}{\pi\epsilon_0 a^3} \right)^2$$

$$E = \int_0^a \frac{1}{2} \epsilon_0 \cdot \frac{Q^2 r^2}{\pi^2 \epsilon_0^2 a^6} \cdot 4\pi r^2 dr$$

$$= \frac{2}{5} \frac{Q^2}{\pi \epsilon_0 a}$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

57. A single electron orbits around a stationary nucleus of charge $+Ze$ in a hydrogen-like atom, where Z is the atomic number and e is the magnitude of the charge on an electron. It requires 47.25 eV to excite the electron from second Bohr orbit to the third Bohr orbit. Ionization energy of hydrogen atom is 13.6 eV. Then
- the value of Z is 5
 - the energy required to excite the electron from the 3rd orbit to the 4th orbit is 16.53 eV (nearly)
 - the wavelength of electromagnetic radiation required to liberate the electron completely when in the first Bohr orbit is 36.56 Å
 - the angular momentum of an electron in the second Bohr orbit is 1.056×10^{-33} Js

Answer (a, b, c)

Sol. $E_3 - E_2 = 47.25$

$$\Rightarrow \frac{-13.6z^2}{9} + \frac{13.6z^2}{4} = 47.25$$

$$\Rightarrow \frac{13.6z^2 \times 5}{36} = 47.25$$

$$z = 5$$

$$\Delta E = E_4 - E_3 = 13.6z^2 \left(\frac{1}{9} - \frac{1}{16} \right)$$

$$\Rightarrow \frac{13.6 \times 25 \times 7}{9 \times 16}$$


$$\Rightarrow 16.53 \text{ eV}$$

$$\frac{hc}{\lambda} = \frac{13.6 \times z^2}{1}$$

$$\Rightarrow \lambda = \frac{hc}{13.6 \times 25} = \frac{1240}{13.6 \times 25} = 36.5 \text{ Å}$$

$$mvr = \frac{mh}{2\pi} = \frac{2 \times h}{2\pi} = 2.11 \times 10^{-34} \text{ Js}$$

58. One mole of an ideal monoatomic gas of molecular mass M undergoes a cyclic process (ABCA) shown in the figure as a density (ρ) versus pressure (P) curve. The correct option(s) is/are



1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)


38
NSEP

34
NSEJS

- (a) Work done on the gas in going from A to B is $W_{AB} = \frac{MP_0}{\rho_0} \ln 2$
- (b) Work done by the gas in the process BC is $W_{BC} = \frac{MP_0}{2\rho_0}$
- (c) Efficiency (η) of the complete cycle ABCA is $\eta = \frac{2}{5} (1 - \ln 2)$
- (d) Heat rejected by the gas in the complete cycle ABCA is $Q_{ABCA} = \frac{MP_0}{\rho_0} (1 - \ln 2)$

Answer (a, c)

Sol. $n = 1$




First, we will convert this curve into PV diagram

$A \rightarrow B \Rightarrow P \propto \rho \therefore T = \text{Constant}$

$\therefore A \text{ to } B \text{ is isothermal.}$

$B \text{ to } C \text{ is isobaric}$

and $C \text{ to } A \text{ isochoric.}$



$$W_{A \text{ to } B} = - \int P dV = mRT \log_e \frac{V_2}{V_1}$$

$$\Rightarrow RT_0 \log_e 2$$

$$\Rightarrow \frac{MP_0}{\rho_0} \log_e 2$$

1484 Students Scored Above MAS

420
Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)
49
NSEA

(Group A & B)
229
NSEB

(Group A & B)
70
NSEC

(Group A & B)
38
NSEP

34
NSEJS

$$W_{B \text{ to } C} = 2P_0 \frac{V_0}{2} = \frac{MP_0}{\rho_0}$$

$$\eta = \frac{W_{\text{net}}}{Q} = \frac{\frac{MP_0}{\rho_0} - \frac{MP_0}{\rho_0} \log_e 2}{\frac{MP_0}{\rho_0} - \frac{MP_0}{\rho_0} \log_e 2} = \frac{\frac{MP_0}{\rho_0} - \frac{MP_0}{\rho_0} \log_e 2}{\frac{5MP_0}{\rho_0}}$$


$$\Rightarrow \frac{2(1 - \log_e 2)}{5}$$

Heat is rejected in C to A and A to B

$$Q_{\text{rejected}} = \frac{MP_0}{\rho_0} \log_e 2 + \frac{3}{2} \frac{MP_0}{\rho_0}$$

59. A small block B of mass $m = 0.25$ kg lying on a frictionless horizontal table, is attached to a massless cord (breaking strength 40 N) passing through a narrow hole C at the center of the table. Initially when the block is revolving in a circle of radius $r_0 = 0.80$ m about a vertical axis through the hole, with a tangential speed of $v_0 = 4.00$ m/s; the tension in the string is T_0 and the kinetic energy of the block is K_0 . The string is then pulled down slowly from below, decreasing the radius of circular path from r_0 to r so that the kinetic energy of the block is now K and the tension in the string is T .

As a result,



(a) the tension $T = T_0 \frac{r_0^4}{r^4}$

(b) the kinetic energy $K = K_0 \frac{r_0^2}{r^2}$

(c) the radius r of the circular path just when the string breaks is 0.40 m

(d) the work done by the tension in the string in reducing the radius of circle from r_0 to $\frac{r_0}{2}$ is $4 K_0$

Answer (b, c)

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB


(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS



$$T_0 = \frac{mV_0^2}{r_0}$$

$$\& K_0 = \frac{1}{2}mV_0^2$$

$$\therefore T_0 = \frac{2K_0}{r_0}$$

$$T = \frac{mV^2}{r}$$

$$K = \frac{1}{2}mV^2$$

$$\therefore T = \frac{2K}{r}$$

and in this process, angular momentum will remain conserved.

$$mVr = mV_0r_0$$

$$\therefore K = K_0 \frac{r_0^2}{r^2}$$

$$T = \frac{mV^2}{r}$$

$$\therefore \frac{T}{T_0} = \frac{V^2 r_0}{r V_0^2} = \frac{r_0^3}{r^3}$$

$$T = 40 = \frac{mV^2}{r}$$

$$\Rightarrow r = \frac{mV^2}{40} = \frac{0.25 \times V_0^2 r_0^2}{40 r^2}$$

$$r = \left(\frac{0.25 \times V_0^2 \times r_0^2}{40} \right)^{1/3} = 0.4$$

$$W = \frac{1}{2}mV^2 - \frac{1}{2}mV_0^2$$

$$\Rightarrow W = \frac{1}{2}mV_0^2 \left(\frac{r_0^2}{r^2} - 1 \right)$$

$$\Rightarrow W = 3K_0$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

60. A circular coil of thin insulated copper wire ($N = 2000$ turns), wrapped around an iron cylinder of cross-section area $\Delta S = 0.001 \text{ m}^2$, is connected to a suspended type moving coil ballistic galvanometer. The suspended rectangular coil of the galvanometer is of mass $m = 80 \text{ g}$, length $\ell = 5 \text{ cm}$, breadth $b = 3 \text{ cm}$ and has $n = 100$ turns of fine copper wire wound on a non-metallic frame of ivory. This rectangular coil of the galvanometer is free to execute torsional oscillations in a radial magnetic field $B = 0.1 \text{ tesla}$.


The galvanometer is being used to measure the charge by employing the formula $q = \frac{T}{2\pi} \frac{c'}{nAB} \theta$. [Given that the moment of inertia of the oscillating coil about the vertical axis is $I = 2.7 \times 10^{-6} \text{ kg m}^2$ and the torsional constant (torsional rigidity) of the suspension fiber is $c = 3.0 \times 10^{-3} \text{ Nm/radian}$: $A = \ell \times b$ is the area of the coil]

When the magnetic induction of $1.0 \text{ weber per meter}^2$, perpendicular to the plane of the circular coil, is reversed (in opposite direction), a deflection of 40 mm is observed on a scale placed 1.0 meter away in front of the reflecting mirror attached with the suspension fiber of the rectangular coil. The correct statement(s) is/are

- (a) the time period of the oscillating rectangular coil is $T = 0.19\text{s}$
- (b) the net change in flux through the circular coil wrapped on the iron cylinder is 4.0 weber
- (c) the induced charge in the circular coil wrapped on the iron cylinder is $q_{\text{ind}} = 240 \mu\text{C}$
- (d) total resistance of the circuit containing the circular coil is $R = 33.3 \text{ k}\Omega$

Answer (a, b, d)

Sol.



Circular
Coil. $N = 2000$

$$T = 2\pi \sqrt{\frac{I}{C}}$$

$$= 2\pi \sqrt{\frac{2.7 \times 10^{-6}}{3 \times 10^{-3}}}$$

$$= 2\pi \sqrt{9 \times 10^{-4}}$$

1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS

$$= 6\pi \times 10^{-2} = 0.19 \text{ s}$$

$$\Rightarrow a$$

$$\Delta\phi = (NBA)^2$$

$$= 2 \times 2000 \times 1 \times 0.001$$

$$= 4 \text{ wb}$$

$$\Rightarrow b$$

$$\theta = \frac{X}{d} = 40 \times 10^{-3} \text{ rad}$$

So deflection of torsion pendulum $\phi = \theta/2$

$$q = \frac{0.19 \times 3 \times 10^{-3} \times 40 \times 10^{-3}}{2\pi \times 100 \times 0.1 \times 15 \times 10^{-4} \times 2}$$

$$= \frac{6\pi \times 10^{-2} \times 120 \times 10^{-6}}{2\pi \times 10 \times 15 \times 10^{-4} \times 2}$$

$$= \frac{720}{30 \times 2} \times 10^{-2-6-1+4} = 120 \mu\text{C}$$

$$(d) \frac{\Delta\phi}{R} = q$$

$$R = \frac{\Delta\phi}{q} = \frac{(2NBA)}{q}$$

$$= \frac{4}{120 \times 10^{-6}}$$

$$= \frac{4000}{120} \times 10^3$$

$$= \frac{800}{24} \times 10^3$$

$$= 33.3 \text{ k}\Omega$$

$$\Rightarrow d$$



1484 Students Scored Above MAS

420

Classroom Students
Qualified in
NSEs 2024-25

(Group A & B)

49
NSEA

(Group A & B)

229
NSEB

(Group A & B)

70
NSEC

(Group A & B)

38
NSEP

34
NSEJS