

DATE: 21/06/2026

Test Booklet Code



**50**

**SUSHRUT**

Corporate Office: 3rd Floor, Incuspaze Campus-2, Plot No. 13,  
Sector-18, Udyog Vihar, Gurugram, Haryana - 122015.

## Questions & Answers

for

Time : 3 hrs. 15 min.

M.M. : 720

## NEET (UG)-2026 (Re-Examination)

### Important Instructions:

1. The test is of **3 hours 15 minutes** duration and the Test Booklet contains **180** multiple choice questions (Four options with a single correct answer) from **Physics, Chemistry & Biology (Botany and Zoology)**.
2. Each question carries **4 marks**. For each correct response, the candidate will get **4 marks**. For each incorrect response, **1 mark** will be deducted from the total scores. The maximum marks are **720**.
3. Use **Blue / Black Ball Point Pen only** for writing particulars on this page / marking responses on Answer Sheet.
4. Rough work is to be done in the space provided for this purpose in the Test Booklet only.
5. On completion of the test, the candidate **must handover the Answer Sheet (original & office copy) to the Invigilator** before leaving the Room / Hall. The candidates are allowed to take away this Test Booklet with them.
6. The **CODE** for this Booklet is **50**.
7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet. Use of white fluid for correction is **NOT** permissible on the Answer Sheet.
8. Each candidate must show on-demand his/her Admit Card to the Invigilator.
9. No candidate, without special permission of the Centre Superintendent or Invigilator, would leave his/her seat.
10. Use of Electronic/Manual Calculator is prohibited.
11. The candidates are governed by all Rules and Regulations of the examination with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of this examination.
12. No part of the **Test Booklet** and **Answer Sheet** shall be detached under any circumstances.
13. The candidates will write the Correct Test Booklet Code as given in the Test Booklet / Answer Sheet in the Attendance Sheet.

**PHYSICS**

1. A particle of mass  $M$  moves along a horizontal  $x$  axis from  $x = 0$  to  $x = L$ . The coefficient of kinetic friction varies as a function of  $x$  as  $\mu_k(x) = \mu_0 - \alpha x$ , where  $\mu_0, \alpha$  are constants of appropriate dimensions, so that  $\mu_k(L) = 0$ . The total work done by the frictional force during the motion is  $n\mu_0 MgL$ , where  $g$  is the acceleration due to gravity. The value of  $n$  is:

- (1) 3 (2) 1  
(3)  $\frac{1}{3}$  (4)  $\frac{1}{2}$

**Answer (4)**


2. The mean free path of molecules in an ideal gas  $A$  is half that of another ideal gas  $B$ . The diameter of the spherical molecules of gas  $A$  is twice the diameter of the molecules of  $B$ . If number densities of the gases  $A$  and  $B$  are  $n_A$  and  $n_B$ , respectively, the correct option is:

- (1)  $n_A = n_B$  (2)  $n_A = 2n_B$   
(3)  $n_A = \frac{1}{4}n_B$  (4)  $n_A = \frac{1}{2}n_B$

**Answer (4)**

3. Two identical inductors are connected in two different configurations  $P$  and  $Q$ , where a time varying current  $I(t)$  is flowing, as shown in the figure. The induced emf between points  $a$  and  $b$  for configuration  $P$  is  $E_P$  and that for configuration  $Q$  is  $E_Q$ . The ratio  $E_P/E_Q$  is:

[Neglect the effect of mutual inductance.]



- (1)  $\frac{1}{4}$  (2)  $\frac{1}{2}$   
(3) 1 (4) 2

**Answer (4)**

4. For sound waves, if the number of nodes for the 5<sup>th</sup> harmonic of an open-ended pipe is  $n$  and that for the 9<sup>th</sup> harmonic of the same pipe with one of its ends closed is  $m$ , the ratio  $\frac{n}{m}$  is

- (1)  $\frac{5}{9}$  (2)  $\frac{9}{5}$   
(3) 1 (4)  $\frac{3}{5}$

**Answer (3)**


5. Consider a long solenoid of length  $l$  and radius  $r$ . If  $n$  is the number of turns per unit length and  $\mu_0$  is the permeability of free space, the inductance of the solenoid is :
- (1)  $\mu_0 \pi n^2 r^2 l$  (2)  $\mu_0 n^2 r^2 l$   
 (3)  $(\mu_0/2\pi)n^2 r^2 l$  (4)  $2\mu_0 \pi n^2 r^2 l$

**Answer (1)**

6. Consider a particle moving along a straight line, whose position as a function of time is given by  $s(t) = \alpha t^2 - \beta t + \gamma$ , where  $\alpha = 1 \text{ ms}^{-2}$ ,  $\beta = 6 \text{ ms}^{-1}$  and  $\gamma = 5 \text{ m}$ . The average speed of the particle, in  $\text{ms}^{-1}$  from  $t = 0$  to  $t = 6 \text{ s}$  is:
- (1) 12 (2) 6  
 (3) 3 (4) 0

**Answer (3)**

7. Consider the following nuclear reaction



Take masses of  ${}^{238}\text{U}$ ,  ${}^{234}\text{Th}$  and  ${}^4\text{He}$  as 238.050 u, 234.043 u and 4.003 u, respectively. The Q value for the reaction, in keV, is:

[Given:  $1 \text{ u} = 931.5 \text{ MeV } c^{-2}$ ]

- (1) 3726 (2) 3730  
 (3) 3736 (4) 3740


**Answer (1)**

8. A beam of light falls on a metal surface such that photo-electrons are generated. If power of the light source starts to decrease linearly with time  $t$ , then variation of the photocurrent  $I$  and magnitude of the stopping potential  $|V|$  with time is best represented by:



**Answer (1)**


9. Three identical capacitors,  $P$ ,  $Q$  and  $S$ , each of the capacitance  $C$ , are connected to a battery of voltage  $V$ , as shown in the figure. If the energy stored in the capacitor  $P$  and total energy stored in the system are  $U_P$  and  $U_T$ , respectively, then the ratio  $\frac{U_P}{U_T}$  is:



- (1)  $2/3$  (2)  $1/3$   
 (3)  $1/2$  (4)  $1/6$

**Answer (4)**


10. Water flows in a streamline motion through a horizontal pipe of circular cross-section as shown in the figure. The pressure difference of water between  $P$  and  $Q$  is  $15 \text{ Nm}^{-2}$ . The area of cross-section at  $P$  and  $Q$  are  $40 \text{ cm}^2$  and  $20 \text{ cm}^2$ , respectively. The rate of flow of water through the pipe, in  $\text{cm}^3\text{s}^{-1}$ , is:  
[Take density of water =  $1000 \text{ kg m}^{-3}$ ]



- (1) 100  
(2) 200  
(3) 300  
(4) 400

**Answer (4)**

11. A current  $I_0$  flows through a metallic circular loop of radius  $r$  as shown in the figure. Resistance of the segment  $ABC$  is half that of  $ADC$ . Magnitude of magnetic field at the centre  $O$  of the loop is :



- (1)  $\frac{\mu_0 I_0}{12r}$   
(2)  $\frac{\mu_0 I_0}{4r}$   
(3)  $\frac{\mu_0 I_0}{2r}$   
(4)  $\frac{\mu_0 I_0}{2\pi r}$

**Answer (1)**

12. In the measurement of viscosity of liquids using terminal velocity experiment, spherical balls of same radius but having different densities are used. The variation of the terminal velocity ( $v$ ) with the ratio of density of spherical ball ( $\sigma$ ) to density of the liquid ( $\rho$ ), is best represented by:



**Answer (1)**

13. Two planets  $P_1$  and  $P_2$  with equal mass have radii  $R_1$  and  $R_2$ , respectively, where  $R_2 = \frac{R_1}{2}$ . The escape speeds of  $P_1$  and  $P_2$  are  $v_1$  and  $v_2$ , respectively. Then  $\frac{v_2}{v_1}$  is:

- (1)  $\frac{1}{\sqrt{2}}$   
(2) 1  
(3)  $\sqrt{2}$   
(4) 2

**Answer (3)**

14. In a solar system, the time-period of revolution of a planet tracing a circular orbit of radius  $R$  is proportional to:
- (1)  $R^{1/2}$  (2)  $R^{3/2}$   
 (3)  $R^2$  (4)  $R^3$


**Answer (2)**

15. Two infinitely long parallel conducting wires  $A$  and  $B$  carry currents  $I$  and  $2I$ , respectively, in the same direction. The wire  $A$  has uniform mass per unit length  $\lambda$  and lies on an insulated floor. The wire  $B$  is kept fixed at a height  $h$  above the floor. The minimum magnitude of  $h$  so that the wire  $A$  does not rise from the floor is: [g is the acceleration due to gravity and  $\mu_0$  is the permeability of free space.]

- (1)  $\frac{\mu_0 I^2}{2\pi\lambda g}$  (2)  $\frac{\mu_0 I^2}{\pi\lambda g}$   
 (3)  $\frac{2\mu_0 I^2}{\pi\lambda g}$  (4)  $\frac{4\mu_0 I^2}{\pi\lambda g}$

**Answer (2)**

16. An ideal Zener diode with breakdown voltage of  $-3$  V is reverse biased with a negative input voltage  $V_i = -5$  V. The magnitude of voltage difference between point  $B$  and  $A$  is:



- (1) 3 V (2) 2 V  
 (3) 1 V (4) 0 V

**Answer (2)**

17. In an adiabatic expansion, the temperature of one mole of an ideal monatomic gas ( $\gamma = 5/3$ ) decreases from 60 K to 50 K. The work done by the gas in the process is:

(Take the universal gas constant as  $R = 8.3 \text{ J mol}^{-1} \text{ K}^{-1}$ )

- (1) 41.5 J (2) 83 J  
 (3) 124.5 J (4) 166 J

**Answer (3)**

18. A ray of light with wavelength  $\lambda$  is incident on three different photoelectric cells namely 1, 2 and 3. The threshold wavelength of these photoelectric cells are  $\lambda_1$ ,  $\lambda_2$ , and  $\lambda_3$ , respectively and the magnitude of stopping potentials of these cells are  $V_1$ ,  $V_2$  and  $V_3$ , respectively. The relation between  $\lambda$  and threshold wavelengths are  $\lambda_1 < \lambda$ ,  $\lambda_2 > \lambda$  and  $\lambda_3 \gg \lambda$ . The correct option is:

- (1)  $V_1 = 0$ ,  $V_2 < V_3$  (2)  $V_1 = 0$ ,  $V_2 > V_3$   
 (3)  $V_1 > V_2$ ,  $V_3 = 0$  (4)  $V_1 < V_2$ ,  $V_3 = 0$

**Answer (1)**

19. A photon and an electron, each of 20 eV energy, move in free space. The ratio of linear momentum of electron  $p_e$  to that of photon  $p_{Ph}$ ,  $\frac{p_e}{p_{Ph}}$  is:

[Take speed of light =  $3 \times 10^8 \text{ ms}^{-1}$ , charge of electron =  $-1.6 \times 10^{-19} \text{ C}$  and mass of electron =  $9 \times 10^{-31} \text{ kg}$ ]

- (1)  $\frac{2}{450}$  (2)  $\frac{1}{250}$   
 (3) 225 (4) 275


**Answer (3)**

20. Which of the following measurements require 'index correction'?
- (1) Measurement of resistance of a wire using meter bridge
  - (2) Measurement of gravitational acceleration using simple pendulum
  - (3) Measurement of focal length of lenses using optical bench
  - (4) Measurement of speed of sound using resonance tube

**Answer (3)**

21. A unit positive point charge is taken slowly through an infinitesimally thin tube that is inside a charged dielectric sphere of radius  $R$ , having uniform positive charge density  $\rho$ , as shown in the figure. The initial and final positions of the charge are marked by  $A$  and  $B$  at distance  $2R$  and  $3R$  respectively, from the centre of the sphere. In this process, the magnitude of the total work done on the point charge is  $\frac{\rho R^2}{n\epsilon_0}$ . The value of  $n$  is :


( $\epsilon_0$  is the permittivity of vacuum)



- (1) 2
- (2) 6
- (3) 9
- (4) 18

**Answer (4)**

22. Consider three media  $P$ ,  $Q$  and  $R$  with refractive indices 1, 1.25, and 1.5 respectively. The medium  $Q$  having a thickness of 5 cm is placed between extended media  $P$  and  $R$  as shown in the figure. An object  $O$  is placed at the centre of medium  $Q$ . If viewed from medium  $P$  near the normal direction, the apparent depth of  $O$  is  $h_1$ . For similar observation from medium  $R$ , the apparent depth is  $h_2$ . The value of  $|h_1 - h_2|$ , in cm, is:




- (1) 0
- (2) 1
- (3) 2
- (4) 3

**Answer (2)**






31. A thin horizontal disc is rotating about a vertical axis passing through its fixed centre  $O$ . Its angular momentum is  $L_A$  and  $L_B$  computed about points  $A$  and  $B$ , respectively, with  $OB = 2 \times OA$ . The value of  $\frac{L_A}{L_B}$  is:



- (1)  $\frac{1}{4}$  (2)  $\frac{1}{2}$   
(3) 1 (4) 2


**Answer (3)**

32. A conducting loop of finite resistance lies on the  $x - y$  plane. There is a constant magnetic field in the  $z$  direction. The area of the loop varies with time  $t$ , as  $A = A_0(1 + \sin t)$  in appropriate units. The figure that correctly indicates the qualitative behaviour of the power  $P$  dissipated in the loop as a function of time is:



**Answer (2)**

33. A point charge  $Q$  is placed inside a cavity within a solid isolated conducting sphere. Consider points  $A$ ,  $B$  and  $C$  as shown in the figure, where the magnitudes of the electric fields are  $E_A$ ,  $E_B$ ,  $E_C$ , respectively. The points  $B$  and  $C$  are at the same distance from the center of the solid sphere. The correct option is :



- (1)  $E_A = 0, E_B = E_C$  (2)  $E_A \neq 0, E_B = E_C$   
(3)  $E_A = 0, E_B > E_C$  (4)  $E_A \neq 0, E_B < E_C$

**Answer (2)**





34. Consider a fixed uniformly charged insulating sphere with radius  $R$  and total charge  $+Q$ . A point charge  $-q$  ( $q \ll Q$ ) with mass  $m$  is released from rest at a distance of  $3R$  from the centre of the charged sphere. When the point charge reaches the surface of the sphere, its speed is:

( $\epsilon_0$  is the permittivity of vacuum, neglect gravitational forces).

- (1)  $\sqrt{\frac{3Qq}{4\pi\epsilon_0 mR}}$  (2)  $\sqrt{\frac{2Qq}{3\pi\epsilon_0 mR}}$   
(3)  $\sqrt{\frac{Qq}{3\pi\epsilon_0 mR}}$  (4)  $\sqrt{\frac{Qq}{4\pi\epsilon_0 mR}}$


**Answer (3)**

35. In Geiger-Marsden experiment, the number of scattered  $\alpha$ -particles  $N(\theta)$  is plotted as a function of scattering angle  $\theta$ . Which of the following options represents the correct plot?



**Answer (3)**

36. Consider two circuits, (A) and (B), each having two resistors. One of them has a positive temperature coefficient of resistance,  $+\alpha$ , while the other one has a negative temperature coefficient,  $-\alpha$ , as shown in the figure. The current through these circuits are denoted by  $I_A$  and  $I_B$ . At initial temperature, the resistance of the two resistors is  $R_0$ . As the temperature is increased, the correct option that describes the variation of current in these circuits is:



- (1)  $I_A$  remains constant while  $I_B$  increases
- (2)  $I_A$  decreases while  $I_B$  increases
- (3)  $I_A$  increases while  $I_B$  decreases
- (4) Both  $I_A$  and  $I_B$  remain constant

**Answer (1)**

37. Consider that  $\sigma_s$ ,  $k_B$ ,  $b$  represents Stefan-Boltzmann constant, Boltzmann constant and Wien's displacement law constant, respectively. The dimension of  $\sigma_s k_B^{-1} b$  is


- (1)  $[L^{-1}T^{-1}K^{-2}]$
- (2)  $[L^{-1}K^{-2}]$
- (3)  $[L^{-1}T^{-1}K^{-3}]$
- (4)  $[L^{-1}T^{-1}K^{-4}]$

**Answer (1)**

38. An electromagnetic wave travelling in a lossless dielectric medium having a dielectric constant,  $\epsilon_r = 9$ , has the electric field,  $E_x = E_0 \sin(kz - 2\pi \times 10^6 t)$  Vm<sup>-1</sup> where  $E_0$  is the amplitude and  $k$  is the wave vector. Among the following options, the **incorrect** choice is
- (1) The speed of the electromagnetic wave inside the medium is  $10^8$  ms<sup>-1</sup>
  - (2) The wavelength of the electromagnetic wave inside the medium is 300 m
  - (3) The magnetic field is given by the relation  $B_y = \frac{B_0}{v} \sin(kz - 2\pi \times 10^6 t)$  where  $v$  is the speed of the electromagnetic wave inside the medium
  - (4) The direction of propagation of the electromagnetic wave is along +z

**Answer (2)**

39. One mole of an ideal monatomic gas undergoes a cyclic process as shown in the figure. The total heat supplied to the gas is:



- (1) 400 J
- (2) 500 J
- (3) 600 J
- (4) 800 J

**Answer (3)**

40. Consider that an electron is revolving in an excited state of Hydrogen atom with velocity  $\sqrt{25.6} \times 10^5$  ms<sup>-1</sup>. The radius of the orbit is  $x \times 10^{-9}$  m. The value of  $x$  is:

[Take the mass of electron to be  $9 \times 10^{-31}$  kg, charge of electron =  $-1.6 \times 10^{-19}$  C and  $\frac{1}{4\pi\epsilon_0} = 9 \times 10^9$  Nm<sup>2</sup>C<sup>-2</sup>]

- (1) 4
- (2) 3
- (3) 2
- (4) 1

**Answer (4)**

41. A car travels on a circular racetrack of radius 50 m, which is banked at an angle  $\theta$ . If the car travels at a speed  $10$  ms<sup>-1</sup>, then the wear and tear on its tyres is minimum. Taking the acceleration due to gravity to be  $10$  ms<sup>-2</sup>, the value of  $\theta$  is:

- (1)  $\tan^{-1}\left(\frac{1}{5}\right)$
- (2)  $\tan^{-1}\left(\frac{2}{5}\right)$
- (3)  $\tan^{-1}(\sqrt{3}/2)$
- (4)  $\tan^{-1}(2\sqrt{3})$

**Answer (1)**

