

CHEMISTRY

46. Match the species in **List-I** with their geometry in **List-II**.

	List-I		List-II
A.	PCl_5	I.	Tetrahedral
B.	BrF_5	II.	Square Planar
C.	BF_4^-	III.	Trigonal bipyramidal
D.	$[\text{Ni}(\text{CN})_4]^{2-}$	IV.	Square pyramidal

Choose the **correct** answer from the options given below:

(1) A-III, B-IV, C-I, D-II

(2) A-III, B-I, C-II, D-IV

(3) A-III, B-II, C-I, D-IV

(4) A-IV, B-III, C-I, D-II

Answer (1)

47. The amino acid that gives a red-blood colour on treating its sodium fusion extract with sodium nitroprusside is

(1) threonine

(2) methionine

(3) serine

(4) leucine

Answer (2)

48. Given below are two statements:

Statement-I : Oxidation of p-nitrotoluene with acidic KMnO_4 gives an acid that is stronger than benzoic acid.

Statement-II : Reduction of p-nitrotoluene with Sn/HCl followed by neutralization gives an amine that is more basic than aniline.

In light of the above statements, choose the **most appropriate** answer from the options given below.

(1) Both **Statement-I** and **Statement-II** are incorrect

(2) **Statement-I** is correct but **Statement-II** is incorrect

(3) **Statement-I** is incorrect but **Statement-II** is correct

(4) Both **Statement-I** and **Statement-II** are correct

Answer (4)

49. The standard electrode potential (E°) for the half-cell reaction $\text{Fe}^{3+} + e^- \rightarrow \text{Fe}^{2+}$ at 298 K is

(Given : $E^\circ(\text{Fe}^{3+}/\text{Fe}) = -0.04 \text{ V}$ and $E^\circ(\text{Fe}^{2+}/\text{Fe}) = -0.44 \text{ V}$ at 298 K)

(1) +0.76 V






(2) -0.48 V

(3) +0.92 V

(4) +0.40 V

Answer (1)

50. The compound that **CANNOT** be obtained from the aldol condensation reaction shown below, is



Answer (4)

51. For a salt **XY**, which is a strong electrolyte, the plot of Λ_m versus \sqrt{c} has a slope of $-90.0 \text{ S cm}^2 \text{ mol}^{-3/2} \text{ L}^{1/2}$ at 298 K. At 0.01 M concentration of **XY**, the value of Λ_m is $145.0 \text{ S cm}^2 \text{ mol}^{-1}$. The limiting molar conductivity of Y^- ion ($\lambda_{\text{Y}^-}^0$, in $\text{S cm}^2 \text{ mol}^{-1}$) at 298 K will be

(Given : $\lambda_{\text{X}^+}^0 = 74.0 \text{ S cm}^2 \text{ mol}^{-1}$)

(1) 100.0

(2) 90.0

(3) 76.0

(4) 80.0

Answer (4)

52. Given below are two statements: One is labelled as **Assertion A** and the other is labelled as **Reason R**.

Assertion A: The first ionization enthalpy of O is lower than that of N and F.

Reason R: The loss of an electron from O leads to stable half-filled p orbital.

In light of the above statements, choose the **most appropriate** answer from the options given below:

(1) Both **A** and **R** are correct and **R** is **NOT** the correct explanation of **A**

(2) **A** is correct but **R** is not correct.

(3) **A** is not correct but **R** is correct

(4) Both **A** and **R** are correct and **R** is the correct explanation of **A**

Answer (4)

53. According to crystal field theory, the correct order of ligands with respect to their decreasing order of field strength is

(1) $\text{CO} > \text{H}_2\text{O} > \text{NH}_3 > \text{Cl}^-$

(2) $\text{Cl}^- > \text{H}_2\text{O} > \text{NH}_3 > \text{CO}$

(3) $\text{Cl}^- > \text{NH}_3 > \text{H}_2\text{O} > \text{CO}$

(4) $\text{CO} > \text{NH}_3 > \text{H}_2\text{O} > \text{Cl}^-$

Answer (4)

54. Given below are two statements:

Statement-I : $[\text{Fe}(\text{ox})_3]^{3-}$ is chiral.

Statement-II : *trans* - $[\text{Cr}(\text{H}_2\text{O})_2(\text{ox})_2]^-$ is chiral.


(Given : $\text{oxH}_2 = \text{HOOC} - \text{COOH}$)

In light of the above statements, choose the **most appropriate** answer from the options given below:

- (1) Both **Statement-I** and **Statement-II** are incorrect
- (2) **Statement-I** is correct but **Statement-II** is incorrect
- (3) **Statement-I** is incorrect but **Statement-II** is correct
- (4) Both **Statement-I** and **Statement-II** are correct

Answer (2)


55. For the following reaction sequence, choose the correct option



- (1) **P** and **Q** are aromatic compounds
- (2) If **P** gives a carboxylic acid on acidification, **Q** gives a poisonous gas on exposure to air and light
- (3) Both **P** and **Q** are carbonyl compounds
- (4) If **P** is the sodium salt of a carboxylic acid, **Q** is a primary alcohol

Answer (2)

56. For an elementary chemical reaction, the Arrhenius plot is given below.



If the energy of activation is 6.64 kJ mol^{-1} and $R = 8.3 \text{ J K}^{-1} \text{ mol}^{-1}$, the temperature at which the rate constant becomes $e^2 \text{ min}^{-1}$, is

- | | |
|-----------|-----------|
| (1) 150 K | (2) 200 K |
| (3) 250 K | (4) 125 K |

Answer (2)

57. Arrange the following compounds in the increasing order of polarity
- $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$
 - $\text{CH}_3\text{CH}_2\text{OH}$
 - CH_3COCH_3
 - CH_3COOH

Choose the correct answer from the options given below.

- $\text{C} < \text{A} < \text{D} < \text{B}$
- $\text{C} < \text{A} < \text{B} < \text{D}$
- $\text{A} < \text{C} < \text{B} < \text{D}$
- $\text{A} < \text{B} < \text{C} < \text{D}$

Answer (3)

58. A protein undergoes reversible thermal denaturation from its initial state **N** to denatured state **D** according to $\text{N} \rightleftharpoons \text{D}$. At 60°C , the concentrations of both **N** and **D** are equal at equilibrium, and the standard enthalpy change of denaturation is 666 kJ mol^{-1} . The standard entropy change (ΔS° in $\text{kJ K}^{-1} \text{ mol}^{-1}$) of the protein upon denaturation at 60°C is closest to

- | | |
|------------|-----------|
| (1) 2000.0 | (2) 333.0 |
| (3) 11.1 | (4) 2.0 |

Answer (4)

59. Consider the following statements about the solutions formed by mixing two liquids.
- An ideal solution thus formed obeys Raoult's law throughout the composition range.
 - Mixture of chloroform and acetone shows negative deviation from Raoult's law.
 - Mixture of aniline and phenol shows positive deviation from Raoult's law.
- B and C only
 - A only
 - A and C only
 - A and B only

Answer (4)

60. Given below are two statements :

Statement-I : Heating NaCl with concentrated H_2SO_4 and MnO_2 results in oxidation of Mn .

Statement-II : Heating NaI with concentrated H_2SO_4 and MnO_2 results in reduction of Mn .


In light of the above statements, choose the **most appropriate** answer from the options given below.

- Both **Statement-I** and **Statement-II** are incorrect
- Statement-I** is correct but **Statement-II** is incorrect
- Statement-I** is incorrect but **Statement-II** is correct
- Both **Statement-I** and **Statement-II** are correct


Answer (3)

61. Given below are two statements :

Statement I : *trans*-But-2-ene upon treatment with Br₂ in CCl₄ gives the following product.



Statement II : *cis*-But-2-ene upon treatment with alkaline KMnO₄ gives the following product.



In the light of the above statements, choose the **most appropriate** answer from the options given below.

- (1) Both **Statement I** and **Statement II** are incorrect
- (2) **Statement I** is correct but **Statement II** is incorrect
- (3) **Statement I** is incorrect but **Statement II** is correct
- (4) Both **Statement I** and **Statement II** are correct


Answer (3)

62. The highest occupied molecular orbital for Ne₂ is

- (1) σ_{2p}
- (2) π_{2p}^*
- (3) σ_{2p}^*
- (4) π_{2p}

Answer (3)

63. The following carbocation is stabilized by the interaction of the empty *p* orbital with



- (1) empty σ and empty π^* orbitals
- (2) empty σ^* and filled π orbitals
- (3) empty σ^* and empty π^* orbitals
- (4) filled σ and filled π orbitals

Answer (4)

64. The green paramagnetic species formed by heating KMnO₄ at 513 K is

- (1) Mn₃O₄
- (2) MnO
- (3) KO₂
- (4) K₂MnO₄

Answer (4)

65. The correct statement is
- (1) Beryllium has three valence orbitals.
 - (2) Magnesium has a maximum covalency of four.
 - (3) Aluminium has five valence orbitals.
 - (4) Boron has a maximum covalency of four.

Answer (4)

66. The correct statement about peptides and proteins is
- (1) Only the proteins having a quaternary structure are biologically active
 - (2) In β -pleated sheet structures, peptide chains are held together by intermolecular hydrogen bonds
 - (3) In α -helices, the polypeptide chain is twisted into a left-handed screw (helix) through intramolecular hydrogen bonds
 - (4) Tertiary structure of proteins has two or more polypeptide subunits

Answer (2)

67. $2A \xrightarrow{k} B$ is a zero-order reaction, where $k = 1.0 \text{ mol L}^{-1} \text{ min}^{-1}$. If the initial concentration of A is 2 M, then the time taken to complete 75% of the reaction will be
- (1) 0.75 min
 - (2) 1.0 min
 - (3) 2.0 min
 - (4) 1.5 min

Answer (1)

68. Given below are two statements: One is labelled as **Assertion A** and the other is labelled as **Reason R**.

Assertion A: Generally, 3d transition metals have high melting points.


Reason R: Involvement of 3d-electrons in addition to 4s-electrons in the interatomic metallic bonding.

In light of the above statements, choose the **most appropriate** answer from the options given below:

- (1) Both **A** and **R** are correct and **R** is **NOT** the correct explanation of **A**
- (2) **A** is correct but **R** is not correct.
- (3) **A** is not correct but **R** is correct
- (4) Both **A** and **R** are correct and **R** is the correct explanation of **A**

Answer (4)

69. Consider the following schematic plots of orbital wavefunction (ψ_r) against distance (r) from the nucleus.







The figure representing two radial nodes in the orbital is

- (1) B (2) C
(3) D (4) A

Answer (2)

70. Identify the reactions which give aniline as the major product.



Choose the correct answer from the options given below.

- (1) B and D only (2) A and C only
(3) C and D only (4) A and B only

Answer (1)

71. Among the species given below, the spin-only magnetic moment is highest for
(Given: Atomic number of Ti = 22, Mn = 25, Fe = 26 and Co = 27)

- (1) $[\text{Fe}(\text{CN})_6]^{3-}$ (2) $[\text{Co}(\text{NH}_3)_6]^{3+}$
(3) $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$ (4) $[\text{Mn}(\text{CN})_6]^{3-}$

Answer (4)

72. The complex which has *facial* and *meridional* isomers is

(Given : py = pyridine and en = H₂N – CH₂ – CH₂ – NH₂)

- (1) [Cr(H₂O)₆]³⁺ (2) [Co(NH₃)₄(H₂O)₂]³⁺
(3) [Ni(en)₂(H₂O)₂]²⁺ (4) [Cr(py)₃(Cl)₃]


Answer (4)

73. A 1 : 3 electrolyte in an aqueous solution is

- (1) [CoCl(NH₃)₅]Cl₂ (2) [Co(NH₃)₆]Cl₃
(3) [Co(NH₃)₃(NO₂)₃] (4) [CoCl₂(NH₃)₄]Cl

Answer (2)

74. Consider the reversible processes for 1.0 mol of an ideal gas as shown in the figure.



w_1, w_2, w_3 and w_4 represent work done (in calories) in the processes 1, 2, 3 and 4, respectively; ΔU_2 and ΔU_4 are changes in the internal energy for the processes 2 and 4, respectively.

[use $R = 2 \text{ cal K}^{-1} \text{ mol}^{-1}$]

The correct option is

- (1) $w_2 + w_4 = \Delta U_2 - \Delta U_4$ (2) $w_1 + w_2 = 2T_1 \ln \frac{V_2}{V_1}$
(3) $w_1 + w_2 + w_3 + w_4 = 0$ (4) $w_1 + w_3 = -2T_1 \ln \frac{V_2}{V_1} - 2T_2 \ln \frac{V_4}{V_3}$

Answer (4)

75. In an acidic medium, 10 mL of 0.25 M oxalic acid is titrated with KMnO_4 solution. If the volume of KMnO_4 solution required to reach end point is 10 mL, the strength of the KMnO_4 solution is
- (1) 0.20 M (2) 0.25 M
 (3) 0.15 M (4) 0.10 M

Answer (4)

76. The lanthanide ion having four unpaired electrons is
 (Given : Atomic numbers of Ce = 58, Nd = 60, Tb = 65 and Ho = 67)
- (1) Ce^{3+} (2) Tb^{3+}
 (3) Ho^{3+} (4) Nd^{3+}

Answer (3)


77. The correct decreasing order of oxidation state of the underlined atom in each molecule is
- (1) $\underline{\text{N}}_2\text{O}_5 > \underline{\text{Al}}_2\text{O}_3 > \underline{\text{H}}_2\underline{\text{S}}$ (2) $\underline{\text{Pb}}\text{O}_2 > \underline{\text{N}}_2\text{O}_3 > \underline{\text{S}}\text{O}_3$
 (3) $\underline{\text{P}}_4\text{O}_6 > \underline{\text{Cl}}_2\text{O}_7 > \underline{\text{Al}}\text{H}_3$ (4) $\underline{\text{P}}_4\text{O}_{10} > \underline{\text{S}}\text{O}_3 > \underline{\text{H}}_2\underline{\text{O}}$

Answer (1)

78. The formula of tetraammineaquachloridocobalt(III) chloride is
- (1) $[\text{Co}(\text{NH}_3)_4]\text{Cl}_3 \times \text{H}_2\text{O}$ (2) $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]\text{Cl}$
 (3) $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]\text{Cl}_2$ (4) $[\text{Co}(\text{NH}_3)_4\text{Cl}_2] \times \text{H}_2\text{O}$

Answer (3)

79. Consider the following reaction, and choose the correct option.



- (1) Compound **P** can be prepared by treating benzene with anhydrous AlCl_3 and CH_3COCl
 (2) On treatment with bromine water, compound **P** gives a white precipitate
 (3) Compound **P** is obtained by the hydrogenation of benzoyl chloride with Pd on BaSO_4
 (4) On treating compound **P** with saturated NaHCO_3 solution, brisk effervescence is observed

Answer (3)

80. **Assertion A:** For an ideal solution formed by mixing liquids **P** and **Q**, $\Delta_{\text{mix}} H = 0$ and $\Delta_{\text{mix}} V = 0$

Reason R: No interactions occur between **P** and **Q**

In the light of the above statements, choose the **most appropriate** answer from the options given below.

- (1) Both **A** and **R** are correct but **R** is **NOT** the correct explanation of **A**
 (2) **A** is correct but **R** is not correct
 (3) **A** is not correct but **R** is correct
 (4) Both **A** and **R** are correct and **R** is the correct explanation of **A**

Answer (2)

81. Match the vitamins in **List I** with their sources in **List II**.

	List I		List II
A.	vitamin A	I.	meat
B.	vitamin B ₁₂	II.	sunflower oil
C.	vitamin E	III.	green leafy vegetables
D.	vitamin K	IV.	carrots

Choose the correct answer from the options given below.

(1) A-IV, B-I, C-II, D-III

(2) A-IV, B-II, C-I, D-III

(3) A-III, B-I, C-IV, D-II

(4) A-II, B-III, C-IV, D-I

Answer (1)

82. Among the following, the compound having conjugated double bonds is

(1) hepta-1,4-diene





(2) hepta-1,5-diene

(3) hepta-1,6-diene

(4) hepta-1,3-diene

Answer (4)

83. One of the products formed in the following reaction is



Answer (3)

84. Two moles of an ideal gas undergo free expansion from 10 L to 100 L at 300 K. The values of ΔS_{system} and $\Delta S_{\text{surroundings}}$ are

(R is universal gas constant)

(1) $\Delta S_{\text{system}} = 4.606 R$; $\Delta S_{\text{surroundings}} = -4.606 R$

(2) $\Delta S_{\text{system}} = 0$; $\Delta S_{\text{surroundings}} = 4.606 R$

(3) $\Delta S_{\text{system}} = 4.606 R$; $\Delta S_{\text{surroundings}} = 0$

(4) $\Delta S_{\text{system}} = 0$; $\Delta S_{\text{surroundings}} = 0$

Answer (3)

85. Among the following options, the correct trend in the electron gain enthalpy is

(1) $\text{Br} > \text{Cl} > \text{F} > \text{I}$

(2) $\text{Cl} > \text{F} > \text{Br} > \text{I}$

(3) $\text{I} > \text{Br} > \text{Cl} > \text{F}$

(4) $\text{F} > \text{Cl} > \text{Br} > \text{I}$

Answer (2)


86. The correct order of solubility of the given salts in water at 298 K is

Salt	K_{sp} at 298 K
AgBr	5.0×10^{-13}
Zn(OH) ₂	1.0×10^{-15}
Hg ₂ Cl ₂	1.3×10^{-18}

- (1) AgBr > Zn(OH)₂ > Hg₂Cl₂ (2) Hg₂Cl₂ > AgBr > Zn(OH)₂
 (3) Zn(OH)₂ > AgBr > Hg₂Cl₂ (4) Hg₂Cl₂ > Zn(OH)₂ > AgBr

Answer (3)

87. Consider the following reaction sequences and choose the correct option.



- (1) K and L are enantiomers
 (2) M and N are geometrical isomers
 (3) M and N are stereoisomers
 (4) K and L are geometrical isomers

Answer (4)

88. The amount of carbon dioxide evolved upon complete combustion of 116 g of n-butane is
 (Given: atomic mass in amu H = 1, C = 12 and O = 16)

- (1) 322 g (2) 176 g
 (3) 362 g (4) 352 g

Answer (4)

89. The numbers 17.0145 and 21.0235 were rounded to three figures after the decimal point. The resulting numbers, respectively, are

- (1) 17.015 and 21.023 (2) 17.014 and 21.024
 (3) 17.015 and 21.024 (4) 17.014 and 21.023

Answer (2)

90. In potash alum, the ratio of K⁺ and SO₄²⁻ ions is

- (1) 2 : 1 (2) 2 : 3
 (3) 3 : 2 (4) 1 : 2

Answer (4)