

DATE: 21/06/2026

Test Booklet Code



80

SUSHRUT

Corporate Office: 3rd Floor, Incuspaze Campus-2, Plot No. 13,
Sector-18, Udyog Vihar, Gurugram, Haryana - 122015.

Answers & Solutions for

Time : 3 hrs. 15 min.

M.M. : 720

NEET (UG)-2026 (Re-Examination)

Important Instructions:

1. The test is of **3 hours 15 minutes** duration and the Test Booklet contains **180** multiple choice questions (Four options with a single correct answer) from **Physics, Chemistry & Biology (Botany and Zoology)**.
2. Each question carries **4 marks**. For each correct response, the candidate will get **4 marks**. For each incorrect response, **1 mark** will be deducted from the total scores. The maximum marks are **720**.
3. Use **Blue / Black Ball Point Pen only** for writing particulars on this page / marking responses on Answer Sheet.
4. Rough work is to be done in the space provided for this purpose in the Test Booklet only.
5. On completion of the test, the candidate **must handover the Answer Sheet (original & office copy) to the Invigilator** before leaving the Room / Hall. The candidates are allowed to take away this Test Booklet with them.
6. The **CODE** for this Booklet is **80**.
7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet. Use of white fluid for correction is **NOT** permissible on the Answer Sheet.
8. Each candidate must show on-demand his/her Admit Card to the Invigilator.
9. No candidate, without special permission of the Centre Superintendent or Invigilator, would leave his/her seat.
10. Use of Electronic/Manual Calculator is prohibited.
11. The candidates are governed by all Rules and Regulations of the examination with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of this examination.
12. No part of the **Test Booklet** and **Answer Sheet** shall be detached under any circumstances.
13. The candidates will write the Correct Test Booklet Code as given in the Test Booklet / Answer Sheet in the Attendance Sheet.

CHEMISTRY

46. Match the species in **List-I** with their geometry in **List-II**.

	List-I		List-II
A.	PCl ₅	I.	Tetrahedral
B.	BrF ₅	II.	Square Planar
C.	BF ₄ ⁻	III.	Trigonal bipyramidal
D.	[Ni(CN) ₄] ²⁻	IV.	Square pyramidal

Choose the **correct** answer from the options given below:

(1) A-III, B-IV, C-I, D-II

(2) A-III, B-I, C-II, D-IV

(3) A-III, B-II, C-I, D-IV

(4) A-IV, B-III, C-I, D-II

Answer (1)

Sol.

	Compound	Hybridization of central atom/ion	Geometry
A.	PCl ₅	sp ³ d	Trigonal bipyramidal
B.	BrF ₅	sp ³ d ²	Square pyramidal
C.	BF ₄ ⁻	sp ³	Tetrahedral
D.	[Ni(CN) ₄] ²⁻	dsp ²	Square planar

47. The amino acid that gives a red-blood colour on treating its sodium fusion extract with sodium nitroprusside is

(1) threonine


(2) methionine

(3) serine

(4) leucine

Answer (2)

Sol. In case both nitrogen and sulphur are present in an organic compound, it gives blood red colour with sodium nitroprusside.



48. Given below are two statements:


Statement-I : Oxidation of p-nitrotoluene with acidic KMnO_4 gives an acid that is stronger than benzoic acid.

Statement-II : Reduction of p-nitrotoluene with Sn/HCl followed by neutralization gives an amine that is more basic than aniline.

In light of the above statements, choose the **most appropriate** answer from the options given below.


- (1) Both **Statement-I** and **Statement-II** are incorrect
- (2) **Statement-I** is correct but **Statement-II** is incorrect
- (3) **Statement-I** is incorrect but **Statement-II** is correct
- (4) Both **Statement-I** and **Statement-II** are correct

Answer (4)

Sol. • Oxidation of p-nitrotoluene with acidic KMnO_4 converts into p-nitrobenzoic acid.  because

of $-I$ effect and $-M$ effect, making p-nitrobenzoic acid as stronger acid than benzoic acid.

- Reduction of p-nitrotoluene with Sn/HCl followed by neutralization reduces the ($-\text{NO}_2$) to an amino group ($-\text{NH}_2$), forming p-toluidine.



i.e., p-methylaniline is more basic than aniline due to the $+I$ effect and hyperconjugation.


49. The standard electrode potential (E°) for the half-cell reaction $\text{Fe}^{3+} + e^- \rightarrow \text{Fe}^{2+}$ at 298 K is

(Given : $E^\circ(\text{Fe}^{3+}/\text{Fe}) = -0.04 \text{ V}$ and $E^\circ(\text{Fe}^{2+}/\text{Fe}) = -0.44 \text{ V}$ at 298 K)

- (1) $+0.76 \text{ V}$
- (2) -0.48 V
- (3) $+0.92 \text{ V}$
- (4) $+0.40 \text{ V}$

Answer (1)

Sol.



$$\Delta G^\circ_{\text{Fe}^{3+}/\text{Fe}} = \Delta G^\circ_{\text{Fe}^{3+}/\text{Fe}^{2+}} + \Delta G^\circ_{\text{Fe}^{2+}/\text{Fe}}$$

$$-nFE^\circ = -nFE^\circ + -nFE^\circ$$

$$-3F(-0.04) = -1FE^\circ + 2F(0.44)$$

$$3F(0.04) = -Fe^\circ + 0.88F$$






$$0.12 = -E^\circ + 0.88$$

$$0.12 - 0.88 = -E^\circ$$



$$-0.76 = -E^\circ$$

$$E^\circ = 0.76 \text{ V}$$

50. The compound that **CANNOT** be obtained from the aldol condensation reaction shown below, is



Answer (4)



Therefore, compound 2 cannot be obtained by aldol condensation of given compounds.

51. For a salt **XY**, which is a strong electrolyte, the plot of Λ_m versus \sqrt{c} has a slope of $-90.0 \text{ S cm}^2 \text{ mol}^{-3/2} \text{ L}^{1/2}$ at 298 K. At 0.01 M concentration of **XY**, the value of Λ_m is $145.0 \text{ S cm}^2 \text{ mol}^{-1}$. The limiting molar conductivity of Y^- ion ($\lambda_{\text{Y}^-}^0$, in $\text{S cm}^2 \text{ mol}^{-1}$) at 298 K will be

(Given : $\lambda_{\text{X}^+}^0 = 74.0 \text{ S cm}^2 \text{ mol}^{-1}$)

- (1) 100.0 (2) 90.0
 (3) 76.0 (4) 80.0

Answer (4)

Sol. $y = mx + c$

slope = -90

According to $\lambda_m = \lambda_m^0 - A\sqrt{c}$

$$145 = \lambda_m^0 - 90 \times 0.1$$

$$\lambda_m^0 = 154 \text{ S cm}^2 \text{ mol}^{-1}$$

$$\lambda_{m(\text{XY})}^0 = \lambda_{\text{X}^+}^0 + \lambda_{\text{Y}^-}^0$$

$$154 = 74 + \lambda_{\text{Y}^-}^0$$

$$\lambda_{\text{Y}^-}^0 = 80 \text{ S cm}^2 \text{ mol}^{-1}$$

52. Given below are two statements: One is labelled as **Assertion A** and the other is labelled as **Reason R**.

Assertion A: The first ionization enthalpy of O is lower than that of N and F.

Reason R: The loss of an electron from O leads to stable half-filled p orbital.

In light of the above statements, choose the **most appropriate** answer from the options given below:

- (1) Both **A** and **R** are correct and **R** is **NOT** the correct explanation of **A**
 (2) **A** is correct but **R** is not correct.
 (3) **A** is not correct but **R** is correct
 (4) Both **A** and **R** are correct and **R** is the correct explanation of **A**

Answer (4)

Sol. N = $1s^2 2s^2 2p^3$

F : $1s^2 2s^2 2p^5$


First ionization enthalpy ($\Delta_i H$)

($\Delta_i H$) for N = 1402 kJ mol^{-1}

$\Delta_i H$ for O = 1314 kJ mol^{-1}

$\Delta_i H$ for F = 1681 kJ/mol


55. For the following reaction sequence, choose the correct option




- (1) P and Q are aromatic compounds
- (2) If P gives a carboxylic acid on acidification, Q gives a poisonous gas on exposure to air and light
- (3) Both P and Q are carbonyl compounds
- (4) If P is the sodium salt of a carboxylic acid, Q is a primary alcohol

Answer (2)

Sol.



56. For an elementary chemical reaction, the Arrhenius plot is given below.



If the energy of activation is 6.64 kJ mol^{-1} and $R = 8.3 \text{ J K}^{-1} \text{ mol}^{-1}$, the temperature at which the rate constant becomes $e^2 \text{ min}^{-1}$, is

- (1) 150 K
- (2) 200 K
- (3) 250 K
- (4) 125 K

Answer (2)

Sol. $\ln k = \ln A - \frac{E_a}{RT}$

$$\ln(e^2) = 6 - \frac{6.64 \times 10^3 \text{ J mol}^{-1}}{8.3 \text{ J K}^{-1} \text{ mol}^{-1} \times T}$$

$$2 = 6 - \frac{6.64 \times 10^3}{8.3 \times T}$$

$$T = \frac{6.64 \times 10^3}{8.3 \times 4} = 200 \text{ K}$$

57. Arrange the following compounds in the increasing order of polarity


- A. $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$
- B. $\text{CH}_3\text{CH}_2\text{OH}$
- C. CH_3COCH_3
- D. CH_3COOH

Choose the correct answer from the options given below.


- (1) $C < A < D < B$
- (2) $C < A < B < D$
- (3) $A < C < B < D$
- (4) $A < B < C < D$

Answer (3)

Sol. On the basis of polarity the correct order is



So correct order is



58. A protein undergoes reversible thermal denaturation from its initial state **N** to denatured state **D** according to $\text{N} \rightleftharpoons \text{D}$. At 60°C , the concentrations of both **N** and **D** are equal at equilibrium, and the standard enthalpy change of denaturation is 666 kJ mol^{-1} . The standard entropy change (ΔS° in $\text{kJ K}^{-1} \text{ mol}^{-1}$) of the protein upon denaturation at 60°C is closest to

- (1) 2000.0
- (2) 333.0
- (3) 11.1
- (4) 2.0

Answer (4)

Sol. $\Delta S = \frac{\Delta H}{T}$

$$= \frac{666}{333}$$

$$= 2 \text{ kJ K}^{-1} \text{ mol}^{-1}$$

59. Consider the following statements about the solutions formed by mixing two liquids.
- An ideal solution thus formed obeys Raoult's law throughout the composition range.
 - Mixture of chloroform and acetone shows negative deviation from Raoult's law.
 - Mixture of aniline and phenol shows positive deviation from Raoult's law.
- B and C only
 - A only
 - A and C only
 - A and B only

Answer (4)

Sol. In the case of phenol and aniline solution, the intermolecular hydrogen bonding between phenolic proton and lone pair on nitrogen atom of aniline is stronger than the respective intermolecular hydrogen bonding between similar molecules. Therefore it shows negative deviation.

60. Given below are two statements :

Statement-I : Heating NaCl with concentrated H_2SO_4 and MnO_2 results in oxidation of Mn.


Statement-II : Heating NaI with concentrated H_2SO_4 and MnO_2 results in reduction of Mn.

In light of the above statements, choose the **most appropriate** answer from the options given below.

- Both **Statement-I** and **Statement-II** are incorrect
- Statement-I** is correct but **Statement-II** is incorrect
- Statement-I** is incorrect but **Statement-II** is correct
- Both **Statement-I** and **Statement-II** are correct


Answer (3)

Sol. Statement-I : (False)



So, reduction of Mn is taking place.


Statement-II : (True)




So, Mn has undergone reduction.

61. Given below are two statements :

Statement I : *trans*-But-2-ene upon treatment with Br_2 in CCl_4 gives the following product.




Statement II : *cis*-But-2-ene upon treatment with alkaline KMnO_4 gives the following product.




In the light of the above statements, choose the **most appropriate** answer from the options given below.

- (1) Both **Statement I** and **Statement II** are incorrect
- (2) **Statement I** is correct but **Statement II** is incorrect
- (3) **Statement I** is incorrect but **Statement II** is correct
- (4) Both **Statement I** and **Statement II** are correct

Answer (3)



Therefore, statement-I is incorrect.



62. The highest occupied molecular orbital for Ne_2 is


- (1) σ_{2p}
- (2) π_{2p}^*
- (3) σ_{2p}^*
- (4) π_{2p}

Answer (3)

Sol. $\text{Ne}_2 : \sigma 1s^2 < \sigma^* 1s^2 < \sigma 2s^2 < \sigma^* 2s^2 < \sigma 2p_z^2 < (\pi 2p_x^2 = \pi 2p_y^2) < \pi^* 2p_x^2 = \pi^* 2p_y^2 < \sigma^* 2p_z^2$

$\sigma^* 2p_z$ i.e. $\sigma^* 2p$ is highest occupied orbital.


63. The following carbocation is stabilized by the interaction of the empty p orbital with



- (1) empty σ and empty π^* orbitals
 (2) empty σ^* and filled π orbitals
 (3) empty σ^* and empty π^* orbitals
 (4) filled σ and filled π orbitals

Answer (4)

Sol.



Due to resonance it is stabilised by filled π orbital.

Due to hyperconjugation, it is stabilised by filled σ orbitals.

64. The green paramagnetic species formed by heating KMnO_4 at 513 K is

- (1) Mn_3O_4
 (2) MnO
 (3) KO_2
 (4) K_2MnO_4

Answer (4)

Sol. $2\text{KMnO}_4 \rightarrow \text{K}_2\text{MnO}_4 + \text{MnO}_2 + \text{O}_2$


K_2MnO_4 will form on heating KMnO_4 which has green colour.

65. The correct statement is

- (1) Beryllium has three valence orbitals.
 (2) Magnesium has a maximum covalency of four.
 (3) Aluminium has five valence orbitals.
 (4) Boron has a maximum covalency of four.

Answer (4)

- Sol.**
- Aluminium has nine valence orbitals
 - Beryllium has four valence orbitals
 - Magnesium has a maximum covalency of six.
 - Boron has maximum covalency of four



66. The correct statement about peptides and proteins is
- (1) Only the proteins having a quaternary structure are biologically active
 - (2) In β -pleated sheet structures, peptide chains are held together by intermolecular hydrogen bonds
 - (3) In α -helices, the polypeptide chain is twisted into a left-handed screw (helix) through intramolecular hydrogen bonds
 - (4) Tertiary structure of proteins has two or more polypeptide subunits

Answer (2)

Sol. In β -pleated sheet structure, peptide chains are held together by intermolecular hydrogen bonds.

This is only correct statement.

Remaining (1), (3) and (4) are incorrect statements.

67. $2A \xrightarrow{k} B$ is a zero-order reaction, where $k = 1.0 \text{ mol L}^{-1} \text{ min}^{-1}$. If the initial concentration of A is 2 M, then the time taken to complete 75% of the reaction will be
- (1) 0.75 min
 - (2) 1.0 min
 - (3) 2.0 min
 - (4) 1.5 min

Answer (1)

Sol. For zero order reaction

$$-\frac{1}{2} \frac{dA}{dt} = k$$

$$t = \frac{A_0 - A_t}{2k}$$

$$t = \frac{2 - 0.5}{2} = 0.75 \text{ min}$$

68. Given below are two statements: One is labelled as **Assertion A** and the other is labelled as **Reason R**.
Assertion A: Generally, 3d transition metals have high melting points.

Reason R: Involvement of 3d-electrons in addition to 4s-electrons in the interatomic metallic bonding.

In light of the above statements, choose the **most appropriate** answer from the options given below:

- (1) Both **A** and **R** are correct and **R** is **NOT** the correct explanation of **A**
- (2) **A** is correct but **R** is not correct.
- (3) **A** is not correct but **R** is correct
- (4) Both **A** and **R** are correct and **R** is the correct explanation of **A**

Answer (4)

Sol. The melting points of 3d transition metals are generally high. This is attributed to the involvement of greater number of electrons from $(n - 1) d$ in addition to the ns electrons in the interatomic metallic bonding.

w_1, w_2, w_3 and w_4 represent work done (in calories) in the processes 1, 2, 3 and 4, respectively; ΔU_2 and ΔU_4 are changes in the internal energy for the processes 2 and 4, respectively.

[use $R = 2 \text{ cal K}^{-1} \text{ mol}^{-1}$]

The correct option is

(1) $w_2 + w_4 = \Delta U_2 - \Delta U_4$

(2) $w_1 + w_2 = 2T_1 \ln \frac{V_2}{V_1}$

(3) $w_1 + w_2 + w_3 + w_4 = 0$

(4) $w_1 + w_3 = -2T_1 \ln \frac{V_2}{V_1} - 2T_2 \ln \frac{V_4}{V_3}$

Answer (4)

Sol. $w_1 \rightarrow$ isothermal reversible process

$$w_1 = -nRT_1 \ln \frac{V_2}{V_1}$$

$$w_1 = -1 \times R \times T_1 \ln \frac{V_2}{V_1}$$

$$w_3 = -n \times R \times T_2 \ln \frac{V_4}{V_3}$$

$$w_3 = -1 \times R \times T_2 \ln \frac{V_4}{V_3}$$

$$w_1 + w_3 = -nRT_1 \ln \left(\frac{V_2}{V_1} \right) - nRT_2 \ln \left(\frac{V_4}{V_3} \right)$$

$$w_1 + w_3 = -2T_1 \ln \left(\frac{V_2}{V_1} \right) - 2T_2 \ln \left(\frac{V_4}{V_3} \right)$$

75. In an acidic medium, 10 mL of 0.25 M oxalic acid is titrated with KMnO_4 solution. If the volume of KMnO_4 solution required to reach end point is 10 mL, the strength of the KMnO_4 solution is


(1) 0.20 M

(2) 0.25 M

(3) 0.15 M

(4) 0.10 M

Answer (4)



No. of equivalent of $\text{KMnO}_4 = \text{No. of equivalent of } \text{H}_2\text{C}_2\text{O}_4$

$$10 \times 5 \times M = 10 \times 0.25 \times 2$$

$$M = \frac{5}{50}$$


$$M = \frac{1}{10} = 0.10 \text{ M}$$

76. The lanthanide ion having four unpaired electrons is
(Given : Atomic numbers of Ce = 58, Nd = 60, Tb = 65 and Ho = 67)

- (1) Ce^{3+} (2) Tb^{3+}
(3) Ho^{3+} (4) Nd^{3+}

Answer (3)





Sol. The outermost electronic configuration is



77. The correct decreasing order of oxidation state of the underlined atom in each molecule is

- (1) $\underline{\text{N}}_2\text{O}_5 > \underline{\text{Al}}_2\text{O}_3 > \underline{\text{H}}_2\underline{\text{S}}$ (2) $\underline{\text{Pb}}\text{O}_2 > \underline{\text{N}}_2\text{O}_3 > \underline{\text{S}}\text{O}_3$
(3) $\underline{\text{P}}_4\text{O}_6 > \underline{\text{Cl}}_2\text{O}_7 > \underline{\text{Al}}\text{H}_3$ (4) $\underline{\text{P}}_4\text{O}_{10} > \underline{\text{S}}\text{O}_3 > \underline{\text{H}}_2\underline{\text{O}}$

Answer (1)




78. The formula of tetraammineaquachloridocobalt(III) chloride is

- (1) $[\text{Co}(\text{NH}_3)_4]\text{Cl}_3 \times \text{H}_2\text{O}$ (2) $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]\text{Cl}$
(3) $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]\text{Cl}_2$ (4) $[\text{Co}(\text{NH}_3)_4\text{Cl}_2] \times \text{H}_2\text{O}$

Answer (3)


Sol. Tetraammineaquachloridocobalt(III) chloride is $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]\text{Cl}_2$.

79. Consider the following reaction, and choose the correct option.



- (1) Compound **P** can be prepared by treating benzene with anhydrous AlCl_3 and CH_3COCl
(2) On treatment with bromine water, compound **P** gives a white precipitate
(3) Compound **P** is obtained by the hydrogenation of benzoyl chloride with Pd on BaSO_4
(4) On treating compound **P** with saturated NaHCO_3 solution, brisk effervescence is observed

Answer (3)



80. **Assertion A:** For an ideal solution formed by mixing liquids **P** and **Q**, $\Delta_{\text{mix}} H = 0$ and $\Delta_{\text{mix}} V = 0$

Reason R: No interactions occur between **P** and **Q**

In the light of the above statements, choose the **most appropriate** answer from the options given below.

- (1) Both **A** and **R** are correct but **R** is **NOT** the correct explanation of **A**
- (2) **A** is correct but **R** is not correct
- (3) **A** is not correct but **R** is correct
- (4) Both **A** and **R** are correct and **R** is the correct explanation of **A**

Answer (2)

Sol. For ideal solution, $\Delta_{\text{mix}} H = 0$ and $\Delta_{\text{mix}} V = 0$

In ideal solution energy required to break interaction between P-P and Q-Q will be same as the energy released on the interaction of P-Q.

81. Match the vitamins in **List I** with their sources in **List II**

	List I		List II
A.	vitamin A	I.	meat
B.	vitamin B ₁₂	II.	sunflower oil
C.	vitamin E	III.	green leafy vegetables
D.	vitamin K	IV.	carrots

Choose the correct answer from the options given below.

- (1) A-IV, B-I, C-II, D-III
- (2) A-IV, B-II, C-I, D-III
- (3) A-III, B-I, C-IV, D-II
- (4) A-II, B-III, C-IV, D-I

Answer (1)

84. Two moles of an ideal gas undergo free expansion from 10 L to 100 L at 300 K. The values of ΔS_{system} and $\Delta S_{\text{surroundings}}$ are

(R is universal gas constant)

- (1) $\Delta S_{\text{system}} = 4.606 R$; $\Delta S_{\text{surroundings}} = -4.606 R$ (2) $\Delta S_{\text{system}} = 0$; $\Delta S_{\text{surroundings}} = 4.606 R$
 (3) $\Delta S_{\text{system}} = 4.606 R$; $\Delta S_{\text{surroundings}} = 0$ (4) $\Delta S_{\text{system}} = 0$; $\Delta S_{\text{surroundings}} = 0$

Answer (3)

Sol. During free expansion

$$\begin{aligned}\Delta S_{\text{system}} &= nR \ln \frac{V_2}{V_1} \\ &= 2.303 \times 2 \times R \times \log \frac{100}{10} \\ &= 4.606 R \\ \Delta S_{\text{surroundings}} &= 0 \text{ as } q = 0\end{aligned}$$

85. Among the following options, the correct trend in the electron gain enthalpy is

- (1) $\text{Br} > \text{Cl} > \text{F} > \text{I}$ (2) $\text{Cl} > \text{F} > \text{Br} > \text{I}$
 (3) $\text{I} > \text{Br} > \text{Cl} > \text{F}$ (4) $\text{F} > \text{Cl} > \text{Br} > \text{I}$

Answer (2)

Sol.

Group 17 Elements	Electron gain enthalpy (kJ/mol)
F	-328
Cl	-349
Br	-325
I	-295


So, the correct order of electron gain enthalpy is $\text{Cl} > \text{F} > \text{Br} > \text{I}$.

86. The correct order of solubility of the given salts in water at 298 K is

Salt	K_{sp} at 298 K
AgBr	5.0×10^{-13}
$\text{Zn}(\text{OH})_2$	1.0×10^{-15}
Hg_2Cl_2	1.3×10^{-18}


- (1) $\text{AgBr} > \text{Zn}(\text{OH})_2 > \text{Hg}_2\text{Cl}_2$
 (2) $\text{Hg}_2\text{Cl}_2 > \text{AgBr} > \text{Zn}(\text{OH})_2$
 (3) $\text{Zn}(\text{OH})_2 > \text{AgBr} > \text{Hg}_2\text{Cl}_2$
 (4) $\text{Hg}_2\text{Cl}_2 > \text{Zn}(\text{OH})_2 > \text{AgBr}$

Answer (3)




$$K_{sp} = 5 \times 10^{-13} = S^2$$

$$S = \sqrt{5 \times 10^{-13}}, S = 7.07 \times 10^{-7} \text{ mol/l}$$



$$K_{sp} = 10^{-15} = (S)(2S)^2$$

$$4S^3 = 10^{-15}, S = 0.63 \times 10^{-5} = 6.3 \times 10^{-6} \text{ mol/l}$$




$$K_{sp} = 1.3 \times 10^{-18} = 4S^3, S = 0.69 \times 10^{-6} = 6.9 \times 10^{-7} \text{ mol/l}$$

The correct order of solubility is:




87. Consider the following reaction sequences and choose the correct option.



- (1) **K** and **L** are enantiomers
- (2) **M** and **N** are geometrical isomers
- (3) **M** and **N** are stereoisomers
- (4) **K** and **L** are geometrical isomers

Answer (4)

Sol.




88. The amount of carbon dioxide evolved upon complete combustion of 116 g of n-butane is

(Given: atomic mass in amu H = 1, C = 12 and O = 16)

- (1) 322 g (2) 176 g
(3) 362 g (4) 352 g

Answer (4)



\therefore 1 mole C_4H_{10} produces 4 mole CO_2

\therefore 58 g $C_4H_{10} \equiv 4 \times 44$ g CO_2

$$\therefore 116 \text{ g } C_4H_{10} \equiv \frac{4 \times 44}{58} \times 116 = 352 \text{ g}$$

89. The numbers 17.0145 and 21.0235 were rounded to three figures after the decimal point. The resulting numbers, respectively, are

- (1) 17.015 and 21.023
(2) 17.014 and 21.024
(3) 17.015 and 21.024
(4) 17.014 and 21.023

Answer (2)


Sol. If the rightmost digit to be removed is 5, then the preceding number is not changed if it is an even number, but it is increased by one if it is an odd number.

90. In potash alum, the ratio of K^+ and SO_4^{2-} ions is

- (1) 2 : 1 (2) 2 : 3
(3) 3 : 2 (4) 1 : 2

Answer (4)

Sol. Potash alum



No. of K^+ = 2

No. of SO_4^{2-} = 4

$$\text{Ratio } \frac{K^+}{SO_4^{2-}} = \frac{2}{4} = \frac{1}{2}$$